

Ecological site F092XY011WI

Moist Loamy Lowlands

Last updated: 4/09/2020
Accessed: 07/11/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 092X–Superior Lake Plain

The Wisconsin portion of the Superior Lake Plain (MLRA 92) corresponds very closely to the Superior Coastal Plain Ecological Landscape published by Wisconsin Department of Natural Resources (WDNR 2015). The following brief overview of this MLRA is borrowed from that publication. The Superior Coastal Plain is bordered on the north by Lake Superior and on the south by the Northwest Sands, Northwest Lowlands, and North Central Forest Ecological Landscapes. The total land area is approximately 1.2 million acres, which mostly consists of privately-owned forestland. The climate is strongly influenced by Lake Superior, resulting in cooler summers, warmer winters, and greater precipitation compared to more inland locations. The most extensive landform in this ecological landscape is a nearly level plain of lacustrine clays that slopes gently northward toward Lake Superior. The coastal plain is cut by deeply incised stream drainages and interrupted by the comparatively rugged Bayfield Peninsula. During the Late Wisconsin glacial period, this area was covered with the advancing and retreating lobes of Superior and Chippewa. The landscape was rippled with moraines, but they were subdued by deposition of lacustrine materials. As the glaciers receded, glacial lakes riddled the landscape—most notably, Glacial Lake Duluth. The glacier receded eastward, exposing the western Lake Superior Basin. The ice covered the eastern basin, blocking the outlet of the lake, and continued to recede and contribute meltwaters that filled the glacial lake. The deep, red clays were deposited during this period of glacial lakes. The meltwaters from the glacier also contained sands which were deposited along the edge of the glacial lakes as beach deposits. Deep, narrow valleys have since been carved by rivers and streams flowing north into Lake Superior. Historically, the Superior Coastal Plain was almost entirely forested. Various mixtures of eastern white pine (*Pinus strobus*), white spruce (*Picea glauca*), balsam fir (*Abies balsamea*), white birch (*Betula papyrifera*), balsam poplar (*Populus balsamifera*), quaking aspen (*Populus tremuloides*), and northern white-cedar (*Thuja occidentalis*) occurred on the fine-textured glacio-lacustrine deposits bordering much of the Lake Superior coast. Sandy soils, sometimes interlayered with clays, occur in some places. Such areas supported forests dominated by eastern white pine and red pine (*Pinus resinosa*). Eastern white pine was strongly dominant in some areas, according to mid-19th century notes left by surveyors of the federal General Land Office (Finley, R. 1976). Dry-mesic to wet-mesic northern hardwoods or hemlock-hardwood forests were prevalent on the glacial tills of the Bayfield Peninsula. Large peatlands occurred along the Lake Superior shoreline, associated with drowned river mouths.

Classification relationships

Habitat Types of N. Wisconsin (Kotar, 2002): Three sites in this ES key out to *Acer rubrum* – *Abies balsamea* / *Vaccinium* spp. – *Cornus canadensis* [ArAbVCo], three sites key out to *Acer rubrum* – *Abies balsamea* / *Sanicula* spp. [ArAbSn], and two sites key out to *Acer saccharum* / *Sanicula* spp. - *Mitchella repens* [ASnMi]. Biophysical Setting (Landfire, 2014): This ES is mapped as Laurentian – Acadian Northern Hardwoods Forest – Hemlock, Laurentian – Acadian Sub-boreal Mesic Balsam Fir-Spruce Forest, and Laurentian – Acadian – Pine – Hemlock – Hardwood Forest. The ES is not well represented by any of these, but most similar to Sub-Boreal Mesic Balsam Fir - Spruce Forest. WDNR Natural Communities (WDNR, 2015): This ES is not well represented by any of the described natural communities, but bears some resemblance to Northern Wet-Mesic Forest and Boreal Forest. USFS Subregions: Superior-Ashland Clay Plain Subsection (212Ya); May contain small areas of Ewen Dissected Lake Plain Subsection (212Jo), Winegar Moraines Subsection (212Jc), Gogebic-Penokee Iron Range Subsection (212Jb), and NorthShore Highlands Subsection (212Lb)* Major Land Resource Area (MLRA): Superior Lake Plain (92)

Ecological site concept

Moist Loamy Lowlands has a small extent across MLRA 92. These sites have soils with some variability, but the central concept is finer materials throughout or at some depth in the soil profile, upland position, and are somewhat poorly drained, so they remain moist except in dry periods. The soils formed in loamy and silty till, in stratified silty glaciolacustrine deposits, in stratified loamy glaciolacustrine or glaciofluvial deposits, or in a sandy glaciofluvial mantle over the loamy to silty deposits. The soils range from extremely acid to moderately alkaline. Historically, shade tolerant balsam fir and white spruce were best represented tree species on this ES, but scattered

white and red pines were also common. Following early logging, trembling aspen became the dominant forest type, but succession to balsam fir and, to a lesser degree, white spruce and red maple is evident everywhere where seed sources are present. While there is no good record of the degree of red maple representation in the pre European settlement forests, the species is well represented and successfully reproducing today. Ground flora includes ferns, sedges, grasses, raspberries, horsetail, and yellow bead-lily. This ES differs from other loamy ES' based on its drainage class and remaining moist for most of the growing season. It differs from Moist Sandy Lowlands and Moist Clayey Lowlands based on soil texture, parent material, and the height of seasonally high water table. Moist Sandy Lowlands often has a seasonally high water table of 30 to 76 cm, the clayey uplands has a depth of 0 to 30 cm, and this loamy uplands has a depth of 0 to 76 cm. The range in characteristics of Moist Loamy Lowlands is based on its range in parent materials—mainly the presence of a sandy mantle on some sites.

Associated sites

F092XY007WI	Wet Loamy or Clayey Lowlands These sites are poorly drained soils formed in mainly clayey though sometimes loamy glaciofluvial and silty glaciolacustrine sediments. They have a seasonally high water table and remain saturated for much of the growing season, creating hydric conditions. HGM criteria: recharge, Depressional. The loamy sites are often adjacent to Moist Loamy Lowlands, but located on a lower landscape position in the drainage sequence.
F092XY014WI	Loamy Uplands These sites are deep, moderately well to well drained loamy soils. They formed in loamy and silty till, glaciolacustrine, or glaciofluvial deposits. Some sites have a sandy mantle. Many sites have a seasonally high water table, but does not remain saturated for the growing season. Soils range from strongly acid to strongly alkaline, and some sites have carbonates present. These sites are often adjacent to Moist Loamy Lowlands, but located on a higher landscape position in the drainage sequence.
F092XY009WI	Loamy Sandstone Uplands These sites are shallow loamy soils that overly sandstone bluffs along the shore of Lake Superior. They are moderately well drained soils, but have a seasonally high water table. The soils range from strongly acid to neutral. These sites may occur adjacent to Moist Loamy Lowlands, but located on a higher landscape position in the drainage sequence.

Similar sites

F092XY010WI	Moist Sandy Lowlands Moist Sandy Lowlands have a sandy mantle overlying finer glaciofluvial materials. The finer materials can cause episaturation in spring and fall, allowing the site to remain moist for some of the growing season, but does not remain saturated, nor does it have hydric conditions. Sites are characterized by the ArAbVCo Habitat Type. These sites are found in a similar landscape to Moist Loamy Lowlands, but are coarser textured and in a different drainage sequence.
F092XY012WI	Moist Clayey Lowlands These sites are somewhat poorly drained soils with fine textures that formed in clayey deposits. Some sites have a sandy or loamy mantle. The fine materials cause episaturation in spring and fall and remain saturated for extended period, but the water table can reach depths of 152cm during dry periods. Soils range from strongly acid to strongly alkaline. Carbonates present in some soils beginning at 30cm. Sites are characterized by the ArAbVCo Habitat Type. These sites are found in a similar landscape to Moist Loamy Lowlands, but are finer textured and in a different drainage sequence.

Table 1. Dominant plant species

Tree	(1) <i>Acer rubrum</i> (2) <i>Abies balsamea</i>
Shrub	(1) <i>Corylus cornuta</i> (2) <i>Alnus incana ssp. rugosa</i>
Herbaceous	(1) <i>Pteridium aquilinum</i> (2) <i>Eurybia macrophylla</i>

Physiographic features

This site occurs on plains, drainageways, terraces, and footslopes located on till plains, lake plains, and outwash plains. Landform shape is predominantly linear, but can be concave on footslopes. Elevation of the landforms range from 185 to 330 meters above sea level. Slopes are 0 to 4 percent. This site occurs on all slope aspects.

Table 2. Representative physiographic features

Landforms	(1) Till plain > Drainageway (2) Till plain > Terrace (3) Lake plain (4) Outwash plain
Runoff class	Medium to high
Elevation	190 – 330 m
Slope	0 %
Water table depth	80 cm
Aspect	Aspect is not a significant factor

Climatic features

The Moist Loamy Lowlands PESD is located throughout the MLRA, but most concentrated along the southern extent with some sites extending up into the Bayfield Peninsula. The annual average precipitation ranges from 29-34 inches, with a range of 56-167 inches of annual average snowfall (PRISM, 1981-2010). The annual average minimum temperature ranges from 28-33oF, and the maximum temperature ranges from 47-52oF (PRISM, 1981-2010). The length of the freeze free period ranges from 156 to 194 days, with an average of 171 days (Table 2). The length of the frost-free period ranges from 130 to 166 days, with an average of 144 days (Table 2).

Table 3 Representative climatic features

Frost-free period (characteristic range)	80-110 days
Freeze-free period (characteristic range)	120-140 days

Precipitation total (characteristic range)	790-810 mm
Frost-free period (actual range)	70-110 days
Freeze-free period (actual range)	110-140 days
Precipitation total (actual range)	760-840 mm
Frost-free period (average)	100 days
Freeze-free period (average)	130 days
Precipitation total (average)	790 mm

- (1) ASHLAND 3S [USC00470347], Ashland, WI
- (2) ASHLAND KENNEDY MEM AP [USW00094929], Ashland, WI
- (3) ASHLAND EXP FARM [USC00470349], Ashland, WI
- (4) BAYFIELD 6 N [USC00470603], Bayfield, WI
- (5) FOXBORO [USC00472889], Foxboro, WI
- (6) MADELINE ISLAND [USC00474953], La Pointe, WI

Influencing water features

Water is received primarily through precipitation, from runoff from adjacent uplands, and from ground water. Water is lost from the site primarily through runoff, evapotranspiration, and ground water recharge.

Permeability of the soil is slow to moderately slow. Runoff potential is medium to high. The hydrologic group of this site is either C or D.

Enough water will percolate into the soil resulting in a perched seasonally high water table (episaturation), or an apparent seasonally high water table (endosaturation), at a depth of 0 to 76 cm during spring, but will range to greater than 152 cm under dry conditions. Water that percolates into the soil is generally lost through plant uptake and evapotranspiration. There is potential for significant ground water recharge.

Soil features

The soils of this site are represented by the Brimley, Iosco, Oronto, Robago, and Spear soil series. These soils are classified as Glossaqualfs (Oronto), Endoaquods (Brimley, Iosco, and Robago), and Glossudalfs (Spear).

This ecological site is characterized by very deep, somewhat poorly drained soils formed in loamy and silty till (Oronto), in stratified silty glaciolacustrine deposits (Spear), in stratified loamy glaciolacustrine or glaciofluvial deposits (Brimley and Robago), or in a sandy glaciofluvial mantle over the loamy to silty deposits (Iosco).

The average gravel content within the soil can be as much as 7 percent, while the average content of cobbles and stones can be as much as 2 percent. Soil reaction (pH) in the upper 100 cm ranges from extremely acid to moderately alkaline. Carbonates occur as shallow as 56 cm, but may be absent within 200 cm.

Table 4. Representative soil features

Parent material	(1) Glaciolacustrine deposits (2) Glaciofluvial deposits
Surface texture	(1) Silty clay loam (2) Silt loam (3) Fine sandy loam (4) Loamy sand
Drainage class	Somewhat poorly drained
Permeability class	Slow to moderately slow
Soil depth	200 cm
Available water capacity (0-152.4cm)	10.62 – 19.25 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 10 %
Soil reaction (1:1 water) (0-101.6cm)	4.4 – 8.1
Subsurface fragment volume ≤3" (0-101.6cm)	0 – 10 %
Subsurface fragment volume >3" (0-101.6cm)	Not specified

Ecological dynamics

Because of relatively poorly drained soils historic fire disturbance has likely been less frequent and less severe than on the better drained sites. This is evident by the presence (historic and current) of shade-tolerant and fire sensitive species such as red maple, balsam fir and white spruce. Aspen stands are common in current communities, but they are largely the result of fires associated with past logging. Red maple and balsam fir are the most obvious succeeding species, but white pine and white spruce may also become more important in the future as seed source availability increases. Although the shade-tolerant sugar maple occurs sporadically in some stands its competitive ability is reduced by excessive soil moisture and relatively low nutrient availability. For these reasons it is likely to remain only as a sporadic associate rather than the dominant component of mature forest communities as is typically the case on all mesic sites throughout northern Wisconsin.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Other references

Cleland, D.T.; Avers, P.E.; McNab, W.H.; Jensen, M.E.; Bailey, R.G., King, T.; Russell, W.E. 1997. National Hierarchical Framework of Ecological Units. Published in, Boyce, M. S.; Haney, A., ed. 1997. Ecosystem Management Applications for Sustainable Forest and Wildlife Resources. Yale University Press, New Haven, CT. pp. 181-200.

County Soil Surveys from Douglas, Bayfield, and Ashland Counties.

Curtis, J.T. 1959. Vegetation of Wisconsin: an ordination of plant communities. University of Wisconsin Press, Madison. 657 pp.

Davis, R.B. 2016. Bogs and Fens, A Guide to the Peatland Plants of Northeastern United States and Adjacent Canada. University Press of New England, Hanover and London. 296 pp.

Finley, R. 1976. Original vegetation of Wisconsin. Map compiled from U.S. General Land Office notes. U.S. Forest Service, North Central Forest Experiment Station, St. Paul, Minnesota.

Hvizdak, David. Personal knowledge and field experience.

Jahnke, J. and Gienckce, A. 2002. MLRA 92 Clay Till Field Investigations. Summary of field day investigations by Region 10 Soil Data Quality Specialists.

NatureServe. 2018. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. U.S.A. Data current as of 28 August 2018.

Kotar, J. 1986. Soil – Habitat Type relationships in Michigan and Wisconsin. J. For. and Water Cons. 41(5): 348-350.

Kotar, J., J.A. Kovach and G. Brand. 1999. Analysis of the 1996 Wisconsin Forest Statistics by Habitat Type. U.S.D.A. For. Serv. N.C. Res. Stn. Gen. Tech. Rept. NC-207.

Kotar, J., J. A. Kovach, and T. L. Burger. 2002. A Guide to Forest Communities and Habitat Types of Northern Wisconsin. Second edition. University of Wisconsin-Madison, Department of Forest Ecology and Management, Madison.

Kotar, J., and T. L. Burger. 2017. Wetland Forest Habitat Type Classification System for Northern Wisconsin: A Guide for Land Managers and landowners. Wisconsin Department of Natural Resources, PUB-FR-627 2017, Madison.

Martin, L. 1965. The physical geography of Wisconsin. Third edition. The University of Wisconsin Press, Madison.

McNab, W.H. and P.W. Avers. 1994. Ecological Subregions of the United States: Section Descriptions. USDA For. Serv. Pub. WO-WSA-5, Washington, D.C.

Radeloff, V.C., D.J. Mladenoff, H.S. He and M.S. Boyce. 1999. Forest landscape change in Northwestern Wisconsin Pine Barrens from pre-European settlement to the present. Can. J. For. Res. 29: 1649-1659.

Schulte, L.A., and D.J. Mladenoff. 2001. The original U.S. public land survey records: their use and limitations in reconstructing pre-European settlement vegetation. Journal of Forestry 99:5–10.

Schulte, L.A., and D.J. Mladenoff. 2005. Severe wind and fire regimes in northern forests: historical variability at the regional scale. Ecology 86(2):431–445.

Soil Survey Staff. Input based on personal experience. Tim Miland, Scott Eversoll, Ryan Bevernitz, and Jason Nemecek.

Stearns, F. W. 1949. Ninety years change in a northern hardwood forest in Wisconsin. Ecology, 30: 350-58.

United States Department of Agriculture, Forest Service. 1989. Proceedings – Land Classification Based on Vegetation: Applications for Management. Gen. Tech. Report INT-527.

United States Department of Agriculture, Forest Service. 1990. Silvics of North America, Vol. 1, Hardwoods. Agricultural Handbook 654, Washington, D.C.

United States Department of Agriculture, Forest Service. 1990. Silvics of North America, Vol. 2, Conifers. Agricultural Handbook 654, Washington, D.C.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

United States Department of Agriculture, Natural Resources Conservation Service. 2008. Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service. Technical Note No. 190-8-76. Washington D.C.

Wilde, S.A. 1933. The relation of soil and forest vegetation of the Lake States Region. Ecology 14: 94-105.

Wilde, S.A. 1976. Woodlands of Wisconsin. University of Wisconsin Cooperative Extension, Pub. G2780, 150 pp.

Wisconsin Department of Natural Resources. 2015. The ecological landscapes of Wisconsin: An assessment of ecological resources and a guide to planning sustainable management. Wisconsin Department of Natural Resources, PUB-SS-1131 2015, Madison.

Contributors

Jacob Prater, Associate Professor at University of Wisconsin Stevens Point

John Kotar, Ecological Specialist, independent contract

Bryant Scharenbroch, Assistant Professor at University of Wisconsin Stevens Point

Approval

Chris Tecklenburg, 4/09/2020

Acknowledgments

Contact for Lead Authors: Jacob Prater (jprater@uwsp.edu) Associate Professor at University of Wisconsin Stevens Point, John Kotar (jkotar@wsic.edu) Ecological Specialist, independent contract, and Bryant Scharenbroch Assistant Professor at University of Wisconsin Stevens Point

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/11/2026
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
