

Ecological site F093BY007MI

Sandy Uplands

Last updated: 9/27/2023

Accessed: 08/28/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 093B–Superior Stony and Rocky Loamy Plains and Hills

The Wisconsin portion of this MLRA is a mixture of high-relief moraines and flat till plains with interspersed glacial meltwater deposits. It is bordered on the north by glaciolacustrine deposits of Glacial Lake Duluth and on the south by extensive pitted and unpitted outwash plains. The approximate land area is just under 600,000 acres (935 sq miles). The Penokee-Gogebic Iron Range runs through the middle of the Wisconsin portion of this MLRA and into Michigan. The range is a hilly, bedrock-controlled moraine. The bedrock outcropping is composed of igneous and metamorphic materials and was created by inland folding and faulting of the ancient Superior continent when it collided with the Marshfield continent about 1.8 billion years ago (Dott & Attig, 2004). Volcanic and intrusive bedrock occurs in some places. This bedrock is overlain by a thin layer of glacial till deposited by the Chippewa Lobe. To the north of the range is a former spillway for Glacial Lake Ontonagon. The flowing meltwater cut deep channels into the morainal systems. Glaciofluvial landforms here include old beaches and dunes. South of the range, along the southern edge of this MLRA, are rolling collapsed end moraines, pushed to their extent by the Chippewa and Ontonagon Lobes. The landscape is dotted with abundant kettle lakes and swamps, especially in the eastern portion. Ice-walled lake plains and eskers are also found along these collapsed moraines. The climate is influenced by Lake Superior in areas near the lake, resulting in cooler summers, warmer winters, and greater precipitation – especially snowfall – compared to more inland locations. Historically, mixtures of eastern hemlock (*Tsuga canadensis*), sugar maple (*Acer saccharum*), yellow birch (*Betula alleghaniensis*), eastern white pine (*Pinus strobus*), and red pine (*Pinus resinosa*) covered the area. In wetter pockets (such as the swamps that dot the moraines to the south) white cedar (*Thuja occidentalis*), black spruce (*Picea mariana*), and tamarack (*Larix laricina*) were common (Finley, R., 1976).

Classification relationships

Relationship to Established Frameworks and Classification Systems: Habitat Types of N. Wisconsin (Kotar, 2002): Two sites key out to *Acer saccharum* – *Tsuga canadensis*/ *Maianthemum canadense* (ATM), though one is difficult to key and may also be represented by *Pinus strobus* – *Acer rubrum*/ *Vaccinium angustifolium* (PARV), and one site keys to *Acer saccharum*/ *Vaccinium angustifolium* – *Viburnum acerifolium* (AVVb). Biophysical Setting (Landfire, 2014): This ES is mapped as Laurentian-Acadian Northern Pine(-Oak) Forest, Laurentian-Acadian Northern Hardwoods Forest – Hemlock, and Laurentian-Acadian Northern Hardwoods Forest; though, it is likely best represented by the latter. WDNR Natural Communities (WDNR (2015): This ES is most similar to the Northern Dry-mesic, Mesic, and Boreal Forests. Hierarchical Framework Relationships: Major Land Resource Area (MLRA): Superior Stony and Rocky Loamy Plains and Hills, Eastern Part (93B) USFS Subregions: Winegar Moraines (212Jc) Small sections occur in the Gogebic-Penokee Iron Range (212Jb) subregion Wisconsin DNR Ecological Landscapes: North Central Forest

Ecological site concept

The Sandy Uplands ecological site found on outwash, till, and lake plains, moraines, hillslopes, and stream terraces. These sites are characterized by very deep, moderately well or well drained soils that formed in sandy glaciofluvial, alluvium, and outwash deposits. Precipitation and runoff from adjacent uplands are the primary sources of water. Soils range from very strongly acid to moderately acid. Sandy Uplands lacks a layer of significant clay accumulation like those found in Alfic Sandy Uplands. These sites have a lower available water capacity than their Alfic counterparts, which limits the plant communities they can support. Other upland sites have loamy materials. Sandy materials often have a lower pH and available water capacity than loamy materials, which may limit vegetative growth.

Associated sites

F093BY004MI	Wet Lowlands Wet Lowlands occur on depressions and drainageways and form in loamy till or loamy alluvium underlain by dense sandy till or sandy and gravelly outwash. These sites are poorly drained and occur lower on the drainage sequence than Sandy Uplands.
F093BY005MI	Moist Lowlands Moist Lowlands occur on footslope positions across the landscape. They are not subject to flooding nor ponding. Soils form in till, lacustrine deposits, or outwash deposits and may be loamy to sandy. These sites are somewhat poorly drained and occur slightly lower on the drainage sequence than Sandy Uplands.
F093BY011MI	Dry Uplands Dry Uplands are found in the sandiest, most permeable soils on the driest landscape positions. They are very deep and excessively drained. These sites occur slightly higher on the drainage sequence than Sandy Uplands.

Similar sites

F093BY011MI	Dry Uplands Dry Uplands are found in the sandiest, most permeable soils on the driest landscape positions. While their landscape positions and textures are quite similar, the moderately well to somewhat excessively drained Sandy Uplands sometimes support mesic (ie. slightly wetter) vegetative communities than Dry Uplands.
F093BY006MI	Alfic Sandy Uplands Alfic Sandy Uplands occur on upland sites in deep sandy drift deposits where an argillic horizon is either present or forming. These soils are well to somewhat excessively drained. Unlike Sandy Uplands, Alfic Sandy Uplands have a horizon of significant clay accumulation and can hold slightly more water than their Sandy counterparts. The vegetative community reflects this, with more mesic and slightly more nutrient-demanding species present on Alfic Sandy Uplands.
F093BY010MI	Loamy Uplands Loamy Uplands occur on upland sites in loamy till, sometimes with a mantle of loess. Like Sandy Uplands, these soils lack significant clay accumulation and are moderately well to well drained. The soil textures of Loamy Uplands are finer than those of Sandy Uplands. Species present on Loamy Uplands are slightly more nutrient-demanding than those present on Sandy Uplands.

Table 1. Dominant plant species

Tree	(1) <i>Acer saccharum</i> (2) <i>Betula alleghaniensis</i>
Shrub	(1) <i>Acer saccharum</i>
Herbaceous	(1) <i>Pteridium aquilinum</i> (2) <i>Eurybia macrophylla</i>

Physiographic features

These sites formed on moraines, till plains, lake plains, outwash fan and plains, and stream terraces in backslope, shoulder, and summit positions, Slope ranges from 0 to 45 percent.

These sites are subject to neither flooding nor ponding. These sites rarely have evidence of a seasonally high water table within 80 inches. Surface runoff is negligible to medium.

Figure 1. Distribution of Sandy Uplands in the Superior Stoney and Rocky Loamy Plains and Hills, Eastern Part (93B).

Table 2. Representative physiographic features

Landforms	(1) Till plain (2) Hillslope (3) Moraine (4) Lake plain (5) Outwash fan (6) Outwash plain (7) Stream terrace (8) Disintegration moraine
Runoff class	Negligible to medium
Flooding frequency	None
Ponding frequency	None
Elevation	200 – 300 m
Slope	0 – 50 %
Water table depth	60 – 200 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

The continental climate of the Superior Stoney and Rocky Loamy Plains and Hills, Eastern Part MLRA is characterized by long, cold winters and short, warm summers where precipitation exceeds evapotranspiration. Neither average annual precipitation nor average annual minimum and maximum temperatures vary greatly within this MLRA, though the climate of the northern tip is somewhat affected by Lake Superior and receives higher annual precipitation in the form of lake effect snow.

Table 3 Representative climatic features

Frost-free period (characteristic range)	90-110 days
Freeze-free period (characteristic range)	120-150 days
Precipitation total (characteristic range)	810-910 mm
Frost-free period (actual range)	80-120 days
Freeze-free period (actual range)	120-160 days
Precipitation total (actual range)	760-910 mm
Frost-free period (average)	100 days
Freeze-free period (average)	130 days
Precipitation total (average)	860 mm

- (1) MELLENE 4 NE [USC00475286], Mellen, WI
- (2) HURLEY [USC00473800], Ironwood, WI
- (3) TWIN LAKES [USC00208345], Toivola, MI
- (4) MARQUETTE [USW00014838], Marquette, MI

Influencing water features

Water is received through precipitation and runoff from adjacent uplands. Water is lost from the site primarily through runoff, evapotranspiration, and groundwater recharge.

Permeability of the soils is slow to rapid. The hydrologic soil group of these sites is A.

Wetland description

Hydrologic Group: A

Hydrogeomorphic Wetland Classification: None

Cowardin Wetland Classification: None

Soil features

These sites are represented by the Pence, Karlin, Neconish, and Manitowish series. Pence is classified as a Typic Haplorthod. Karlin is classified as an Entic Haplorthod. Neconish and Manitowish are both classified as Oxyaquic Haplorthods.

These soils primarily form in sandy outwash, sometimes with a thin mantle of loamy alluvium or loamy eolian deposits. Soils are very deep and moderately well to somewhat excessively drained. They do not meet hydric soil requirements.

Surface textures are sandy loam to loamy fine sand. Subsurface textures range from coarse sand to fine sandy loam. Subsurface fragments are found on nearly all sites. Soil pH ranges from very strongly acid to moderately acid with values of 4.5 to 6.0. Carbonates are generally absent in these sites.

Table 4. Representative soil features

Parent material	(1) Glaciofluvial deposits (2) Outwash (3) Alluvium
Surface texture	(1) Loamy fine sand (2) Fine sandy loam (3) Sandy loam
Drainage class	Moderately well drained to somewhat excessively drained
Permeability class	Slow to rapid
Soil depth	200 cm
Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (Depth not specified)	10.11 – 12.29 cm
Calcium carbonate equivalent (Depth not specified)	Not specified
Soil reaction (1:1 water) (Depth not specified)	4.5 – 10
Subsurface fragment volume <=3" (Depth not specified)	0 – 20 %
Subsurface fragment volume >3" (Depth not specified)	0 – 10 %

Ecological dynamics

Historically, mature forests on this ecological site were dominated by shade tolerant sugar maple and hemlock, often with an admixture of yellow birch (Wilde, 1933, Finley, 1976). This association was self-maintained with new cohorts of advance regeneration gaining canopy status through gaps formed by small-scale disturbances and natural mortality in the dominant canopy. Scattered large individuals of less

shade tolerant white pine also were common component of mesic hardwood forests. These presumably became established following relatively rare disturbances that included fire (Schulte and Mladenoff, 2005).

Current stands on this ecological site represent the entire array of potential successional stages from pure aspen, or aspen-white birch, stands to sugar maple dominated mixed northern hardwoods stands. Succession to sugar maple dominance is evident everywhere that seed sources are present. However, hemlock regeneration is scarce. In old forests, hemlock finds optimal conditions for germination and seedling establishment on rotten logs, stumps and mounds that normally have warmer surfaces and better moisture retention than the forest floor (USDA, 1990). Most present forest communities lack these conditions.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Sites key out primarily to ATM, but species are not very well represented by this site. Sites are drier than ATM, and are likely best represented by a dry-mesic type like AVVb. STM designed on ATM and AVVb habitat types. (Wisconsin Forest Habitat Type Classification System)

Other references

Cleland, D.T.; Avers, P.E.; McNab, W.H.; Jensen, M.E.; Bailey, R.G., King, T.; Russell, W.E. 1997. National Hierarchical Framework of Ecological Units. Published in, Boyce, M. S.; Haney, A., ed. 1997. Ecosystem Management Applications for Sustainable Forest and Wildlife Resources. Yale University Press, New Haven, CT. pp. 181-200.

Curtis, J.T. 1959. Vegetation of Wisconsin: an ordination of plant communities. University of Wisconsin Press, Madison. 657 pp.

Dott, R. H., & Attig, J. W. 2004. Roadside geology of Wisconsin. pp. 40. Mountain Press Pub.

Finley, R. 1976. Original vegetation of Wisconsin. Map compiled from U.S. General Land Office notes. U.S. Forest Service, North Central Forest Experiment Station, St. Paul, Minnesota.

NatureServe. 2018. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. U.S.A. Data current as of 28 August 2018.

Kotar, J., J. A. Kovach, and T. L. Burger. 2002. A Guide to Forest Communities and Habitat Types of Northern Wisconsin. Second edition. University of Wisconsin-Madison, Department of Forest Ecology and Management, Madison.

Kotar, J., and T. L. Burger. 2017. Wetland Forest Habitat Type Classification System for Northern Wisconsin: A Guide for Land Managers and landowners. Wisconsin Department of Natural Resources, PUB-FR-627 2017, Madison.

Schulte, L.A., and D.J. Mladenoff. 2001. The original U.S. public land survey records: their use and limitations in reconstructing pre-European settlement vegetation. Journal of Forestry 99:5–10.

Schulte, L.A., and D.J. Mladenoff. 2005. Severe wind and fire regimes in northern forests: historical variability at the regional scale. *Ecology* 86(2):431–445.

Schulte, L.A., and D.J. Mladenoff. 2005. Severe wind and fire regimes in northern forests: historical variability at the regional scale. *Ecology* 86(2):431–445.

United States Department of Agriculture, Forest Service. 1990. *Silvics of North America*, Vol. 1, Hardwoods. Agricultural Handbook 654, Washington, D.C.

United States Department of Agriculture, Forest Service. 1990. *Silvics of North America*, Vol. 2, Conifers. Agricultural Handbook 654, Washington, D.C.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

United States Department of Agriculture, Natural Resources Conservation Service. 2008. Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service. Technical Note No. 190-8-76. Washington D.C.

Wilde, S.A. 1933. The relation of soil and forest vegetation of the Lake States Region. *Ecology* 14: 94-105.

Wilde, S.A. 1976. *Woodlands of Wisconsin*. University of Wisconsin Cooperative Extension, Pub. G2780, 150 pp.

Wisconsin Department of Natural Resources. 2015. The ecological landscapes of Wisconsin: An assessment of ecological resources and a guide to planning sustainable management. Wisconsin Department of Natural Resources, PUB-SS-1131 2015, Madison.

Contributors

Jacob Prater (jprater@uwsp.edu) Associate Professor at University of Wisconsin Stevens Point

Joel Gebhard (jgebhard@uwsp.edu) Associate Research Specialist at University of Wisconsin Stevens Point

Bryant Scharenbroch Assistant Professor at University of Wisconsin Stevens Point

John Kotar (jkotar@wsic.edu) Ecological Specialist, independent contractor

Approval

Suzanne Mayne-Kinney, 9/27/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Contact for Lead Authors: Jacob Prater (jprater@uwsp.edu) Associate Professor at University of Wisconsin Stevens Point, Joel Gebhard (jgebhard@uwsp.edu) Associate Research Specialist at University of Wisconsin Stevens Point, Bryant Scharenbroch Assistant Professor at University of Wisconsin Stevens Point, and John Kotar (jkotar@wsic.edu) Ecological Specialist, independent contractor
Contact for lead author	

Date	09/27/2023
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
