

Ecological site R102CY044NE

WET LAND

Last updated: 12/10/2024
Accessed: 08/30/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 102C—Loess Uplands

Most of this area is in the Dissected Till Plains part of the Central Lowland Province of the Interior Plains. This MLRA has broad, undulating to rolling ridgetops and hilly to steep valley sides. The valleys are generally narrow, but broad flood plains and terraces are along the major rivers and the large tributaries. Elevation ranges from 335 to 610 meters (1,100 to 2,000 feet) increasing from southeast to northwest. Peorian age loess covers most of the area with depths ranging from 2 to 20 meters (6 to 70 feet). Glacial till underlies the loess in most areas. Bedrock can be found at or near the surface predominantly along the Missouri River valley found on the eastern side of the MLRA, but some bedrock can also be found in the northern part of 102C in Minnesota and South Dakota. The soils are predominantly Mollisols but Entisols are prominent in the floodplains of the area. Nearly all the area is farmed with 70% of the area being used as cropland for corn and soybeans. Feed grains and hay crops are also grown. The major resource concerns are wind erosion, water erosion, maintenance of organic matter and soil tilth, and soil moisture management. (USDA/NRCS 2006)

Classification relationships

NE Natural Heritage Program/NE Game & Parks Commission: "Lowland Tallgrass Prairie" General information for MLRA 102C: *Fenneman (1916) Physiographic Regions* Division - Interior Plains East: Province - Central Lowland Section - Till Plains West: Province - Great Plains Section - High Plains *USFS (2007) Ecoregions* Domain - Humid Temperate Division - Prairie Province - Prairie Parkland (Temperate) Section - North-Central Glaciated Plains (251B) *EPA Ecoregions (Omernik 1997)* I - Great Plains (9) II - Temperate Prairies (9.2) III - Western Corn Belt Plains (9.2.3) IV - Loess Prairies (47a) IV - Northeastern Nebraska Loess Hills (47k) IV - Transitional Sandy Plain (47l)

Ecological site concept

This ecological site is a wetland with a seasonal high water table, poorly to very poorly drained soils and a reference community of sedges and rushes. The foremost diagnostic feature of this site is the seasonally high-water table from 0 to 51 centimeters (0 to 20 inches), with additional moisture received from higher adjacent areas as run-on. The high-water table and potential frequent ponding can limit the types of plants that can occupy these areas. This site mainly occurs on nearly level to gently sloping floodplains and stream terraces (0-6% slopes). A few areas are on upland drains or interdunal areas in uplands that can range up to 5% slopes. It usually receives runoff from adjacent sites, has a seasonally high-water table from November-May, may flood occasionally for a brief duration and may frequently pond to four inches. These are predominantly very deep, very poorly to poorly drained soils. The surface texture covers a wide range of silty, loamy, and sandy textures from 15 to 76 centimeters (6 to 30 inches) and the Subsurface Texture Groups are Fine, Fine-Loamy, Fine-Silty or Sandy from 30 to 203 centimeters (12 to 80 inches). The reference plant community consists of 80-95% grasses and grass-like, 5-10% forbs and 0-5% shrubs. Dominant grass-like include Hardstem bulrush, Broadleaf arrowhead and various other members of the sedge family (sedges, rushes, bulrushes and spikesedges).

Associated sites

R102CY046NE	Subirrigated Occurs on higher relief with a seasonally high-water table between 24 and 40 inches and a marked decrease in hydrophytic vegetation.
-------------	---

R102CY048NE	Loamy Overflow Occurs on surrounding higher areas.
R102CY045NE	WET SUBIRRIGATED Occurs on adjacent slightly higher area with seasonally high-water table between 0 and 20 inches with a slight decrease in hydrophytic vegetation.

Similar sites

R102CY043NE	SHALLOW MARSH Occurs most commonly on closed upland depressions. Similar vegetation and water tables / ponding.
--------------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Schoenoplectus acutus</i> (2) <i>Sagittaria latifolia</i>

Physiographic features

This site mainly occurs on nearly level to gently sloping floodplains and stream terraces (0-6% slopes). A few areas are on upland drains or interdunal areas in uplands that can range up to 5% slopes. It usually receives runoff from adjacent sites, has a seasonally high-water table from November-May, may flood occasionally for a brief duration and may pond frequently to a depth of four inches.

Table 2. Representative physiographic features

Landforms	(1) Flood plain (2) Swale (3) Terrace
Runoff class	Very low to low
Flooding duration	Extremely brief (0.1 to 4 hours) to brief (2 to 7 days)
Flooding frequency	None to frequent
Ponding frequency	None to frequent
Elevation	370 – 450 m

Slope	0 – 10 %
Ponding depth	0 – 10 cm
Water table depth	0 – 50 cm
Aspect	Aspect is not a significant factor

Climatic features

Most of the rainfall occurs as high-intensity, convective thunderstorms during the growing season. The maximum precipitation occurs from the middle of spring to early in autumn. Precipitation in winter occurs as snow. The annual snowfall ranges from about 60 centimeters (24 inches) in the southern part of the area to 85 centimeters (34 inches) in the northern part. The average annual temperature gradient trends higher from north (45°F) to south (51°F), and the average annual precipitation gradient trends higher from northwest (25") to southeast (31").

The following data summary includes weather stations representing the full geographic extent of the MLRA and is based on 70% probabilities. This means that actual observed climate conditions may fall outside these ranges 30% of the time. Furthermore, climatic events can manifest many ways. For example, abnormally dry periods could occur as 3 consecutive drought years out of 10, 3 individual years separated by "normal" years, or some combination. Tree-ring records indicate that portions of the Great Plains have also historically experienced droughts lasting several decades, so plant community response will largely depend on the way climatic variability is realized in interaction with past and current land management.

Table 3 Representative climatic features

Frost-free period (characteristic range)	130-130 days
Freeze-free period (characteristic range)	150-150 days
Precipitation total (characteristic range)	690-740 mm
Frost-free period (actual range)	130-130 days
Freeze-free period (actual range)	150-160 days
Precipitation total (actual range)	690-740 mm
Frost-free period (average)	130 days
Freeze-free period (average)	150 days
Precipitation total (average)	710 mm

- (1) YANKTON 2 E [USW00094911], Yankton, SD
- (2) ELGIN [USC00252595], Elgin, NE

- (3) GENOA 2 W [USC00253185], Genoa, NE
- (4) WAYNE [USC00259045], Wayne, NE
- (5) NORFOLK 4W [USC00255997], Norfolk, NE

Influencing water features

This site has a seasonal high-water table from November to May and may flood occasionally for a brief duration.

Soil features

These are predominantly very deep, very poorly to poorly drained soils. The surface texture covers a wide range of silty, loamy, and sandy textures from 15 to 76 centimeters (6 to 30 inches) and the Subsurface Texture Groups are Fine, Fine-Loamy, Fine-Silty or Sandy from 30 to 203 centimeters (12 to 80 inches). Rills, gullies, and water flow patterns are not inherent to this site. Pedestalling is none to slight. Soil aggregate stability will vary based on the varying textural classes in this ecological site.

Major soils assigned to this site include: Calco, Arlo, Barney, Kezan, James, Loup, Zook, Baltic, Norway, Clarno, Clamo, Colo and Lamo.

Table 4. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Silt loam (2) Loam (3) Fine sandy loam (4) Loamy fine sand (5) Silty clay loam
Drainage class	Very poorly drained to poorly drained
Permeability class	Very slow to rapid
Soil depth	200 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (Depth not specified)	2.54 – 22.86 cm
Calcium carbonate equivalent (Depth not specified)	0 – 40 %

Electrical conductivity (Depth not specified)	0 – 10 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0 – 20
Soil reaction (1:1 water) (Depth not specified)	5.6 – 10
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The foremost diagnostic feature of this site is the seasonally high-water table from 0 to 51 centimeters (0 to 20 inches), with additional moisture received from higher adjacent areas as run-on. The high-water table and ponding can limit the types of plants that can occupy these areas but also can “buffer” variability in plant production caused by fluctuating weather conditions (droughts). Relatively minor changes in local elevation can dramatically affect the plant community due to changes in the depth to the water table. This site often occurs in complex with other sites, particularly Wet Land, Subirrigated, and Loamy Overflow. Plant community composition may also experience similar changes through disturbances that affect the water table itself, such as extended dry or wet cycles.

This site developed with occasional fires being part of the ecological processes. It is presumed that the historic fires generally occurred every 3-4 years, were randomly distributed, and started by lightning at various times throughout the season when thunderstorms were likely to occur. It is also believed that pre-European inhabitants may have used fire as a management tool for attracting herds of large migratory herbivores (bison, elk, and/or deer.) The impact of fire over the past 100 years has been relatively insignificant due to the human control of wildfires and the lack of acceptance of prescribed fire as a management tool.

The degree of herbivory (feeding on herbaceous plants) has a significant impact on the dynamics of the site. Historically, periodic grazing by herds of large migratory herbivores was a primary influence. Secondary influences of herbivory by species such as grasshoppers and root feeding organisms impacted the vegetation historically and continue to this day. The management of herbivory by humans through grazing of domestic livestock and/or manipulation of wildlife populations has been a major influence on the ecological dynamics of the site. This management coupled with climate largely dictates the plant communities for the site.

The plant community for this site is dynamic due to the complex interaction of many ecological processes. The interpretive plant community for this site is the reference state. The reference state has been determined by the study of rangeland relic areas, areas protected from excessive disturbance and areas under long term rotational grazing strategies. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a diagram that illustrates the common plant communities that can occur on the site and the transition pathways among communities. The ecological processes will be discussed in more detail in the plant community descriptions following the diagram.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Information presented here has been derived from RANGE-417 archives, Rangeland NRI, and other inventory data. Meetings and comments of local, state and regional experts convene and created the ES concepts and STMs. Field observations from range-trained personnel were also used. Field observations from range-trained personnel were also used. In addition to the multitude of NRCS field office employees and private landowners that helped with site visits and local knowledge, those involved in developing this site include: Nebraska NRCS: Nadine Bishop, State Rangeland Management Specialist Patrick Cowser, Resource Soil Scientist Cassidy Gerdes, Biologist Dirk Schultz, Soil Conservationist Dan Shurtliff, Asst. State Soil Scientist (retired) South Dakota NRCS: Stan Boltz, State Rangeland Management Specialist Shane Deranleau, Area Rangeland Management Specialist Kevin Luebke, State Biologist Iowa NRCS: Jess Jackson, Area Grazing Specialist Minnesota NRCS: Lance Smith, Area Grazing Specialist MLRA Office 10: Stu McFarland, Ecological Site Inventory Specialist Stacey Clark, Ecological Site Inventory Specialist, QA Michael Whited, Region 10 Director (retired) Jo Parsley, Soil Scientist/ Soil Data Quality Specialist National Soil Survey Center: Mike Kucera, National Agronomist, Soil Quality & Ecosystems Steve Peaslee, GIS Specialist, Soil Survey Interpretations Nebraska Game & Parks Commission: Gerry Steinauer, Botanist Scott Wessel, Biologist Russ Hamer, Biologist Rebekah Jessen, Biologist Nebraska Forest Service: Steve Rasmussen, District Forester

Other references

USDA/NRCS 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

USDA/NRCS. 2012. Field Office Technical Guide (Nebraska, Natural Resources Information, Statewide Soil and Site Information, Rangeland Interpretations, Nebraska Range Site Descriptions – Vegetative Zones 3 and 4), U.S. Department of Agriculture, Natural Resources Conservation Service, Nebraska Ecological Sciences.

Fenneman, Nevin M. 1916. Physiographic Subdivision of the United States. Annals of the Association of American Geographers.

Omerik, J.M. 1997. Ecoregions of the Conterminous United States. Annals of the Association of American Geographers, v.77, no. 1, p.118-125.

Cleland_2007, USDA/USFS. 2007. Ecological Subregions: Sections and Subsections for the Conterminous United States. Washington, DC: USDA - Forest Service.

Contributors

Stan Boltz
Dana Larsen
Nadine Bishop
Greg Clark

Approval

Acknowledgments

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident. Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English. To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, available online and at any USDA office, or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632- 9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Nadine Bishop, Emily Helms, Jeff Nichols
Contact for lead author	jeffrey.nichols@usda.gov
Date	12/04/2024
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills: None. Rills are not expected on this site.
2. Presence of water flow patterns: None. Water flow patterns are not expected on this site.
3. Number and height of erosional pedestals or terracettes: None. Pedestals and terracettes are not expected to occur on this site.

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare ground is 5 percent or less. During periods of above average precipitation and run-on, this site may be ponded for longer than normal durations and the typical vegetation will be temporarily reduced. This situation will create areas of bare ground for relatively short periods of time (1 growing season or less) after which early successional forbs, grasses, and grass-like will occupy the site. Bare ground is exposed mineral soil that is not covered by vegetation (basal and/or foliar canopy), litter, standing dead vegetation, gravel/rock, and visible biological crust (e.g., lichen, mosses, algae).

5. Number of gullies and erosion associated with gullies: None. Gullies are not expected on this site.

6. Extent of wind scoured, blowouts and/or depositional areas: None. Wind-scoured and/or depositional areas should not be present.

7. Amount of litter movement (describe size and distance expected to travel): None. Litter falls into place. Litter movement is not expected on this site.

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil stability ratings will be 5 to 6, typically 6. There is typically a high root content and organic matter in the soil surface. Soil surface is very resistant to erosion.

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): The thickness of the A-horizon varies significantly with soil series and ranges from 6 to 50 inches (15 to 127 cm) thick. Soil colors may be very dark gray, dark gray, gray, or grayish brown (10YR 2 to 6/0, 1 or 2 or 2.5Y 3/2 or 5Y 4/1) when dry and black, very dark gray, dark gray, very dark grayish brown (10YR 2 to 4/1 or 2 or 5Y 3/1 or N 2.5/) when moist. Soil structures in the A-horizon are weak fine subangular blocky parting to weak fine granular, weak medium subangular blocky parting to weak fine granular, moderate fine prismatic structure parting to moderate fine subangular blocky weak fine or weak fine and medium granular, or weak thin platy. See Official Soils Descriptions for additional details; major soil series correlated to the site are Calco, Arlo, Barney, Kezan, James, Loup, Zook, Baltic, Norway, Clarno, Clamo, Colo, and Lamo.

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Plant community composition is approximately 80-95 percent grasses or grass-like plants, 5 to 10 percent forbs, and 0 to 5 percent shrubs which optimizes infiltration on the site. The grass and grass-like component is composed of grass-like, cool-season (C3) bunchgrass, warm-season (C4), tallgrass. The functional/structural groups provide a combination of rooting depths and structure which positively influences infiltration. Invasion of introduced cool-season grasses such as reed canarygrass and creeping foxtail may have an adverse impact infiltration and runoff.

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): Typically, none. Physical impact during wet or ponded periods may cause temporary compaction, but this limited compaction will not restrict root development in the reference state.

12. **Functional/Structural Groups** (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Phase 1.1 1. Grass-likes (4 species minimum): hardstem bulrush, sedges, rushes, bulrushes, spikesedges. 2. Native, perennial, cool-season, bunchgrass (3 species minimum): northern reedgrass, bluejoint reedgrass, plains bluegrass. Phase 1.2 1. Native forbs (4 species minimum): water knotweed, nodding beggartick, and other forbs which vary from location to location. 2. Grass-likes (3 species minimum): rushes, sedges, bullrushes, spike-sedges. 3. Native, perennial, cool-season, bunchgrass (2 species minimum): northern reedgrass, bluejoint reedgrass, plains bluegrass.

Sub-dominant: Phase 1.1 1. Native, perennial, warm-season tallgrass (1 species minimum): big bluestem, Indiangrass, prairie cordgrass, switchgrass. 2. Native forbs (4 species minimum): broadleaf arrowhead, and other forbs which vary from location to location. Phase 1.2 1. Native, perennial, warm-season tallgrass (1 species minimum): big bluestem, Indiangrass, prairie cordgrass, switchgrass.

Other: Minor - Phase 1.1 1. Native annual grass: rough barnyard grass, scratchgrass. 2. Shrubs: willows, indigobush, and other shrubs that vary from location to location. Minor - Phase 1.2 1. Native annual grass: rough barnyard grass, scratchgrass. 2. Shrubs: willows, indigobush, and other shrubs that vary from location to location.

Additional: The Tapertip Flatsedge – Rough barnyardgrass – Common Rivergrass – Slough Sedge Community or Reference Community (1.1) includes six F/S groups which include in order of relative abundance, grass-likes; native, perennial, cool-season, bunchgrass; native annual grass; native, perennial, warm-season, tallgrass; native forbs; and shrubs. The Willow - Prairie Cordgrass – Big Bluestem Community (1.2) includes six F/S groups which include in order of relative abundance, native, perennial, warm-season (C4) tallgrass; grass-likes; shrubs; native, perennial, cool-season (C3) bunchgrass; native, annual grasses; and native forbs.

13. **Amount of plant mortality and decadence** (include which functional groups are expected to show mortality or decadence): Bunchgrasses have strong, healthy centers with few (less than 3 percent) dead centers. Shrubs may show some dead branches (less than 5 percent) as plants age.
-

14. **Average percent litter cover (%) and depth (in):** Plant litter cover is evenly distributed throughout the site and is expected to be 80 to 90 percent and at a depth of 0.5 to 1.5 inches (1.25 to 3.75 cm). Reed canarygrass or creeping foxtail excessive litter can negatively impact the functionality of this site.
-

15. **Expected annual annual-production** (this is TOTAL above-ground annual-production, not just forage annual-production): The representative value (RV) for annual production is 4,500 pounds per acre in a year with normal precipitation and temperatures. Low and High production years should yield 3,000 and 6,000 pounds per acre respectively.
-

16. **Potential invasive (including noxious) species** (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: No non-native invasive species are present. Canada thistle, reed canarygrass, creeping foxtail, common watercress, leafy spurge, quackgrass, redtop, and purple loosestrife are known invasives that have the potential to become dominant or co-dominant on the site. Note: species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants.
-

- 17. Perennial plant reproductive capability:** All perennial species exhibit high vigor relative to climatic conditions. Perennial grasses should have vigorous rhizomes or tillers; vegetative and reproductive structures are not stunted. All perennial species should be capable of reproducing annually.
-