

Ecological site R107XA202IA

Calcareous Till Upland Prairie

Last updated: 5/21/2020
Accessed: 09/25/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 107X—Iowa and Missouri Deep Loess Hills

The Iowa and Minnesota Loess Hills (MLRA 107A) includes the Northwest Iowa Plains, Inner Coteau, and Coteau Moraines landforms (Prior 1991; MDNR 2005). It spans two states (Iowa, 89 percent; Minnesota, 11 percent), encompassing approximately 4,470 square miles (Figure 1). The elevation ranges from approximately 1,700 feet above sea level (ASL) on the highest ridges to about 1,115 feet ASL in the lowest valleys. Local relief is mainly 10 to 100 feet. However, some valley floors can range from 80 to 200 feet, while some upland flats only range between 3 and 6 feet. The eastern half of the MLRA is underlain by Wisconsin-age till, deposited between 20,000 and 30,000 years ago and is known as the Sheldon Creek Formation. The western half is underlain by Pre-Illinoian glacial till, deposited more than 500,000 years ago and has since undergone extensive erosion and dissection. Both surfaces are covered by approximately 4 to 20 feet of loess on the hillslopes, and Holocene alluvium covers the till in the drainageways. Cretaceous bedrock, comprised of sandstone and shale, lies beneath the glacial material (USDA-NRCS 2006). The vegetation in the MLRA has undergone drastic changes over time. Spruce forests dominated the landscape 30,000 to 21,500 years ago. As the last glacial maximum peaked 21,500 to 16,000 years ago, they were replaced with open tundras and parklands. The end of the Pleistocene Epoch saw a warming climate that initially prompted the return of spruce forests, but as the warming continued, spruce trees were replaced by deciduous trees (Baker et al. 1990). Not until approximately 9,000 years ago did the vegetation transition to prairies as climatic conditions continued to warm and subsequently dry. Between 4,000 and 3,000 years ago, oak savannas began intermingling within the prairie landscape, while the more wooded and forested areas maintained a foothold in sheltered areas. This prairie-forest transition ecosystem formed the dominant landscapes until the arrival of European settlers (Baker et al. 1992).

Classification relationships

U.S. Forest Service Ecological Subregions: North Central Glaciated Plains (251B) Section, Outer Coteau des Prairies (251Bb), Northwest Iowa Plains (251Bd) Subsections (Cleland et al. 2007) U.S. EPA Level IV Ecoregion: Loess Prairies (47a) (USEPA 2013) National Vegetation Classification – Ecological Systems: Northern Tallgrass Prairie (CES205.686) (NatureServe 2015) National Vegetation Classification - Plant Associations: *Andropogon gerardii* – *Hesperostipa spartea* – *Sporobolus heterolepis* (CEGL002202) (NatureServe 2015) Biophysical Settings: Northern Tallgrass Prairie (BpS 4214200) (LANDFIRE 2009) Natural Resources Conservation Service – Iowa Plant Community Species List: Prairie, Northern Mesic Tallgrass (USDA-NRCS 2007) Iowa Department of Natural Resources: Blacksoil Tallgrass Prairie (INAI 1984) Minnesota Department of Natural Resources: Ups23a Mesic Prairie (Southern) (MDNR 2005)

Ecological site concept

Calcareous Till Upland Prairies are located within the green areas on the map (Figure 1). They occur on upland summits and shoulders, and the soils are Mollisols, Entisols, and Inceptisols that are well-drained and deep, formed in calcareous glacial till. These fine-loamy soils have organic-rich surfaces with high base saturation and will intermittently dry out for periods during the summer season (MDNR 2005; NatureServe 2015). The historic pre-European settlement vegetation on this site was dominated by herbaceous tallgrass species. Big bluestem (*Andropogon gerardii* Vitman) and little bluestem (*Schizachyrium scoparium* (Michx.) Nash) are the dominant grasses of this ecological site. Other grasses that may occur include Indiangrass (*Sorghastrum nutans* (L.) Nash), porcupinegrass (*Hesperostipa spartea* (Trin.) Barkworth), and prairie dropseed (*Sporobolus heterolepis* (A. Gray) A. Gray) (MDNR 2005). Forbs typical of an undisturbed plant community associated with this ecological site include white prairie clover (*Dalea candida* Michx. ex Willd.), purple prairie clover (*Dalea purpurea* Vent.), and meadow zizia (*Zizia aptera* (A. Gray) Fernald) (Drobney et al. 2001; MDNR 2005). Patches of shrubs can be present and generally include leadplant (*Amorpha canescens* Pursh) and prairie rose (*Rosa arkansana* Porter) (MDNR 2005). Fire and herbivory are the primary disturbance factors that maintain this site, while drought is a secondary factor (MDNR 2005; LANDFIRE 2009).

Associated sites

R107XA204IA	Calcareous Till Protected Backslope Savanna Glacial till on backslopes with north and east-facing aspects including Cornell, Moneta, Steinauer, Steinauer variant, and soils that are moderately deep to carbonates
R107XA201IA	Loess Upland Prairie Loess on uplands including Annieville, Galva, McCreath, Primghar, Primghar variant, Ransom, Sac, Sac variant, and Wilmonton
R107XA205IA	Loamy Sediment Upland Prairie Loamy sediments on uplands including Bolan, Bolan variant, Dickman, Everly, Fostoria, and Ocheyedon
R107XA203IA	Calcareous Till Exposed Backslope Prairie Glacial till on backslopes with south and west-facing aspects including Cornell, Moneta, Steinauer, Steinauer variant, and soils that are moderately deep to carbonates

Similar sites

R107XA201IA	Loess Upland Prairie Loess Upland Prairies are derived from fine-silty loess
R107XA205IA	Loamy Sediment Upland Prairie Loamy Sediment Upland Prairies are derived from fine-loamy sediments
R107XA203IA	Calcareous Till Exposed Backslope Prairie Calcareous Till Exposed Backslope Prairies are derived from the same parent material but occur on slopes greater than 20 percent and have a south and west aspect
R107XA204IA	Calcareous Till Protected Backslope Savanna Calcareous Till Protected Backslope Savannas are derived from the same parent material but occur on slopes greater than 20 percent and have a north and east aspect

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Amorpha canescens</i> (2) <i>Rosa arkansana</i>
Herbaceous	(1) <i>Andropogon gerardii</i> (2) <i>Schizachyrium scoparium</i>

Physiographic features

Calcareous Till Upland Prairies occur on upland summits and shoulders (Figure 2). They are situated on elevations ranging from approximately 1000 to 1801 feet ASL. The site does not experience flooding but rather generates runoff to adjacent, downslope ecological sites.

Figure 1. Figure 1. Location of Calcareous Till Upland Prairie ecological site within MLRA 107A.

Figure 2. Figure 2. Representative block diagram of Calcareous Till Upland Prairie and associated ecological sites.

Table 2. Representative physiographic features

Slope shape across	(1) Convex
Slope shape up-down	(1) Convex
Landforms	(1) Upland
Runoff class	Medium to high
Elevation	310 – 550 m
Slope	10 – 20 %
Water table depth	200 cm
Aspect	Aspect is not a significant factor

Climatic features

The Iowa and Minnesota Loess Hills falls into the hot humid continental climate (Dfa) Köppen-Geiger climate classification (Peel et al. 2007). In winter, dry, cold air masses periodically shift south from Canada. As these air masses collide with humid air, snowfall and rainfall result. In summer, moist, warm air masses from the Gulf of Mexico migrate north, producing significant frontal or convective rains. Occasionally, hot, dry winds originating from the Desert Southwest will stagnate over the region, creating extended droughty periods in the summer from unusually high temperatures. Air masses from the Pacific Ocean can also spread into the region and dominate producing mild, dry weather in the autumn known as Indian Summers (NCDC 2006).

The soil temperature regime of MLRA 107A is classified as mesic, where the mean annual soil temperature is between 46 and 59°F (USDA-NRCS 2006). Temperature and precipitation occur along a north-south gradient, where temperature and precipitation increase the further south one travels. The average freeze-free period of this ecological site is about 162 days, while the frost-free period is about 143 days (Table 2). The majority of the precipitation occurs as rainfall in the form of convective thunderstorms during the growing season. Average annual precipitation is approximately 31 inches, which includes rainfall plus the water equivalent from snowfall. The average annual low and high temperatures are 36 and 58°F, respectively (Table 3).

Climate data and analyses are derived from 30-year averages gathered from four National Oceanic and Atmospheric Administration (NOAA) weather stations contained within the range of this ecological site (Table 4).

Table 3 Representative climatic features

Frost-free period (characteristic range)	130-130 days
--	--------------

Freeze-free period (characteristic range)	140-150 days
Precipitation total (characteristic range)	740-790 mm
Frost-free period (actual range)	130-130 days
Freeze-free period (actual range)	140-150 days
Precipitation total (actual range)	710-840 mm
Frost-free period (average)	130 days
Freeze-free period (average)	150 days
Precipitation total (average)	760 mm

- (1) HOLSTEIN [USC00133909], Holstein, IA
- (2) SIOUX CTR 2 SE [USC00137700], Sioux Center, IA
- (3) HAWARDEN [USC00133718], Hawarden, IA
- (4) LE MARS [USC00134735], Le Mars, IA

Influencing water features

Calcareous Till Upland Prairies are not influenced by wetland or riparian water features. Precipitation is the main source of water for this ecological site. Infiltration is slow (Hydrologic Group C) for undrained soils, and surface runoff is medium to high. Precipitation infiltrates the soil surface and percolates downward through the horizons unimpeded by any restrictive layer. The Dakota bedrock aquifer underlying this ecological site is typically deep and confined, leaving it generally unaffected by recharge (Prior et al. 2003). Surface runoff contributes some water to downslope ecological sites (Figure 5).




Figure 7. Figure 5. Hydrologic cycling in Calcareous Till Upland Prairie ecological site.

Soil features

Soils of Calcareous Till Upland Prairies are in the Mollisols, Entisols, and Inceptisols orders, further classified as Entic Hapludolls, Typic Udorthents, and Typic Eutrudepts with slow infiltration and medium to high runoff potential. The soil series associated with this site includes Moneta and Steinauer (Figure 6). The parent material is calcareous glacial till, and the soils are well-drained and deep. Soil pH classes are slightly to moderately alkaline. No rooting restrictions are noted for the soils of this ecological site (Table 5).

Figure 8. Figure 6. Profile sketches of soil series associated with Calcareous Till Upland Prairie.

Table 4. Representative soil features

Parent material	(1) Till
Family particle size	(1) Fine-loamy
Drainage class	Well drained
Permeability class	Slow
Soil depth	200 cm

Ecological dynamics

The information in this Ecological Site Description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

MLRA 107A is defined by a relatively low relief landscape that experiences lower rainfall amounts and available moisture compared to other MLRAs occurring to the south and east. As a result, prairie vegetation communities dominate the uplands, while forested communities are restricted to medium and large streams (Prior 1991; Eilers and Roosa 1994; MDNR 2017a, b). Calcareous Till Upland Prairies form an aspect of this vegetative continuum. This ecological site occurs on upland summits and shoulders on well-drained fine-loamy soils. Plants characteristic of this ecological site consist of sun-loving, fire-adapted herbaceous vegetation.

Fire and grazing are the dominant ecosystem drivers for maintaining the vegetation of Calcareous Till Upland Prairies. Fire intensity typically consisted of periodic, high severity surface fires occurring every 1 to 5 years (LANDFIRE 2009). Ignition sources included summertime lightning strikes from convective storms and bimodal, human ignitions during the spring and fall seasons. Native Americans regularly set fires to improve sight lines for hunting, driving large game, improving grazing and browsing habitat, agricultural clearing, and enhancing vital ethnobotanical plants (Barrett 1980). Bison (*Bos bison*) and possibly prairie elk (*Cervus elaphus*) were the main herbivores in northern tallgrass prairies. Herbivory occurred via mob grazing with large herds of animals rapidly moving across the prairie as they grazed (LANDFIRE 2009). These continuous disturbances provided critical conditions for perpetuating the native prairie ecosystem (MDNR 2005).

Drought has also played a role in shaping this ecological site. The periodic episodes of reduced soil moisture in conjunction with the well-drained soils have favored the proliferation of plant species tolerant of such conditions. Drought can also slow the growth of plants and result in dieback of certain species. When coupled with fire and herbivory, periods of drought can greatly delay the establishment of woody vegetation (Pyne et al. 1996).

Today, Calcareous Till Upland Prairies are limited in their extent, having been converted to agricultural production land. Corn (*Zea mays* L.) and soybeans (*Glycine max* (L.) Merr.) are the dominant crops grown on this ecological site, but small patches of forage land are present. A return to the historic plant community may not be possible following extensive land modification, but long-term conservation agriculture or prairie reconstruction efforts can help to restore some biotic diversity and ecological function. The state-and-transition model that follows provides a detailed description of each state, community phase, pathway, and transition. This model is based on available experimental research, field observations, literature reviews, professional consensus, and interpretations.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 3.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 13. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 14. Community 4.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

No field plots were available for this site. A review of the scientific literature and professional experience were used to approximate the plant communities for this provisional ecological site. Information for the state-and-transition model was obtained from the same sources. All community phases are considered provisional based on these plots and the sources identified in ecological site description.

Other references

Baker, R.G., C.A. Chumbley, P.M. Witinok, and H.K. Kim. 1990. Holocene vegetational changes in eastern Iowa. *Journal of the Iowa Academy of Science* 97: 167-177.

Baker, R.G., L.J. Maher, C.A. Chumbley, and K.L. Van Zant. 1992. Patterns of Holocene environmental changes in the midwestern United States. *Quaternary Research* 37: 379-389.

Barret, S.W. 1980. Indians and fire. *Western Wildlands Spring*: 17-20.

Bharati, L., K.-H. Lee, T.M. Isenhardt, and R.C. Schultz. 2002. Soil-water infiltration under crops, pasture, and established riparian buffer in Midwestern USA. *Agroforestry Systems* 56: 249-257.

Brudvig, L.A., C.M. Mabry, J.R. Miller, and T.A. Walker. 2007. Evaluation of central North American prairie management based on species diversity, life form, and individual species metrics. *Conservation Biology* 21: 864-874.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.

Drobney, P.D., G.S. Wilhelm, D. Horton, M. Leoschke, D. Lewis, J. Pearson, D. Roosa, and D. Smith. 2001. Floristic Quality Assessment for the State of Iowa. Neal Smith National Wildlife Refuge and Ada Hayden Herbarium, Iowa State University, Ames, IA. 123 pps.

Eilers, L. and D. Roosa. 1994. The Vascular Plants of Iowa: An Annotated Checklist and Natural History. University of Iowa Press, Iowa City, IA. 319 pps.

Fierer, N., J. Ladau, J.C. Clemente, J.W. Leff, S.M. Owens, K.S. Pollard, R. Knight, J.A. Gilbert, and R.L. McCulley. 2013. Reconstructing the microbial diversity and function of pre-agricultural tallgrass prairie soils in the United States. *Science* 342: 621-624.

Franzluebbers, A.J., J.A. Stuedemann, H.H. Schomberg, and S.R. Wilkinson. 2000. Soil organic C and N pools under long-term pasture management in the Southern Piedmont USA. *Soil Biology and Biochemistry* 32:469-478.

Iowa Natural Areas Inventory [INAI]. 1984. An Inventory of Significant Natural Areas in Iowa: Two Year Progress Report of the Iowa natural Areas Inventory. Iowa Natural Areas Inventory, Iowa Department of Natural Resources, Des Moines, IA.

Kardol, P. and D.A. Wardle. 2010. How understanding aboveground-belowground linkages can assist restoration ecology. *Trends in Ecology and Evolution* 25: 670-679.

LANDFIRE. 2009. Biophysical Setting 4214200 Northern Tallgrass Prairie. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.

Leake, J., D. Johnson, D. Donnelly, G. Muckle, L. Boddy, and D. Read. 2004. Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Canadian Journal of Botany* 82: 1016-1045.

Martin, L.M. and B.J. Wilsey. 2006. Assessing grassland restoration success: relative roles of seed additions and native ungulate activities. *Journal of Applied Ecology* 43: 1098-1110.

Martin, L.M. and B.J. Wilsey. 2012. Assembly history alters alpha and beta diversity, exotic-native proportions and functioning of restored prairie plant communities. *Journal of Applied Ecology* 49: 1436-1445.

Minnesota Department of Natural Resources [MDNR]. 2005. Field Guide to the Native Plant Communities of Minnesota: The Prairie Parkland and Tallgrass Aspen Parklands Provinces. Ecological Land Classification Program, Minnesota County Biological Survey, and Natural Heritage and Nongame Research Program. Minnesota Department of Natural Resources, St. Paul, Minnesota.

Minnesota Department of Natural Resources [MDNR]. 2017a. Coteau Moraines Subsection. Minnesota Department of Natural Resources: Ecological Classification System. Available at: <http://www.dnr.state.mn.us/ecs/251Bb/index.html>. (Accessed 10 October 2017).

Minnesota Department of Natural Resources [MDNR]. 2017b. Inner Coteau Subsection. Minnesota Department of Natural Resources: Ecological Classification System. Available at: <http://www.dnr.state.mn.us/ecs/251Bc/index.html>. (Accessed 10 October 2017).

National Climate Data Center [NCDC]. 2006. Climate of Iowa. Central Region Headquarters, Climate Services Branch, National Climatic Data Center, Asheville, NC.

NatureServe. 2015. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1 NatureServe, Arlington, VA. Available at <http://explorer.natureserve.org>. (Accessed 13 February 2017).

Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633-1644.

Prior, J.C. 1991. Landforms of Iowa. University of Iowa Press for the Iowa Department of Natural Resources, Iowa City, IA. 153 pps.

Prior, J.C., J.L. Boekhoff, M.R. Howes, R.D. Libra, and P.E. VanDorpe. 2003. Iowa's Groundwater Basics: A Geological Guide to the Occurrence, Use, & Vulnerability of Iowa's Aquifers. Iowa Department of Natural Resources, Iowa Geological Survey Educational Series 6. 92 pps.

Pyne, S.J., P.L. Andrews, and R.D. Laven. 1996. Introduction to Wildland Fire, Second Edition. John Wiley and Sons, Inc. New York, New York. 808 pps.

Skinner, R.H. 2008. High biomass removal limits carbon sequestration potential of mature temperate pastures. *Journal for Environmental Quality* 37: 1319-1326.

Smith, D.D. 1998. Iowa prairie: original extent and loss, preservation, and recovery attempts. *The Journal of the Iowa Academy of Sciences* 105: 94-108.

Smith, D.D., D. Houseal, and K. Henderson. 2010. The Tallgrass Prairie Center Guide to Prairie Restoration in the Upper Midwest. University of Iowa Press, Iowa City, IA. 338 pps.

Society for Ecological Restoration [SER] Science & Policy Working Group. 2002. SER Primer on Ecological Restoration. Available at: <http://www.ser.org/>. (Accessed 28 February 2017).

Teague, W.R., S.L. Dowhower, S.A. Baker, N. Haile, P.B. DeLaune, and D.M. Conover. 2011. Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems and Environment* 141: 310-322.

Tomer, M.D., D.W. Meek, and L.A. Kramer. 2005. Agricultural practices influence flow regimes of headwater streams in western Iowa. *Journal of Environmental Quality* 34: 1547-1558.

Undersander, D., B. Albert, D. Cosgrove, D. Johnson, and P. Peterson. 2002. Pastures for Profit: A Guide to Rotational Grazing (A3529). University of Wisconsin-Extension and University of Minnesota Extension Service. 43 pps.

United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2003. National Range and Pasture Handbook, Revision 1. Grazing Lands Technology Institute. 214 pps.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. 682 pps.

United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2007. Iowa NRCS Plant Community Species Lists. Des Moines, IA. Available at: https://www.nrcs.usda.gov/wps/portal/nrcs/detail/ia/technical/ecoscience/bio/?cid=nrcs142p2_008160. (Accessed 1 February 2018).

U.S. Environmental Protection Agency [EPA]. 2013. Level III and Level IV Ecoregions of the Continental United States. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1: 3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-andiv-ecoregions-continental-united-states>. (Accessed 1 March 2017).

Williams, D.A., L.L. Jackson, and D.D. Smith. 2007. Effects of frequent mowing on survival and persistence of forbs seeded into a species-poor grassland. *Restoration Ecology* 18: 628-637.

Wilsey, B.J. 2008. Productivity and subordinate species response to dominant grass species and seed source during restoration. Restoration Ecology 18: 628-637.

Contributors

Lisa Kluesner
Dan Pulido

Approval

Chris Tecklenburg, 5/21/2020

Acknowledgments

This project could not have been completed without the dedication and commitment from a variety of partners and staff (Table 6). Team members supported the project by serving on the technical team, assisting with the development of state and community phases of the state-and-transition model, providing peer review and technical editing, and conducting quality control and quality assurance reviews. Table 6. List of primary contributors and reviewers. Organization Name Title Location Drake University: Dr. Tom Rosburg, Professor of Ecology and Botany, Des Moines, IA Iowa Department of Natural Resources: John Pearson, Ecologist, Des Moines, IA LANDFIRE (The Nature Conservancy): Randy Swaty, Ecologist, Evanston, IL Natural Resources Conservation Service: Rick Bednarek, Iowa State Soil Scientist, Des Moines, IA Patrick Chase, Area Resource Soil Scientist, Fort Dodge, IA Stacey Clark, Regional Ecological Site Specialist, St. Paul, MN James Cronin, State Biologist, Des Moines, IA Tonie Endres, Senior Regional Soil Scientist, Indianapolis, IN John Hammerly, Soil Data Quality Specialist, Indianapolis, IN Lisa Kluesner, Ecological Site Specialist, Waverly, IA Sean Kluesner, Earth Team Volunteer, Waverly, IA Jeff Matthias, State Grassland Specialist, Des Moines, IA Louis Moran, PhD, Area Resource Soil Scientist, Sioux City, IA Kevin Norwood, Soil Survey Regional Director, Indianapolis, IN Doug Oelmann, Soil Scientist, Des Moines, IA James Phillips, GIS Specialist, Des Moines, IA Dan Pulido, Soil Survey Leader, Atlantic, IA Jason Steele, Area Resource Soil Scientist, Fairfield, IA Doug Wallace, Ecological Site Specialist, Columbia, MO

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/25/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
