

Ecological site R111XD011IN

Mollic Till Ridge

Last updated: 9/11/2024

Accessed: 09/26/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 111X–Indiana and Ohio Till Plain

A PROVISIONAL ECOLOGICAL SITE is a conceptual grouping of soil map unit components within a Major Land Resource Area (MLRA) based on the similarities in response to management. Although there may be wide variability in the productivity of the soils grouped into a Provisional Site, the soil vegetation interactions as expressed in the State and Transition Model are similar and the management actions required to achieve objectives, whether maintaining the existing ecological state or managing for an alternative state, are similar. Provisional Sites are likely to be refined into more precise group during the process of meeting the APPROVED ECOLOGICAL SITE DESCRIPTION criteria. This PROVISIONAL ECOLOGICAL SITE has been developed to meet the standards established in the National Ecological Site Handbook. The information associated with this ecological site does not meet the Approved Ecological Site Description Standard, but it has been through a Quality Control and Quality Assurance processes to assure consistency and completeness. Further investigations, reviews and correlations are necessary before it becomes an Approved Ecological Site Description. 111D – Indiana and Ohio Till Plain, Western Part. This MLRA occurs in two separate areas. One area is in the west-central part of Indiana (73 percent), and the other is in southwestern Ohio (27 percent). The MLRA makes up 5,355 square miles (13,880 square kilometers). It includes the towns of Crawfordville, Delphi, Frankfort, Lafayette, and Liberty, Indiana, and Hamilton, Lebanon, Middletown, and Wilmington, Ohio. Interstates 65 and 74 cross the part of this area in Indiana, and Interstates 71 and 75 cross the part in Ohio. Shades and Turkey Run State Parks are in the part in Indiana, and Caesar Creek and Hueston Woods State Parks are in the part in Ohio. A small portion of the Wright-Patterson Air Force Base, in Ohio, is in the northern part of the area. This area is in the Till Plains Section of the Central Lowland Province of the Interior Plains. It is dominated by loess hills and flats that are broken in places by moraines, kames, outwash plains, and stream terraces. Narrow, shallow valleys commonly are along the few large streams in the area. Elevation ranges from 530 to 1,050 feet (160 to 320 meters), increasing gradually from southwest to northeast. Relief is mainly a few meters, but in some areas hills rise as much as 100 feet (30 meters) above the adjoining plains. The extent of the major Hydrologic Unit Areas (identified by four-digit numbers) that make up this MLRA is as follows: Wabash (0512), 68 percent; Great Miami (0508), 15 percent; Middle Ohio (0509), 14 percent; Scioto (0506), 2 percent; and Upper Illinois (0712), 1 percent. Wildcat Creek in Indiana and the Little Miami River in Ohio have been designated as National Wild and Scenic Rivers. Sugar Creek and Walnut Creek occur in the part of the area in northern Indiana, and the Whitewater River is in the part in southeastern Indiana. The Sevenmile, Fourmile, and Great Miami Rivers cross the part of the area in Ohio. Most of the eastern part of this MLRA is underlain by Late Ordovician shale and limestone. The western part is underlain by shale, siltstone, sandstone, limestone, and dolostone ranging in age from Middle Pennsylvanian to Silurian. Surficial materials include glacial deposits of till, outwash, and lacustrine sediments from Wisconsin and older glacial periods. A thin or moderately thick mantle of loess overlies much of the area.

Classification relationships

Major Land Resource Area (USDA-Natural Resources Conservation Service, 2006) USFS Ecological Regions (USDA, 2007): Sections –Central Till Plains, Beech Maple (222H), Interior Low Plateau-Shawnee Hills (223D), Interior Low Plateau-Bluegrass (223F), Central Till Plains-Oak Hickory (223G), Central Till Plains and Grand Prairies (251D) Subsections -Bluffton Till Plains (222Ha), Miami-Scioto Plain-Tipton Till Plain (222Hb), Little Miami Old Drift Plain (222Hc), Mad River Interlobate Plains (222Hd), Crawford Uplands (223De), Crawford Escarpment (223Df), Northern Bluegrass (223Fd), Lower Wabash Alluvial Plain (223Gc), Southwest Indiana Glaciated Lowlands (223Ge), Eastern Grand Prairie (253Dd). NatureServe Systems anticipated (NatureServe, 2011): Agriculture - Cultivated Crops and Irrigated Agriculture, Agriculture - Pasture/Hay, Allegheny-Cumberland Dry Oak Forest and Woodland, Central Interior Acidic Cliff and Talus, Central Interior Highlands Calcareous Glade and Barrens, Central Tallgrass Prairie, Clearcut - Grassland/Herbaceous, Introduced Upland Vegetation – Treed, Managed Tree Plantation, Mississippi River Riparian Forest, North-Central Interior and Appalachian Acidic Peatland, North-Central Interior Beech-Maple Forest, North-Central Interior Dry-Mesic Oak Forest and Woodland, North-Central Interior Dry Oak Forest and Woodland, North-Central Interior Floodplain, North-Central Interior Freshwater Marsh, North-Central Interior Maple-Basswood Forest, North-Central Interior Oak Savanna, North-Central Interior Wet Flatwoods, North-Central Interior Wet Meadow-Shrub Swamp, North-Central Oak Barrens, Northern Atlantic Coastal Plain Hardwood Forest, Ruderal Forest, Ruderal Upland - Old Field,

South-Central Interior / Upper Coastal Plain Wet Flatwoods, South-Central Interior Large Floodplain, South-Central Interior Mesophytic Forest, South-Central Interior Small Stream and Riparian, Southern Appalachian Oak Forest, Southern Interior Low Plateau Dry-Mesic Oak Forest, Successional Shrub/Scrub LANDFIRE Biophysical Settings anticipated (USGS, 2010): Allegheny-Cumberland Dry Oak Forest and Woodland, Bluegrass Savanna and Woodland, Central Interior and Appalachian Floodplain Systems, Central Interior and Appalachian Riparian Systems, Central Interior and Appalachian Shrub-Herbaceous Wetland Systems, Central Interior and Appalachian Swamp Systems, Central Interior Highlands Calcareous Glade and Barrens, Central Interior Highlands Dry Acidic Glade and Barrens, Central Tallgrass Prairie, Great Lakes Coastal Marsh Systems, Mississippi River Alluvial Plain Dry-Mesic Loess Slope Forest, North-Central Interior Beech-Maple Forest, North-Central Interior Dry-Mesic Oak Forest and Woodland, North-Central Interior Dry Oak Forest and Woodland, North-Central Interior Maple-Basswood Forest, North-Central Interior Oak Savanna, North-Central Interior Wet Flatwoods, Paleozoic Plateau Bluff and Talus, Pennyroyal Karst Plain Prairie and Barrens, South-Central Interior Mesophytic Forest, South-Central Interior/Upper Coastal Plain Flatwoods, Southern Appalachian Oak Forest, Southern Interior Low Plateau Dry-Mesic Oak Forest

Ecological site concept

This site is an upland site formed on glacial till parent materials located in a swell (convex) landscape position. It is located on summits, shoulders and backslopes. The soil surface color is dark(3/2 Munsell or darker) but does not extend beyond 10 inches, making them taxonomically mollic integrades. The characteristic vegetation of this site is that of an oak savanna. This site exhibits a co-dominance of woody and herbaceous species. Black oak is the most abundant woody species, with white oak also being present as well as a very few hickory trees. The herbaceous vegetation was dominated by little bluestem, big bluestem, switchgrass and strongly resembled the richness and diversity of species found in adjacent, more xeric prairies. Fire frequency and intensity were the principle drivers that maintained the site. Frequent low intensity ground fires kept the tree species from becoming more dominant and allowed the herbaceous species to be co-dominant on the site. Lengthening the fire return interval transitions this site to a woodland or forest state. Due to the productivity of the soils of this site as well as the gentle topography, the majority of this site is currently in agriculture production.

Associated sites

R111XD012IN	Till Ridge Prairie Located on an adjacent landscape position; dark soil surface color extends more than 10 inches
F111XD010IN	Till Ridge Located on an adjacent landscape position; soils surface color is lighter than 3/2 Munsell; soils are moderately well to well drained.
F111XD009IN	Wet Till Ridge Located on an adjacent landscape positon; soils surface color is lighter than 3/2 Munsell; soils are poorly to somewhat poorly drained.

Similar sites

R111XD006IN	Mollic Till Depression A savanna site, but with a greater tree influence. Located on till depressions. More overall production.
R111XD019IN	Outwash Integrate Savanna site with lesser amount of trees. Developed on outwash parent materials. Less overall production.
R111XD027IN	Sand Dune Drier savanna vegetation type due to lighter soils and increased drainage

Table 1. Dominant plant species

Tree	(1) <i>Quercus velutina</i> (2) <i>Quercus alba</i>
Shrub	Not specified
Herbaceous	(1) <i>Andropogon gerardii</i> (2) <i>Schizachyrium scoparium</i>

Physiographic features

This ecosite is found till landscapes in MLRA 111D: Indiana and Ohio Till Plain, Western Part on swells or convex slopes.

Table 2. Representative physiographic features

Landforms	(1) Till plain (2) Ground moraine (3) Recessional moraine
Runoff class	Low to high
Flooding frequency	None
Ponding frequency	None
Elevation	140 – 290 m
Slope	0 – 10 %
Ponding depth	0 cm
Water table depth	10 – 120 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

The average annual precipitation in this area is 36 to 43 inches (915 to 1,090 millimeters). Most of the rainfall occurs as convective thunderstorms during the growing season. About half or more of the precipitation occurs during the freeze-free period. Snowfall is common in winter. The average annual temperature is 49 to 54 degrees F (10 to 12 degrees C). The freeze-free period averages about 200 days and ranges from 180 to 215 days.

Table 3 Representative climatic features

Frost-free period (characteristic range)	130-140 days
Freeze-free period (characteristic range)	170-180 days
Precipitation total (characteristic range)	990-1,090 mm
Frost-free period (actual range)	130-150 days
Freeze-free period (actual range)	170-190 days
Precipitation total (actual range)	990-1,120 mm
Frost-free period (average)	140 days
Freeze-free period (average)	180 days
Precipitation total (average)	1,040 mm

- (1) BOSWELL 4WNW [USC00120858], Fowler, IN
- (2) LAFAYETTE 8 S [USC00124715], Lafayette, IN
- (3) TERRE HAUTE INDIANA ST [USC00128723], Terre Haute, IN
- (4) WABASH [USC00129138], Wabash, IN
- (5) HAMILTON BUTLER CO RGNL AP [USW00053855], Fairfield, OH
- (6) WILMINGTON 3 N [USC00339219], Wilmington, OH
- (7) ATTICA 2E [USC00120331], Attica, IN

Influencing water features

This ecological site is not influenced by wetland or riparian water features.

Soil features

The soil series associated with this site are: Wingate, Toronto, Throckmorton, Octagon, Montmorenci, Mellott, Markham. They are very deep, somewhat poorly drained to well drained, and very slow to moderate permeable soils, with very strongly acidic to slightly alkaline soil reaction, that formed in Glaciofluvial deposits, Loess, Till from Limestone.

Table 4. Representative soil features

Parent material	(1) Till – limestone
Surface texture	(1) Loam (2) Silty clay loam (3) Clay loam

Family particle size	(1) Loamy
Drainage class	Somewhat poorly drained to well drained
Permeability class	Very slow to moderate
Soil depth	70 – 150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	11.18 – 20.57 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 30 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	5.3 – 7.9
Subsurface fragment volume <=3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The historic plant community of the Mollic Till Ridge ecological site is that of a savanna. This site is characterized by the ebb and flow of the co-dominant tall grass prairie species and oak species, primarily black and white oak. This dynamic was driven by the fire frequency and intensity. The canopy tree cover varied from about 10-60% and was dominated by black oak. Other prominent tree species included

white oak and in some cases bur oak. The ground layer is dominated by graminoids principally big bluestem and little bluestem, but also included Indiangrass and switchgrass. Ground fires were the most common disturbance for this site. These frequent, but lower intensity fires, in conjunction with weather related tree mortality maintained the function of the site. Replacement fires occurred approximately every 200 years. Since settlement, the majority of this site has been converted to agricultural use with the majority being row crop agriculture where corn and soybean production is the most common practice.

State and transition model

Figure 7. Mollic Till Ridge

Figure 8. Legend

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Other references

Betz, R. (1973). The prairies of Indiana. Proceedings of the Fifth Midwest Prairie Conference (pp. 34-31). Ames: Iowa State University.

Braun, E. Lucy. 2001. Deciduous forests of eastern North America. Caldwell, N.J.: Blackburn Press.

Homoya, M. A., Abrell, D. B., Aldrich, J. R., & Post, T. W. (1985). The Natural Regions of Indiana. Indiana Academy of Science , 94, 245-269.

Kartesz, J. T. (2011). Density Gradient Map Samples Produced From BONAP's Floristic Synthesis. Retrieved 12 12, 2011, from Biota of North America Program: <http://bonap.org/diversity/diversity/diversity.html>

NatureServe. (2011). An online encyclopedia of life [web application]. NatureServe, Arlington, VA, USA [Online: [www. natureserve. org/explorer](http://www.natureserve.org/explorer)] .

Jackson, Marion T. 1997. The Natural heritage of Indiana. Bloomington: Indiana University Press, published in association with the Indiana Department of Natural Resources and the Indiana Academy of Science.

Johnson, Paul S., Stephen R. Shifley, and Robert Rogers. 2002. The ecology and silviculture of oaks. Wallingford, Oxon: CABI

Transeau, E. (1935). The prairie peninsula. Ecology vol. 16 (3) , 423-437.

Upland Oak Ecology Symposium, and Martin A. Spetich. 2004. Upland Oak Ecology Symposium: history, current conditions, and sustainability : Fayetteville, Arkansas, October 7-10, 2002. [Asheville, NC]: [Southern Research Station].

USDA. (2007). Ecological Subregions: Sections and Subsections for the Conterminous United States. Washington, DC: USDA - Forest Service.

USDA. (2006). Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U. S. Department of Agriculture, Natural Resources Conservation Service. U. S. Department of Agriculture Handbook 296.

USGS. (2010). LANDFIRE Biophysical Settings. Retrieved from <http://www.landfire.gov>

Whitaker, John O., Charles J. Amlaner, Marion T. Jackson, George R. Parker, and Peter Evans Scott. 2012. Habitats and ecological communities of Indiana presettlement to present. Bloomington: Indiana University Press.

Contributors

Tyler Staggs

Approval

Greg Schmidt, 9/11/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	TYLER STAGGS
Contact for lead author	
Date	02/01/2022
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. Perennial plant reproductive capability:
-