

Ecological site F113XY913IL

Clayey Till Backslope Woodland

Last updated: 5/17/2024
Accessed: 08/31/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 113X–Central Claypan Areas

The eastern Illinois portion of the Central Claypan Areas MLRA is in the Till Plains Section of the Central Lowland Province of the Interior Plains (USDA-NRCS, 2006) and includes the Southern Till Plain Natural Division of the natural divisions of Illinois (Schwegman, 1973; 1997; IDNR, 2018) in south-central Illinois. South-central Illinois is a dissected Illinoian till plain south of the terminal Wisconsin moraine. This region consists of nearly level to gently sloping, old till plains. Stream valleys are shallow and generally are narrow. Elevation is about 660 feet (200 meters), increasing gradually from south to north. Local relief is generally low on the broad, flat till plains and flood plains and high on the dissected hills bordering rivers or drainage systems. The Kaskaskia, Little Muddy, Little Wabash, Embarras, and Skillet Fork rivers are part of this area. This region is covered with loess, which overlies old glacial drift (Illinoian till) that has a high content of clay. Fragipans are also present. Pennsylvanian limestone and shale bedrock underlay the glacial till. The dominant soil orders in this region are Alfisol and Mollisol. The soils in the area predominantly have a mesic soil temperature regime, an aquic or udic soil moisture regime, and mixed or smectitic mineralogy. They generally are very deep, well drained to poorly drained, and loamy or clayey. (USDA-NRCS, 2006).

Classification relationships

Major Land Resource Area (MLRA) (USDA-NRCS, 2006): 113 – Central Claypan Areas, Eastern Part U.S. Forest Service Ecoregions (Cleland et al. 2007): Domain: Humid Temperate Domain Division: Hot Continental Division Province: Eastern Broadleaf Forest (Continental) Province Code: 222 Section: Central Till Plains, Oak-Hickory Section Section Code: 222G

Ecological site concept

This woodland community type is found in south-central Illinois in the northern region of the Central Claypan Areas MLRA. Clayey Till Backslope Woodland ecological sites are scattered along transitional areas between broad summit ecological sites and steeper hillslope ecological sites. This ecological site occurs on soils that formed in clayey glacial till containing a strongly developed brownish colored paleosol. The historic pre-European settlement vegetation on this ecological site had an open tree canopy dominated by white oak (*Quercus alba* L.)*, bur oak (*Quercus macrocarpa* Michx.), or less commonly swamp white oak (*Quercus bicolor* Willd.) (White, 1978). The trees are generally shorter in stature than those occurring in a forested situation, and older trees typically exhibit spreading lower branches. Shagbark hickory (*Carya ovata* (Mill.) K. Koch), pin oak (*Quercus palustris* Münchh.), post oak (*Quercus stellata* Wangenh.), black oak (*Quercus velutina* Lam.), and red maple (*Acer rubrum* L.) were occasional canopy associates. Shrubs were usually sparse, sometimes scattered ericaceous species, but more typically New Jersey tea (*Ceanothus americanus* L.) or American hazelnut (*Corylus americana* Walter). The herbaceous flora is usually dominated by grasses and forbs such as little bluestem (*Schizachyrium scoparium* (Michx.) Nash), Indian grass (*Sorghastrum nutans* (L.) Nash), big bluestem (*Andropogon gerardii* Vitman), sundial lupine (*Lupinus perennis* L.), sunflower (*Helianthus* spp.), butterfly weed (*Asclepias tuberosa* L.), ticktrefoil *Desmodium* spp., and lespedeza (*Lespedeza* spp.) and Pennsylvania sedge (*Carex pensylvanica* Lam.) (Anderson, 1975; Brugam et.al., 2016; Coates, 1992; Edgin, 1996, 2002, 2003; Anderson et.al., 2007; Taft et.al., 1994; Edgin and Ebinger. 1997; White, 1978). The canopy was probably moderately tall but less dense and less structurally diverse than lower backslope forest ecological sites (White, 1978). Increased light from an open canopy causes a diversity of woodland ground flora species to flourish, especially prairie grasses. Woodlands are distinguished from forest, by their relatively open understory, and the presence of sun-loving ground flora species (White, 1994). Fire was the primary disturbance factor that maintains this ecological site, while drought, windthrow, and grazing were secondary factors (LANDFIRE 2009). * All plant common and scientific names in this document were obtained from the U.S. Department of Agriculture – Natural Resources Conservation Service National PLANTS Database (USDA NRCS, 2018).

Associated sites

F113XY911IL	Loamy Till Backslope Forest Forest ecological site on loess over till loamy soils with slopes greater than 5 percent.
R113XY904IL	Upland Prairie Prairie ecological site is often upslope but on dark colored soils associated with nearly level till plains.
F113XY912IL	Wet Clayey Till Backslope Woodland Wet Clayey Till Backslope Woodland ecological sites are mapped as a complex with Clayey Till Backslope Woodlands.

Similar sites

F113XY912IL	Wet Clayey Till Backslope Woodland Wet Clayey Till Backslope Woodland ecological sites also support an oak-hickory community, but poorer drainage results in a wetter growth environment that is less productive with more moisture tolerant species present.
--------------------	---

Table 1. Dominant plant species

Tree	(1) <i>Quercus alba</i> (2) <i>Quercus macrocarpa</i>
Shrub	(1) <i>Ceanothus americanus</i> (2) <i>Corylus americana</i>
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Lespedeza</i>

Physiographic features

This site is on convex slopes along drainageways in dissected uplands with slopes of 5 to 15 percent. The site generates runoff to adjacent, downslope ecological sites. This site does not flood.

Table 2. Representative physiographic features

Hillslope profile	(1) Backslope
Landforms	(1) Upland > Till plain
Runoff class	High
Elevation	160 – 210 m
Slope	10 – 20 %

Water table depth	150 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

The soil temperature regime of MLRA 113 is classified as mesic, where the mean annual soil temperature is between 47 and 59°F. Temperature and precipitation occur along a north-south gradient, where temperature and precipitation increase the further south you travel (USDA-NRCS 2006). The majority of the precipitation occurs as rainfall in the form of convective thunderstorms during the growing season.

Table 3 Representative climatic features

Frost-free period (characteristic range)	150-160 days
Freeze-free period (characteristic range)	190-190 days
Precipitation total (characteristic range)	1,090-1,120 mm
Frost-free period (actual range)	150-170 days
Freeze-free period (actual range)	190-190 days
Precipitation total (actual range)	1,090-1,120 mm
Frost-free period (average)	160 days
Freeze-free period (average)	190 days
Precipitation total (average)	1,120 mm

- (1) CHARLESTON [USC00111436], Charleston, IL
- (2) NEWTON [USC00116157], Newton, IL
- (3) EFFINGHAM 3SW [USW00093816], Effingham, IL

Influencing water features

Clayey Till Backslope Woodlands are not influenced by wetland or riparian water features. Precipitation is the main source of water for this ecological site. Infiltration is slow (Hydrologic Group C/D), and surface runoff is moderate to high (SSS NRCS WSS, 2018). Surface runoff contributes some water to downslope ecological sites. These areas have a claypan or clay layer at or near the surface, with a slow rate of water transmission. Water seeps from the loess up slope and runs onto these soils at the interface of materials at the surface of the till.

Soil features

These soils are very deep, well drained, and slowly permeable. They formed in glacial till that contains a strongly developed brown mottled paleosol. They are very strongly acid to moderately acid. The dominant clay mineral in the upper part of the control section is montmorillonite, and in the lower part of the control section and substratum it is illite. The control section averages between 35 and 45 percent clay, 15 and 35 percent sand, and 2 and 3 percent gravel. Soils of this ecological site are in the Alfisol order, further classified as fine, smectitic, mesic Chromic Vertic Hapludalfs. Soil series associated with this site include Ursa.

Table 4. Representative soil features

Parent material	(1) Till
Surface texture	(1) Loam
Family particle size	(1) Fine
Drainage class	Well drained
Permeability class	Very slow
Soil depth	150 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (Depth not specified)	7.62 – 21.34 cm
Calcium carbonate equivalent (Depth not specified)	0 – 10 %
Electrical conductivity (Depth not specified)	Not specified
Sodium adsorption ratio (Depth not specified)	Not specified
Soil reaction (1:1 water) (Depth not specified)	4.5 – 7.8

Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The MLRA lies within the transition zone between the eastern deciduous forests and the tallgrass prairies. The heterogeneous topography of the area results in variable microclimates and fuel matrices that in turn are able to support prairies, savannas, woodlands, and forests. Clayey Till Backslope Woodlands form an aspect of this vegetative continuum. This ecological site occurs on convex slopes along drainageways in dissected uplands with slopes of 5 to 15 percent on well drained, slowly permeable soils. Species characteristic of this ecological site consist of an open canopy of oaks and hickories with a continuous understory of herbaceous vegetation.

Fire is a critical factor that maintains Clayey Till Backslope Woodlands. Fire typically consisted of low- to moderate-severity surface fires every 15 to 25 years ((Anderson, 1975; Brugam et.al., 2016; LANDFIRE 2009). Ignition sources included summertime lightning strikes from convective storms and bimodal, human ignitions during the spring and fall seasons. Native Americans regularly set fires to improve sight lines for hunting, drive large game, improve grazing and browsing habitat, agricultural clearing, and enhance vital ethnobotanical plants (Barrett 1980; LANDFIRE 2009).

Drought, grazing, and windthrow have also played a role in shaping this ecological site. Clayey Till Backslope Woodland ecological sites were also subjected to occasional disturbances from wind and ice, as well as grazing by native large herbivores, such as bison (*Bos bison*), prairie elk (*Cervus elaphus*), and white-tailed deer (*Odocoileus virginianus*). Wind and ice would have periodically opened the canopy up by knocking over trees or breaking substantial branches off canopy trees.

Grazing by large native herbivores would have effectively kept understory conditions more open, creating conditions more favorable to oak reproduction and woodland ground flora species (Anderson, 1982; Irland 2000; Peterson 2000). When coupled with fire, periods of drought, herbivory, and high wind events can greatly delay the establishment and maturation of woody vegetation (Pyne et al. 1996).

Extensive conversion for agriculture has fragmented this system. These ecological sites are moderately productive. Today, many of these ecological sites have been cleared and converted to pasture and cropland. Uncontrolled domestic grazing has also impacted these communities, further diminishing the diversity of native plants and introducing species that are tolerant of grazing, such as coralberry (*Symphoricarpos orbiculatus* Moench), gooseberry (*Ribes* spp.), and Virginia creeper (*Parthenocissus quinquefolia* (L.) Planch.). Grazed sites also have a more open understory. In addition, soil compaction and soil erosion can be a problem and lower productivity. In the long term absence of fire, woody species, especially hickory (*Carya* spp.) and hophornbeam (*Ostrya virginiana* (Mill.) encroach into these woodlands (IDNR, 2018). Once established, these woody plants can quickly fill the existing understory increasing shade levels with a greatly diminished ground flora (Dey and Kabrick, 2015). Remaining woodland ecological sites have a younger (50 to 80 years) canopy layer whose species composition and quality has been altered by timber harvesting practices.

A provisional state and transition diagram is depicted. Detailed descriptions of each state, transition, plant community, and pathway follow the model. This model is based on available experimental research, field observations, professional consensus, and interpretations. It may change as knowledge increases.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

No field plots were available for this site. A review of the scientific literature and professional experience were used to approximate the plant communities and ecological dynamics for this provisional ecological site. Information for the state-and-transition model was obtained from the same sources. All community phases are considered provisional based on the sources identified in ecological site description.

References

- Anderson, R.C. and M.R. Anderson. 1975. The presettlement vegetation of Williamson County, Illinois.. *Castanea* 40:345–363.
- Anderson, R.C. 1982. An evolutionary model summarizing the roles of fire, climate, and grazing animals in the origin and maintenance of grasslands. Pages 297–308 in , , and , editors. *Grasses and grasslands: systematics and ecology*.
- Anderson R. C., J. S. Fralish, and J. M. Baskin. 2007. Presettlement forests of Illinois. G. V. Burger, J. E. Ebinger, and G. S. Wilhelm, eds., *Proceedings of the Oak Woods Management Workshop* 9–19.
- Barrett, S.W. 1980. Indians and fire.. *Western Wildlands* 17–20.
- Briggs, J.M., A.K. Knapp, and B.L. Brock. 2002. Expansion of woody plants in tallgrass prairie: a fifteen- year study of fire and fire-grazing interactions. *The American Midland Naturalist* 147:287–294.
- Brinson, M.M. 1993. A hydrogeomorphic classification for wetlands.
- Brugam, R.B., P.D. Kilburn, and L.L. Luecking. 2016. Pre-settlement Vegetation of Greene, Jersey and Macoupin Counties along the Prairie/Forest Border in Illinois.. *Transactions of the Illinois State Academy of Science* 109:9–17.
- Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 1–92.
- Coates, D.T., K.J. Lyman, and J.E. Ebinger. 1992. Woody vegetation structure of a post oak flatwoods in Illinois.. *Castanea* 57:196–201.
- Comer, P.J., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003 (Date accessed). *Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems*.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep water habitats of the United States.. U.S. Dept. of Interior, Fish & Wildlife Service, Office of Biological Services, Washington DC. FWS/OBS-79/31 1–142.
- Dey, D.C. and J.M. Kabrick. 2015. Restoration of Midwestern oak woodlands and savannas.. Pages 401–428 in *Restoration of Boreal and Temperate Forests*, Second Edition. CRC Press, Boca Raton, Florida, USA..
- Edgin, B. 1996. Barrens of pre-settlement Lawrence County, Illinois.. Pages 59–65 in *Proceedings of the 15th North American Prairie Conference*.
- Edgin, B. and J.E. Ebinger. 1997. Barrens and the pre-settlement prairie/forest interface in Crawford County, Illinois.. *Castanea* 62:260–267.
- Edgin, B., R. Beadles, and J.E. Ebinger. 2002. Woody Composition and Structure of Karcher's Post Oak Woods Nature Preserve, Hamilton County, Illinois.. *Transactions of the Illinois State Academy of Science* 95:251–259.

Edgin B., W. E. McClain, R. Gillespie, and J. E. Ebinger. 2003. Vegetation composition and structure of Eversgerd Post Oak Flatwoods, Clinton County, Illinois.. *Northeast Naturalist* 10:111–118.

Illinois Department of Natural Resources (IDNR). March 2018 (Date accessed). Natural Divisions - Southern Till Plain..

Irland, L.C. 2000. Ice storms and forest impacts.. *The Science of the Total Environment* 262:231–242.

Kilburn, P. and R.B. Brugam. 2014. Inventory of Vegetation Studies in Illinois Based on the Public Land Survey Records.. *Transactions of the Illinois State Academy of Science* 107:13–17.

USGS. 2009 (Date accessed). **Landfire National Vegetation Dynamics Models**. <http://www.LANDFIRE.gov/index.php>.

Mohlenbrock R. H. and D. M. Ladd. 1978. Distribution of Illinois Vascular Plants. Southern Illinois Univ. Press, Carbondale and Edwardsville, IL. 281p.

Mohlenbrock R. H. 2003. Vascular Flora of Illinois. Vascular Flora of Illinois, 3rd edition. Southern Illinois University Press, Carbondale, Illinois. 1–736.

National Cooperative Soil Survey (NCSS). 2018 (Date accessed). **National Cooperative Soil Characterization Database**. <https://ncsslabsdatamart.sc.egov.usda.gov/>.

National Oceanic and Atmospheric Administration (NOAA). 2018 (Date accessed). **Climate Data 1980-2010**. <https://www.ncdc.noaa.gov/data-access/land-based-station-data/find-station>.

NatureServe. 2018 (Date accessed). **Association Detail Report: CEGLO02427**. <http://explorer.natureserve.org>.

Nelson, P. 2010. The Terrestrial Natural Communities of Missouri. Revised edition. Missouri Natural Areas Committee, Department of Natural Resources and the Department of Conservation, Jefferson City. 549p.

Pyne, S.J., P.L. Andrews, and R.D. Laven. 1996. Introduction to Wildland Fire, Second Edition. Introduction to Wildland Fire, Second Edition. John Wiley and Sons, Inc. New York, New York. 1–808.

Schwegman, J.E., G.B. Fell, M.D. Hutchinson, G. Paulson, W.M. Shephard, and J. White. 1973. The natural divisions of Illinois. Comprehensive plan for the Illinois Nature Preserve system. Part 2. Illinois Nature Preserves Commission, Rockford, IL 1–32.

. 2018 (Date accessed). **Web Soil Survey (SSS NRCS WSS)**. <https://websoilsurvey.sc.egov.usda.gov/>.

SSS NRCS OSD and . 2018 (Date accessed). **Official Soil Series Descriptions**. <https://soilseries.sc.egov.usda.gov/osdname.aspx>.

Taft, J.B., M.W. Schwartz, and L.R. Philippe. 1995. Vegetation ecology of flatwoods on the Illinoian till plain. *Journal of Vegetation Science* 6:647–666.

United States Department of Agriculture, . 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin... *USDA Handbook* 296 1–682.

USDA, N. 2018 (Date accessed). **The PLANTS Database**. <http://plants.usda.gov>.

Voigt J. W. and R. H. Mohlenbrock. 1964. Plant communities of southern Illinois. Plant communities of southern Illinois. Southern Illinois University Press, Carbondale 1–202.

White J. 1978. Natural Areas Inventory Technical Report. Natural Areas Inventory Technical Report: Volume I, Survey Methods and Results. Illinois Natural Areas Inventory, Department of Landscape Architecture, University of Illinois at Urbana/Champaign 1–426.

White, J. and M. Madany. 1978. Classification of natural communities in Illinois (Appendix 30). In J. White, Illinois Natural Areas Inventory Technical Report. Volume 1: Survey Methods and Results. Illinois Natural Areas Inventory, Department of Landscape Architecture, University of Illinois at Urbana/Champaign. 310–405.

Other references

Relationship to other established ecological classifications:

Biophysical Setting (LANDFIRE, 2009); the reference community of this ecological site is most similar to: North-Central Interior Dry-Mesic Oak Forest and Woodland (CES202.046)

National Vegetation Classification System (NatureServe, 2018): the reference community of this ecological site is most similar to: *Quercus stellata* - *Quercus velutina* / *Schizachyrium scoparium* Woodland (CEGL005281) and *Quercus alba* - *Quercus macrocarpa* - *Quercus bicolor* Woodland Alliance (A3324)

Illinois Natural Areas Survey (INAS) (White, 1978); the reference community of this ecological site is most similar to: INAS Community Class – Savanna; Natural community – Dry-Mesic Savanna

Contributors

Douglas Wallace
Ralph Tucker
Zach Weber

Approval

Suzanne Mayne-Kinney, 5/17/2024

Acknowledgments

Contact information for primary authors: Ralph Tucker (ralph.tucker@mo.usda.gov), Soil Scientist, United States Department of Agriculture - Natural Resources Conservation Service (USDA-NRCS), Union, MO; Zach Weber (zach.weber@il.usda.gov), Soil Scientist, USDA-NRCS, Olney, IL; Douglas Wallace (doug.wallace@mo.usda.gov), Ecologist, USDA-NRCS, Columbia, MO.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	08/31/2026
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
