

Ecological site F114XA302IN

Acidic Upland Forest

Last updated: 9/26/2024
Accessed: 09/24/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 114X–Southern Illinois and Indiana Thin Loess and Till Plain

MLRA 114A makes up about 4,550 square miles (11,795 square kilometers). The three parts of this MRLA are mostly in the Till Plains Section of the Central Lowland Province of the Interior Plains. The western third of the western part is in the Highland Rim Section of the Interior Low Plateaus Province of the Interior Plains. The eastern half of the eastern part is in the Kanawha Section of the Appalachian Plateaus Province of the Appalachian Highlands. Both large and small tributaries of the Ohio River dissect the nearly level to very steep glaciated uplands in this area. The major streams and rivers have well defined valleys with broad flood plains and numerous stream terraces. The flood plains along the smaller streams are narrow. Broad summits are nearly level to gently sloping. Elevation ranges from 320 feet (100 meters) on the southernmost flood plain along the Ohio River to 1,250 feet (380 meters) on the highest ridges. Local relief is mainly 10 to 50 feet (3 to 15 meters), but it can be 50 to 100 feet (15 to 30 meters) along drainageways and streams. Also, the Ohio River bluffs are as much as 300 feet (90 meters) above the river valley floor.

Classification relationships

USFS: 222 Eastern Broadleaf Forest (Continental) Province. Homoya's Natural Regions of Indiana: Bluegrass Region The following NatureServe Explorer Ecological System has a high level of probability to match the ecological site found on these soils. Southern Interior Low Plateau Dry-Mesic Oak Forest-Unique Identifier: CES202.898

Ecological site concept

Acidic Upland Forest sites were historically a mature hardwood forest community with oaks and hickories being the dominant canopy species. These upland sites are found on acidic soils. Decades of disturbance (logging, grazing, invasive vegetation, fire suppression, etc.) have impacted most sites that are still wooded. These disturbances have transitioned most sites to a mixed hardwood forest with less of an oak component that was historically present and an increase in maple, ash and tulip poplar. The majority of areas have now been converted for agricultural use.

Associated sites

F114XA502IN	Till Uplands Till Upland Forest. These sites are on upland till plains.
-------------	---

Similar sites

F114XA305IN	Non-Acidic Upland Forest The non-acidic upland forest sites also are dominated by a oak-hickory forest; however the species on these sites are those adapted to calcareous soils.
-------------	---

Table 1. Dominant plant species

Tree	(1) <i>Quercus velutina</i> (2) <i>Carya glabra</i>
Shrub	(1) <i>Vaccinium</i>
Herbaceous	(1) <i>Agrimonia</i> (2) <i>Desmodium</i>

Physiographic features

These sites are generally located on hillsides, ridges, and till plains.

Table 2. Representative physiographic features

Landforms	(1) Upland > Hill (2) Upland > Ridge (3) Till plain > Till plain
Runoff class	Low to high
Flooding frequency	None
Ponding frequency	None
Elevation	100 – 370 m
Slope	10 – 20 %
Water table depth	20 – 150 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	Not specified

Slope	0 – 30 %
Water table depth	Not specified

Climatic features

About 60 percent of the precipitation falls during the freeze-free period. Most of the rainfall occurs as high-intensity, convective thunderstorms during summer. Snowfall is common in winter. The freeze-free period averages about 180 days.

Table 4 Representative climatic features

Frost-free period (characteristic range)	150-160 days
Freeze-free period (characteristic range)	180-180 days
Precipitation total (characteristic range)	1,170-1,220 mm
Frost-free period (actual range)	150-160 days
Freeze-free period (actual range)	180-180 days
Precipitation total (actual range)	1,170-1,220 mm
Frost-free period (average)	160 days
Freeze-free period (average)	180 days
Precipitation total (average)	1,190 mm

- (1) SEYMOUR 2 N [USC00127935], Seymour, IN
- (2) LEXINGTON 3N [USC00124977], Lexington, IN
- (3) MILFORD [USC00335268], Milford, OH
- (4) NORTH VERNON 2 ESE [USC00126435], North Vernon, IN

Influencing water features

There are no riparian or wetland features influencing these sites.

Soil features

Soils associated with this site are acidic, often on black or gray shale. Series include Deputy, Gilpin, Jennings, Jessietown, Loudon, Loudonville, Mechanicburg, Rarden Scottsburg Trappist, Weddel, and Whitcomb.

Table 5. Representative soil features

Parent material	(1) Residuum – shale (2) Till (3) Loess (4) Slope alluvium
Surface texture	(1) Silt loam (2) Silty clay (3) Silty clay loam
Drainage class	Somewhat poorly drained to well drained
Permeability class	Slow to moderately rapid
Depth to restrictive layer	60 – 180 cm
Soil depth	40 – 150 cm
Surface fragment cover ≤3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (Depth not specified)	10.16 – 15.24 cm
Soil reaction (1:1 water) (Depth not specified)	4.5 – 10
Subsurface fragment volume ≤3" (Depth not specified)	0 – 10 %
Subsurface fragment volume >3" (Depth not specified)	0 – 10 %

Table 6. Representative soil features (actual values)

Drainage class	Not specified
----------------	---------------

Permeability class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (Depth not specified)	7.62 – 17.78 cm
Soil reaction (1:1 water) (Depth not specified)	Not specified
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The historic plant community of this ecological site is an oak-hickory forest. Common species on these sites may include black oak (*Quercus velutina*), white oak (*Q. alba*), chestnut oak (*Q. montana*), shagbark hickory (*Carya ovata*), pignut hickory (*Carya glabra*), scarlet oak (*Q. coccinea*), flowering dogwood (*Cornus florida*), hophornbeam (*Ostrya virginiana*), and sassafras (*Sassafras albidum*). This site is naturally dominated by stable, uneven-aged forests with the canopy dynamics being driven by gap-phase regeneration.

Decades of disturbance (logging, grazing, fire suppression, etc.) have impacted these sites. Many sites now have a mixed hardwood forest with much less of an oak component that was historically present. Severe or long-term disturbances have transitioned sites toward a community of shade-tolerant species such as maple, ash and tulip poplar.

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

No field monitoring was conducted as part of this PES development. Future ESD development will result in plant community edits, soil mapunits being added or removed from this grouping, and/or additions or modifications to the narratives, tables, vegetation descriptions and state and transition model.

Other references

Braun, E. Lucy. 2001. Deciduous forests of eastern North America. Caldwell, N.J.: Blackburn Press.

Homoya, M. A., Abrell, D. B., Aldrich, J. R., & Post, T. W. (1985). The Natural Regions of Indiana. Indiana Academy of Science , 94, 245-269.

Kartesz, J. T. (2011). Density Gradient Map Samples Produced From BONAP's Floristic Synthesis. Retrieved 12 12, 2011, from Biota of North America Program: <http://bonap.org/diversity/diversity/diversity.html>

NatureServe. (2011). An online encyclopedia of life [web application]. NatureServe, Arlington, VA, USA [Online: [www. NatureServe.org/explorer](http://www.NatureServe.org/explorer)] .

Jackson, Marion T. 1997. The Natural heritage of Indiana. Bloomington: Indiana University Press, published in association with the Indiana Department of Natural Resources and the Indiana Academy of Science.

USDA. (2007). Ecological Subregions: Sections and Subsections for the Conterminous United States. Washington, DC: USDA - Forest Service.

USDA. (2006). Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U. S. Department of Agriculture, Natural Resources Conservation Service. U. S. Department of Agriculture Handbook 296.

USGS. (2010). LANDFIRE Biophysical Settings. Retrieved from <http://www.landfire.gov>

Whitaker, John O., Charles J. Amlaner, Marion T. Jackson, George R. Parker, and Peter Evans Scott. 2012. Habitats and ecological communities of Indiana presettlement to present. Bloomington: Indiana University Press.

Approval

Greg Schmidt, 9/26/2024

Acknowledgments

PES documents developed for adjacent MLRAs in Indiana and Ohio served as a source of information as these regions often shared similar soil series with MLRA 114A. NRCS county soil surveys where a valuable reference including tree species observed on site by NRCS staff. Soil Survey and NRCS Indiana resource soil scientists contributed field observation, field notes, and extensive soil mapping expertise. In accordance with Federal Civil Rights law and U.S. Department of Agriculture (USDA) Civil Rights regulations and policies, the USDA, its agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior credible activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident. Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English. To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](http://www.usda.gov/howtofile) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov. USDA is an equal opportunity provider, employer, and

lender.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	A Arends, ESI Specialist
Contact for lead author	
Date	09/24/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
