

Ecological site F114XB502IN

Wet Till Upland Forest

Last updated: 11/16/2023

Accessed: 09/24/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 114X–Southern Illinois and Indiana Thin Loess and Till Plain

This MLRA is a loess-covered till plain with broad, nearly level summits and steeper slopes in areas dissected by tributaries of the Ohio and Mississippi Rivers. It is used to produce cash crops, feed grain, and livestock. This MLRA is in Indiana (47 percent), Illinois (38 percent), and Ohio (15 percent) in four separate areas. It makes up about 10,388 square miles (26,904 square kilometers). This area is in the Till Plains section of the Central Lowland province of the Interior Plains. Well defined valleys with broad flood plains and numerous stream terraces are along the major streams and rivers. The flood plains along the smaller streams are narrow. Broad summits are nearly level or gently sloping. Steep slopes are along rivers and streams. Elevation ranges from 310 feet (90 meters) on the southernmost flood plains to 1,340 feet (410 meters) on the highest ridges. Local relief is mainly 10 to 50 feet (3 to 15 meters) but can be 50 to 100 feet (15 to 30 meters) along drainageways and streams. The Little Miami River flows through the part of this MLRA in Ohio. The Ohio River flows along the southernmost boundary in some parts of this area in Ohio. The Kaskaskia River flows through the part of this area in Illinois. Tributaries to the Mississippi and Ohio Rivers drain this MLRA. This area is covered dominantly by loess and Illinoian-age till or outwash. Most of the loess is Late Wisconsin-age Peoria Loess. In some places the Peoria Loess is underlain by Early Wisconsin-age Roxana Silt or by sandier or grittier loess. The loess ranges from 3 to 7 feet (1 or 2 meters) in thickness on stable summits and does not occur on some of the steeper slopes. The underlying Illinoian-age till and outwash commonly contain a paleosol. Meltwater outwash and lacustrine and alluvial deposits are on some of the stream terraces along the major tributaries. The till and outwash are underlain by several bedrock systems. Mississippian and Pennsylvanian bedrock occurs mostly in the western part of the MLRA. Ordovician, Silurian, and Devonian bedrock occurs mostly in the central part. Bedrock outcrops are common on the bluffs along the large rivers and their major tributaries. They also are evident at the base of steep slopes along minor streams and drainageways. The average annual precipitation ranges from 39 to 47 inches (990 to 1,190 millimeters) with a mean of 42 inches (1,060 millimeters). The annual temperature ranges from 53 to 56 degrees F (11.8 to 13.6 degrees C) with a mean of 55 degrees F (13 degrees C). The freeze-free period ranges from 185 to 215 days with a mean of 200 days. The dominant soil orders are Alfisols and Entisols. The soils in the area have a mesic soil temperature regime, an aquic or udic soil moisture regime, and mixed or smectitic mineralogy. They are deep or very deep, poorly drained to well drained, and loamy, silty, or clayey. Although limited in extent, some soils have a natric horizon in the part of the MLRA in Illinois. The main soils and their series: Albaqualfs that formed in loess or loess over pedisodiment on till plains (Marine series); Endoaqualfs that formed in loess or loess over pedisodiment on till plains (Ocone series); Fluvaquents that formed in alluvium on flood plains (Wakeland series); Fragiudalfs that formed in loess over pedisodiment over till (Cincinnati series) and loess over till (Rossmoyne series) on till plains; Glossaqualfs that formed in loess over till on till plains (Avonburg, Clermont, and Cobbsfork series) Hapludalfs that formed in till (Hickory series) and loess over pedisodiment (Homen series) on till plains. The soils on uplands support natural hardwoods. Oak, hickory, beech, and sugar maple are the dominant species. Native grasses grow in some scattered areas between the trees. The soils in low-lying areas support mixed forest vegetation. Pin oak, shingle oak, sweetgum, and black oak are the dominant species on the wetter sites. White oak, black oak, northern red oak, hickory, yellow-poplar, ash, sugar maple, and black walnut grow on the better drained sites. Honey locust is dominant on soils that formed in shaly limestone residuum. Silver maple, eastern cottonwood, American sycamore, pin oak, elm, and sweetgum grow along rivers and streams. Black walnut is abundant on very deep, well drained soils on some small flood plains. Sedge and grass meadows and scattered trees are on some low-lying sites. Most of this MLRA is in farms and used to produce corn, soybeans, and livestock. Some small grains, including winter wheat, oats, and grain sorghum, also are grown. A small acreage is used for specialty crops, such as popcorn and apple orchards. The grassland supports introduced and native grasses. The forested areas are mainly on steep valley sides and in low-lying parts of flood plains. Surface coal mines make up a small acreage. (USDA, Natural Resources Conservation Service. 2022)

LRU notes

LRU 114XB is in two separate areas in Illinois (66 percent) and Indiana (34 percent). It makes up about 7,005 square miles (18,150 square kilometers). It includes the towns of Brazil, Bloomfield, Cloverdale, and Spencer, Indiana, and Carlyle, Nashville, Hillsboro, Greenville, Vandalia, and Pinckneyville, Illinois. Interstates 55, 64, and 70 cross the part of the MLRA in Illinois. They converge in St.

Louis, which is just west of this MLRA. The east edge of the Scott Air Force Base is on the western edge of the area in Illinois. This area is in the Till Plains Section of the Central Lowland Province of the Interior Plains. Both large and small tributaries of the West Fork of the White River, the Eel River, the Kaskaskia River, and the Little Muddy River dissect the nearly level to very steep uplands. Well defined valleys with broad flood plains and numerous stream terraces are along the major streams and rivers. The flood plains along the smaller streams are narrow. Broad summits are nearly level to gently sloping. Elevation ranges from 350 feet (105 meters) on the southernmost flood plains along the Ohio and Wabash Rivers to 1,190 feet (365 meters) on the highest ridges. Local relief is mainly 10 to 50 feet (3 to 15 meters), but it can be 50 to 100 feet (15 to 30 meters) along drainageways and streams. It generally is low on broad, flat till plains and flood plains and high on the dissected hills bordering rivers or drainage systems.

Classification relationships

Major Land Resource Area (MLRA) (USDA-NRCS, 2022): 114X–Southern Illinois and Indiana Thin Loess and Till Plain U.S. Forest Service Ecoregions (Cleland et al. 2007): Domain: Humid Temperate Domain Division: Hot Continental Division Province: Eastern Broadleaf Forest (Continental) Province Code: 222

Ecological site concept

The historic reference plant community had a wide variety of dominant deciduous hardwood trees. Species were those tolerant of seasonally wet soils including swamp white oak green ash, and northern red oak. Sites in Indiana often have pin oak and/or sweetgum present. Understory species on these sites were extremely diverse and varied depending on shade, microtopography and fine soil characteristics. Today, disturbances such as logging, grazing, oak removal, non-native vegetation and hydrological modifications have created many different forest composition variations. Wetter micro-sites will favor more pin oak, swamp white oak, green ash, bitternut hickory, and maple. Better drained micro-sites will include more mesic species such as northern red oak, bur oak, white ash, and Shumard's oak.

Associated sites

F114XB503IN	Till Upland Forest The Till Upland Forest ecological site and the Wet Till Upland Forest ecological site occur on similar landscape positions and are often intermixed. The Till Upland Forest does not have a water table between 0-42 inches, whereas the Wet Till Upland Forest ecological site does.
-------------	---

Similar sites

F114XB403IN	Wet Outwash Upland Forest Wet Outwash Upland Forest has a parent material of outwash and occurs on flood plains, stream terraces, terraces and fans, Wet Till Upland Forest ecological sites has a parent material of till and occurs on till plains, and ground moraines.
-------------	---

Table 1. Dominant plant species

Tree	(1) <i>Quercus</i> (2) <i>Fraxinus pennsylvanica</i>
Shrub	(1) <i>Cornus</i> (2) <i>Asimina triloba</i>
Herbaceous	(1) <i>Carex</i> (2) <i>Elymus virginicus</i>

Physiographic features

These sites are found on multiple landforms including till plains, hills, and/or ground moraines.

Table 2. Representative physiographic features

Landforms	(1) Till plain (2) Ground moraine (3) Draw (4) Hillslope (5) Ridge
Runoff class	Low to high
Flooding frequency	None
Ponding frequency	None
Elevation	90 – 310 m
Slope	0 – 20 %
Water table depth	0 – 110 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

About 60 percent of the precipitation falls during the freeze-free period. Most of the rainfall occurs as high-intensity, convective thunderstorms during summer. Snowfall is common in winter. The freeze-free period averages about 183 days.

Table 3 Representative climatic features

Frost-free period (characteristic range)	150-170 days
Freeze-free period (characteristic range)	180-190 days
Precipitation total (characteristic range)	1,040-1,170 mm
Frost-free period (actual range)	140-170 days
Freeze-free period (actual range)	170-190 days
Precipitation total (actual range)	1,040-1,190 mm

Frost-free period (average)	150 days
Freeze-free period (average)	180 days
Precipitation total (average)	1,120 mm

- (1) SHAKAMAK SP [USC00127959], Jasonville, IN
- (2) SPENCER [USC00128290], Spencer, IN
- (3) CARBONDALE SOUTHERN IL AP [USW00093810], De Soto, IL
- (4) WATERLOO [USC00119002], Waterloo, IL
- (5) JERSEYVILLE 2 SW [USC00114489], Jerseyville, IL
- (6) PANA 3E [USC00116579], Pana, IL

Influencing water features

These sites have seasonally high water tables.

Soil features

The soil series associated with this site are: Atlas, Blair, Fishhook, Grantfork, and Vigo. Parent materials is till but can be overlaid by loess, paleosol and/or pedisediment. Atlas and Fishhook have restrictive layer (densic material) in the soil from 30 to 61 inches in depth.

Table 4. Representative soil features

Parent material	(1) Till
Surface texture	(1) Silt loam (2) Silty clay loam
Family particle size	(1) Fine (2) Fine-silty
Drainage class	Somewhat poorly drained
Permeability class	Slow to moderate
Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	12.7 – 22.86 cm

Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	4.5 – 7.8
Subsurface fragment volume <=3" (0-101.6cm)	0 – 10 %
Subsurface fragment volume >3" (0-101.6cm)	Not specified

Ecological dynamics

Wet Till Upland reference sites include swamp white oak, bur oak, northern red oak , and green ash. Indiana sites often include sweetgum, pin oak and/or tulip poplar.

Wetter micro-sites will favor more pin oak, swamp white oak, green ash, bitternut hickory, and maple. Better drained micro-sites will include more mesic species such as northern red oak, bur oak, white ash, and Shumard's oak.

These disturbances will transition this community to a woodland dominated by fast-growing, shade-tolerant species such as maple, ash, poplar, beech, and elm. The resulting dense canopy shade on these sites often results in a more sparse understory community.

Many forest sites in Indiana and Illinois have been invaded by non-native plant species. Many species may be present depending on disturbances and seed sources.

Few high-quality, old-growth communities remain. Agriculture is the largest use of these soils in MLRA 114x.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

No field monitoring was conducted as part of this PES development. Future ESD development may result in plant community edits, soil mapunits being added or removed from this grouping, and/or additions or modifications to the narratives, tables, vegetation descriptions and state and transition model. There are multiple Atlas complex mapunits included in this initial PES group. Atlas soils have a high clay content, so dominant species would have included swamp white oak, bur oak, pin oak, green ash, hackberry, and elm. For further information on Atlas soils refer to "Wet Clayey Till Backslope Woodland" (F113XY912IL) developed in the adjacent MLRA 113.

Other references

Anderson, R. C., ?J. S. Fralish, ?Jerry M. Baskin. 2007. Presettlement forests of Illinois. In Proceedings of the Oak Woods Management Workshop, ed. G. V. Burger, J. E. Ebinger, and G. S. Wilhelm, pp. 9-19. Charleston, Ill.: Eastern Illinois University.

Barrett, S.W. 1980. Indians and fire. *Western Wildlands* Spring: 17-20.

Braun, E. Lucy. 2001. *Deciduous forests of eastern North America*. Caldwell, N.J.: Blackburn Press.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. *Ecological Subregions: Sections and Subsections of the Conterminous United States*. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pp.

Comer PJ, Faber-Langendoen D, Evans R, Gawler SC, Josse C, Kittel G, Menard S, Pyne M, Reid M, Schulz K, Snow K, and Teague J. 2003. *Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems*. NatureServe, Arlington, Virginia.

Cowardin, L.M., V. Carter, F.C. Golet, & E.T. LaRoe. 1979. *Classification of wetlands and deep water habitats of the United States*. U.S. Dept. of Interior, Fish & Wildlife Service, Office of Biological Services, Washington DC.

Federal Geographic Data Committee. 2013. *Classification of wetlands and deepwater habitats of the United States*. FGDC-STD-004-2013. Second Edition. Wetlands Subcommittee, Federal Geographic Data Committee and U.S. Fish and Wildlife Service, Washington, DC

Homoya, M. A., Abrell, D. B., Aldrich, J. R., & Post, T. W. (1985). The Natural Regions of Indiana. *Indiana Academy of Science* , 94, 245-269.

Illinois Department of Natural Resources (IDNR). 2018. Natural Divisions - Southern Till Plain. Accessed; March 2018. <https://www.dnr.illinois.gov/conservation/IWAP/Documents/NaturalDivisions/SouthernTillPlain>

Jackson, Marion T. 1997. *The Natural heritage of Indiana*. Bloomington: Indiana University Press, published in association with the Indiana Department of Natural Resources and the Indiana Academy of Science.

Keyser, Tara L.; Arthur, Mary; Loftis, David L. 2017. Repeated burning alters the structure and composition of hardwood regeneration in oak-dominated forests of eastern Kentucky, USA. *Forest Ecology and Management*. 393: 1-11. <https://doi.org/10.1016/j.foreco.2017.03.015>.

Kilburn, P. and R. B. Brugam. 2014. Inventory of Vegetation Studies in Illinois Based on the Public Land Survey Records. *Transactions of the Illinois State Academy of Science*. Vol. 107, pp. 13-17.

Landfire (Landfire National Vegetation Dynamics Database). 2009. *Landfire National Vegetation Dynamics Models*. Landfire Project, USDA Forest Service, U.S. Department of Interior. (<http://www.LANDFIRE.gov/index.php>: accessed 22 February 2018).

Mohlenbrock, R. H. and D. M. Ladd. 1978. *Distribution of Illinois Vascular Plants*. Southern Illinois Univ. Press, Carbondale and Edwardsville, Ill. 282 pp.

Mohlenbrock, R. H. 2014. *Vascular Flora of Illinois*, 4rd edition. Carbondale, Illinois: Southern Illinois University Press. 736 pp.

National Cooperative Soil Survey (NCSS). National Cooperative Soil Characterization Database. Available online: <https://ncsslabdatamart.sc.egov.usda.gov/>. Accessed: February 2018.

National Oceanic and Atmospheric Administration (NOAA). 1980-2010. <https://www.ncdc.noaa.gov/data-access/land-based-station-data/find-station>.

NatureServe. 2018. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia.

Available <http://explorer.natureserve.org>. (Association Detail Report: CEGLO02427) (Accessed: May 22, 2018).

Nowacki, Gregory J.; Abrams, Marc D. 2008. The demise of fire and "mesophication" of forests in the eastern United States. *BioScience*. 58(2): 123-138.

Schwegman, J. E., G. B. Fell, M. D. Hutchinson, G. Paulson, W. M. Shephard, and J. White. 1973. Comprehensive plan for the Illinois Nature Preserve system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, IL. 32 pp.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions (SSS NRCS OSD). Available online. Accessed 2019.

USDA. 2019. The PLANTS Database (<http://plants.usda.gov>, 1 March 2018). National Plant Data Team, Greensboro, NC 27401-4901 USA.

USDA, Natural Resources Conservation Service. 2022. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture, Agriculture Handbook 296.

USDA-NRCS. 2008. Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service. Technical Note No. 190–8–76. Washington D.C.

USGS. (2010). LANDFIRE Biophysical Settings. Retrieved from <http://www.landfire.gov>

Voigt, J. W., and R. H. Mohlenbrock. 1964. Plant communities of southern Illinois. Southern Illinois University Press, Carbondale. 202 pp.

Whitaker, John O., Charles J. Amlaner, Marion T. Jackson, George R. Parker, and Peter Evans Scott. 2012. Habitats and ecological communities of Indiana presettlement to present. Bloomington: Indiana University Press.

White, J. 1994. How the terms savanna, barrens, and oak openings were used in early Illinois. In J.S. Fralish, R. C. Anderson, J.E. Ebinger and R. Szafoni, eds., *Proceedings of the North American Conference on Barrens and Savannas*, Illinois State University, Normal Illinois.

White, J. 1978. Classification of natural communities in Illinois. Natural Areas Inventory Technical Report: Volume I, Survey Methods and Results. Illinois Natural Areas Inventory, Department of Landscape Architecture, University of Illinois at Urbana/Champaign. 426 pp.

Contributors

John Allen, Acting Soil Survey Office Leader, USDA-NRCS, Indiana
Dena Anderson, Resource Soil Scientist, USDA-NRCS, Indiana
Ralph Tucker, Soil Survey Office Leader, USDA-NRCS, Missouri
Anita Arends, Ecological Site Specialist, USDA-NRCS, Illinois

Approval

Suzanne Mayne-Kinney, 11/16/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/24/2026

Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
