

Ecological site F116BY008MO

Interbedded Sedimentary Upland Woodland

Last updated: 10/06/2020

Accessed: 09/01/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 116B–Springfield Plain

The Springfield Plain is in the western part of the Ozark Uplift. It is primarily a smooth plateau with some dissection along streams. Elevation is about 1,000 feet in the north to over 1,700 feet in the east along the Burlington Escarpment adjacent to the Ozark Highlands. The underlying bedrock is mainly Mississippian-aged limestone, with areas of shale on lower slopes and structural benches, and intermittent Pennsylvanian-aged sandstone deposits on the plateau surface.

Classification relationships

Terrestrial Natural Community Type in Missouri (Nelson, 2010): The reference state for this ecological site is most similar to a Dry Chert Woodland. Missouri Department of Conservation Forest and Woodland Communities (Missouri Department of Conservation, 2006): The reference state for this ecological site is most similar to a Post Oak Woodland. National Vegetation Classification System Vegetation Association (NatureServe, 2010): The reference state for this ecological site is most similar to a *Quercus stellata* - *Quercus velutina* / *Schizachyrium scoparium* Woodland (CEGL005281). Geographic relationship to the Missouri Ecological Classification System (Nigh & Schroeder, 2002): This ecological site occurs primarily within the following Land Type Associations: Stockton Prairie/Savanna Dissected Plain Upper Sac River Oak Savanna/Woodland Low Hills Little Sac River Oak Savanna/Woodland Low Hills Upper Pomme de Terre Oak Savanna/Woodland Dissected Plain James River Oak Savanna/Woodland Low Hills

Ecological site concept

NOTE: This is a “provisional” Ecological Site Description (ESD) that is under development. It contains basic ecological information that can be used for conservation planning, application and land management. After additional information is collected, analyzed and reviewed, this ESD will be refined and published as “Approved”. Interbedded Sedimentary Upland Woodlands occur primarily along the northeast edge of the Springfield Plain, typically on lower hillslopes where interbedded shale, mudstone and limestone is near the surface. A few isolated areas to the southwest are in the Elk River watershed in McDonald County, Missouri. Soils are moderately deep to deep over sedimentary bedrock, and typically have shale and mudstone fragments in clayey subsoils. The reference plant community is woodland with an overstory dominated by white oak and black oak, and a ground flora of native grasses and forbs.

Associated sites

F116BY004MO	Low-Base Chert Upland Woodland Low-base Chert Upland Woodlands are upslope.
F116BY012MO	Interbedded Sedimentary Protected Backslope Forest Interbedded Sedimentary Protected Backslope Forests are often downslope on steep northern and eastern aspects.

F116BY036MO	Interbedded Sedimentary Exposed Backslope Woodland Interbedded Sedimentary Exposed Backslope Woodlands are often downslope on steep southern and western aspects.
--------------------	--

Similar sites

F116BY003MO	Chert Upland Woodland Chert Upland Woodlands have similar overstory species composition and landscape positions but are generally more productive.
--------------------	---

Table 1. Dominant plant species

Tree	(1) <i>Quercus alba</i> (2) <i>Quercus velutina</i>
Shrub	(1) <i>Rhus aromatica</i>
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Carex pensylvanica</i>

Physiographic features

This site is on upland summit crests, shoulders and backslopes with slopes of 3 to 20 percent. The site generates runoff to adjacent, downslope ecological sites. This site does not flood.

The following figure (adapted from Dodd, 1990) shows the typical landscape position of this ecological site, and landscape relationships with other ecological sites. Interbedded Sedimentary Upland Woodland sites are within the area labeled as “3”, and are typically on lower slopes and structural benches where the shale and mudstone crops out downslope from the overlying limestone. Steeper, lower backslopes within this area are in the Interbedded Sedimentary Backslope ecological sites. Low-base Chert Upland Woodland sites are typically upslope, and areas of Low-base Chert Upland sites are shown on upper slopes within the area labeled as “3”. Several soils are included within the Low-base Chert Upland Woodland area labeled as “2”, as indicated by the dashed line within the delineation.

Figure 1. Landscape relationships for this ecological site

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Ridge (3) Interfluve
Flooding frequency	None
Ponding frequency	None
Slope	0 – 20 %

Water table depth	40 – 100 cm
Aspect	Aspect is not a significant factor

Climatic features

The Springfield Plain has a continental type of climate marked by strong seasonality. In winter, dry-cold air masses, unchallenged by any topographic barriers, periodically swing south from the northern plains and Canada. If they invade reasonably humid air, snowfall and rainfall result. In summer, moist, warm air masses, equally unchallenged by topographic barriers, swing north from the Gulf of Mexico and can produce abundant amounts of rain, either by fronts or by convectional processes. In some summers, high pressure stagnates over the region, creating extended droughty periods. Spring and fall are transitional seasons when abrupt changes in temperature and precipitation may occur due to successive, fast-moving fronts separating contrasting air masses.

The Springfield Plain experiences few regional differences in climates. The average annual precipitation in this area is 41 to 45 inches. Snow falls nearly every winter, but the snow cover lasts for only a few days. The average annual temperature is about 55 to 58 degrees F. The lower temperatures occur at the higher elevations. Mean July maximum temperatures have a range of only one or two degrees across the area.

Mean annual precipitation varies along a west to east gradient. Seasonal climatic variations are more complex. Seasonality in precipitation is very pronounced due to strong continental influences. June precipitation, for example, averages three to four times greater than January precipitation. Most of the rainfall occurs as high-intensity, convective thunderstorms in summer.

During years when precipitation comes in a fairly normal manner, moisture is stored in the top layers of the soil during the winter and early spring, when evaporation and transpiration are low. During the summer months the loss of water by evaporation and transpiration is high, and if rainfall fails to occur at frequent intervals, drought will result. Drought directly affects plant and animal life by limiting water supplies, especially at times of high temperatures and high evaporation rates.

Superimposed upon the basic MLRA climatic patterns are local topographic influences that create topoclimatic, or microclimatic variations. In regions of appreciable relief, for example, air drainage at nighttime may produce temperatures several degrees lower in valley bottoms than on side slopes. At critical times during the year, this phenomenon may produce later spring or earlier fall freezes in valley bottoms. Deep sinkholes often have a microclimate significantly cooler, moister, and shadier than surrounding surfaces, a phenomenon that may result in a strikingly different ecology. Higher daytime temperatures of bare rock surfaces and higher reflectivity of these unvegetated surfaces may create distinctive environmental niches such as glades and cliffs. Slope orientation is an important topographic influence on climate. Summits and south-and-west-facing slopes are regularly warmer and drier than adjacent north- and east-facing slopes. Finally, the climate within a canopied forest is measurably different from the climate of a more open grassland or savanna areas.

Source: University of Missouri Climate Center - <http://climate.missouri.edu/climate.php>; Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin, United States Department of Agriculture Handbook 296 - <http://soils.usda.gov/survey/geography/mlra/>

Table 3 Representative climatic features

Frost-free period (average)	190 days
Freeze-free period (average)	210 days
Precipitation total (average)	1,190 mm

- (1) SPRINGFIELD [USW00013995], Springfield, MO
- (2) MARSHFIELD [USC00235307], Marshfield, MO
- (3) STOCKTON DAM [USC00238082], Stockton, MO

Influencing water features

This ecological site is not influenced by wetland or riparian water features. This site generates runoff to adjacent, downslope ecological sites. This site does not flood.

The water features of this upland ecological site include evapotranspiration, surface runoff, and drainage. Each water balance component fluctuates to varying extents from year-to-year. Evapotranspiration remains the most constant. Precipitation and drainage are highly variable between years. Seasonal variability differs for each water component. Precipitation generally occurs as single day events. Evapotranspiration is lowest in the winter and peaks in the summer. Water stored as ice and snow decreases drainage and surface runoff rates throughout the winter and increases these fluxes in the spring. The surface runoff pulse is greatly influenced by extreme events. Conversion to cropland or other high intensities land uses tends to increase runoff, but also decreases evapotranspiration. Depending on the situation, this might increase groundwater discharge, and decrease baseflow in receiving streams.

Soil features

These soils are underlain by interbedded shale, siltstone and limestone bedrock between 40 and 70 inches, although the site definition allows for soils as shallow as 20 inches. The subsoils are not low in bases. The soils were formed under woodland vegetation, and have thin, light-colored surface horizons. Parent material is slope alluvium over residuum derived from siltstone and shale. They have silt loam surface horizons that are often gravelly, and clayey subsoils with varying amounts of shale fragments. These soils are not affected by seasonal wetness. Soil series associated with this site include Alsup and Boskydell.

The accompanying picture of the Alsup series shows a thin gravelly silt loam surface horizon over a yellowish brown, clayey subsoil. The olive yellow colors below about 80 cm in this picture are inherited from the shale parent material. Soft shale is below one meter, at the bottom of this picture. Scale is in centimeters. Picture from Henderson (2004).

Figure 6. Alsup series

Table 4. Representative soil features

Parent material	(1) Residuum – shale and siltstone (2) Slope alluvium
Surface texture	(1) Silt loam (2) Gravelly silt loam (3) Very gravelly silty clay loam
Family particle size	(1) Clayey
Drainage class	Moderately well drained
Permeability class	Very slow
Soil depth	100 – 180 cm
Surface fragment cover <=3"	10 – 50 %
Surface fragment cover >3"	0 – 20 %

Available water capacity (0-101.6cm)	7.62 – 15.24 cm
Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	4.5 – 7.8
Subsurface fragment volume <=3" (Depth not specified)	20 – 50 %
Subsurface fragment volume >3" (Depth not specified)	0 – 20 %

Ecological dynamics

Information contained in this section was developed using historical data, professional experience, field reviews, and scientific studies. The information presented is representative of very complex vegetation communities. Key indicator plants, animals and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The Reference Plant Community is not necessarily the management goal. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

The reference plant community is woodland dominated by an overstory of white oak and black oak. This woodland type is moderate in canopy closure (50 to 80 percent), with an open understory and a dense, diverse herbaceous ground flora. Historically, white oak dominated the canopy, along with black oak and occasional hickory, bur oak and post oak. Woodlands are distinguished from forest, by their relatively open understory, and the presence of sun-loving ground flora species including the dominant prairie grasses. Characteristic plants in the ground flora can be used to gauge the restoration potential along with remnant open-grown old-age trees, and tree height growth.

Fire played an important role in the maintenance of these systems. Because Interbedded Sedimentary Upland Woodlands normally occur next to the prairie edge, it is likely that these ecological sites burned at least once every 3 to 5 years. These periodic fires kept woodlands open, removed the litter, and stimulated the growth and flowering of the grasses and forbs. During fire free intervals, woody understory species increased and the herbaceous understory diminished. The return of fire would open the woodlands up again and stimulate the abundant ground flora.

Today, this community has either been cleared and converted to pasture or cropland, or has grown dense in the absence of fire. Most occurrences today exhibit canopy closure of 80 to 100 percent. In addition, the sub-canopy and understory layers are better developed. Black oak and hickory now share dominance with white oak and there are considerable more saplings in the understory. Under these denser, more shaded conditions, the original sun-loving ground flora has diminished in diversity and cover. While some woodland species

persist in the ground flora, many have been replaced by more shade-tolerant species.

In the long term absence of fire, woody species, especially hickory, hophornbeam and gooseberry encroach into these woodlands. Once established, these woody plants can quickly fill the existing understory increasing shade levels greatly diminishing the ground flora. Removal of the younger understory and the application of prescribed fire have proven to be effective management tools.

Uncontrolled domestic grazing has also impacted these communities, further diminishing the diversity of native plants and introducing species that are tolerant of grazing, such as coralberry, gooseberry, and Virginia creeper. Grazed sites also have a more open understory. In addition, soil compaction and soil erosion related to grazing can be a problem and lower site productivity.

Interbedded Sedimentary Upland Woodlands, if managed properly, can be a source for timber products especially white oak. Most areas on this ecological site have been repeatedly logged and high graded. Even-age management, using clearcut, or shelterwood and seed-tree harvest systems without fire will perpetuate the overly dense, shaded conditions. Thinning and/or occasional partial cuts, using an uneven-age management system can provide sunlight to the woodland floor, stimulating native woodland ground flora. However, in the absence of fire and continual cultural treatments, oak sprouting creates a dense stand, again shading out the sun-loving ground flora. Partial cutting and prescribed fire can restore the more open structure and diversity of ground flora species. This type of site with proper management can provide timber products, wildlife habitat, and potential native forage.

A State and Transition Diagram follows. Detailed descriptions of each state, transition, plant community, and pathway follow the model. This model is based on available experimental research, field observations, professional consensus, and interpretations. It is likely to change as knowledge increases.

State and transition model

Figure 7. State and transition diagram for this ecological site

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height M	Canopy Cover (%)	Diameter Cm	Basal Area (square M/hectare)
Tree							
white oak	QUAL	Quercus alba	Native	–	30-60	–	–
black oak	QUVE	Quercus velutina	Native	–	30-60	–	–
post oak	QUST	Quercus stellata	Native	–	10-30	–	–
mockernut hickory	CATO6	Carya tomentosa	Native	–	10-30	–	–
shagbark hickory	CAOV2	Carya ovata	Native	–	10-20	–	–

Table 7. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (m)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
little bluestem	SCSC	Schizachyrium scoparium	Native	–	5-20
Pennsylvania sedge	CAPE6	Carex pensylvanica	Native	–	5-20
hairy woodland brome	BRPU6	Bromus pubescens	Native	–	5-20
oval-leaf sedge	CACE	Carex cephalophora	Native	–	5-20
Virginia wildrye	ELVI3	Elymus virginicus	Native	–	5-20
fuzzy wuzzy sedge	CAHI6	Carex hirsutella	Native	–	5-20
Bosc's panicgrass	DIBO2	Dichanthelium boscii	Native	–	5-20
rock muhly	MUSO	Muhlenbergia sobolifera	Native	–	5-20
Forb/Herb					
American lopseed	PHLE5	Phryma leptostachya	Native	–	5-20
three-lobed violet	VITR2	Viola triloba	Native	–	5-20

Baldwin's ironweed	VEBA	<i>Vernonia baldwinii</i>	Native	–	5-20
hairy sunflower	HEHI2	<i>Helianthus hirsutus</i>	Native	–	5-20
eastern purple coneflower	ECPU	<i>Echinacea purpurea</i>	Native	–	5-20
Virginia spiderwort	TRVI	<i>Tradescantia virginiana</i>	Native	–	5-20
elmleaf goldenrod	SOUL2	<i>Solidago ulmifolia</i>	Native	–	5-20
manyray aster	SYAN2	<i>Symphyotrichum anomalum</i>	Native	–	5-20
pointedleaf ticktrefoil	DEGL5	<i>Desmodium glutinosum</i>	Native	–	5-20
smooth small-leaf ticktrefoil	DEMA2	<i>Desmodium marilandicum</i>	Native	–	5-20
fourleaf milkweed	ASQU	<i>Asclepias quadrifolia</i>	Native	–	5-20
eastern beebalm	MOBR2	<i>Monarda bradburiana</i>	Native	–	5-20
nakedflower ticktrefoil	DENU4	<i>Desmodium nudiflorum</i>	Native	–	5-20
Shrub/Subshrub					
fragrant sumac	RHAR4	<i>Rhus aromatica</i>	Native	–	5-20
American hazelnut	COAM3	<i>Corylus americana</i>	Native	–	5-20
New Jersey tea	CEAM	<i>Ceanothus americanus</i>	Native	–	5-20

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 13. Community 5.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 14. Community 6.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

Wildlife (MDC 2006): Wild turkey, white-tailed deer, and eastern gray squirrel depend on hard and soft mast food sources and are typical upland game species of this type. Oaks provide hard mast; scattered shrubs provide soft mast; native legumes provide high-quality wildlife food. Sedges and native cool-season grasses provide green browse; patchy native warm-season grasses provide cover and nesting habitat; and a diversity of forbs provides a diversity and abundance of insects. Post-burn areas can provide temporary bare-ground – herbaceous cover habitat important for turkey poults and quail chicks. Bird species associated with mature communities include Indigo Bunting, Red-headed Woodpecker, Eastern Bluebird, Northern Bobwhite, Eastern Wood-Pewee, Broad-winged Hawk, Great-Crested Flycatcher, Summer Tanager, and Red-eyed Vireo. Reptile and amphibian species associated with this site include tiger salamander, small-mouthed salamander, ornate box turtle, northern fence lizard, five-lined skink, broad-headed skink, flat-headed snake, and rough earth snake.

Other information

Forestry (NRCS 2002; 2014): Management: Field measured site index values for oak range from 51 for post oak, 62 for red oak and 54 for white oak. Timber management opportunities are fair to good. Create group openings of at least 2 acres. Large clearcuts should be minimized if possible to reduce impacts on wildlife and aesthetics. Uneven-aged management using single tree selection or group selection cuttings of ½ to 1 acre are other options that can be used if clear cutting is not desired or warranted. These sites respond well to prescribed fire as a management tool. Limitations: Clay in lower portion of soil profile; Exposed clayey soils have reduced traction and compact easily when wet. Unsurfaced roads and skid trails may be impassable during rainy periods. Restrict activities to dry periods or surfaced areas. Seedling mortality may be high during the summer because of lack of adequate soil moisture, especially on south facing slopes. Erosion is a hazard when slopes exceed 15 percent. On steep slopes greater than 35 percent, traction problems increase and equipment use is not recommended.

Inventory data references

Potential Reference Sites: Interbedded Sedimentary Upland Woodland Plot STLACE02 – Boskydell soil Located in Stockton Lake COE/CA, Cedar County, MO Latitude: 37.606618 Longitude: -93.702388

Other references

Anderson, R.C. 1990. The historic role of fire in North American grasslands. Pp. 8-18 in S.L. Collins and L.L. Wallace (eds.). Fire in North American tallgrass prairies. University of Oklahoma Press, Norman.

Batek, M.J., A.J. Rebertus, W.A. Schroeder, T.L. Haithcoat, E. Compas, and R.P. Guyette. 1999. Reconstruction of early nineteenth-century vegetation and fire regimes in the Missouri Ozarks. *Journal of Biogeography* 26:397-412.

Dodd, Jerry A. 1990. Soil Survey of Webster County, Missouri. U.S. Dept. of Agric. Soil Conservation Service.

Harlan, J.D., T.A. Nigh and W.A. Schroeder. 2001. The Missouri original General Land Office survey notes project. University of Missouri, Columbia.

Henderson, Richard L. 2004. Soil Survey of Cedar County, Missouri. U.S. Dept. of Agric. Natural Resources Conservation Service.

Ladd, D. 1991. Reexamination of the role of fire in Missouri oak woodlands. Pp. 67-80 in G.V. Brown, James K.; Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Missouri Department of Conservation. 2006. Missouri Forest and Woodland Community Profiles. Missouri Department of Conservation, Jefferson City, Missouri.

NatureServe. 2010. Vegetation Associations of Missouri (revised). NatureServe, St. Paul, Minnesota.

Natural Resources Conservation Service. 2002. Woodland Suitability Groups. Missouri FOTG, Section II, Soil Interpretations and Reports. 30 pgs.

Natural Resources Conservation Service. Site Index Reports. Accessed May 2014. https://esi.sc.egov.usda.gov/ESI_Forestland/pgFSWelcome.aspx

Nelson, Paul W. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.

Nigh, Timothy A., and Walter A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, Missouri.

Schoolcraft, H.R. 1821. Journal of a tour into the interior of Missouri and Arkansas from Potosi, or Mine a Burton, in Missouri territory, in a southwest direction, toward the Rocky Mountains: performed in the years 1818 and 1819. Richard Phillips and Company, London.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. 682 pgs.

Contributors

Doug Wallace
Fred Young

Approval

Acknowledgments

Missouri Department of Conservation and Missouri Department of Natural Resources personnel provided significant and helpful field and technical support in the development of this ecological site.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/16/2020
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
