

Ecological site F126XY003OH

Moist Ridge

Last updated: 9/27/2024
Accessed: 10/05/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 126X—Central Allegheny Plateau

These sites generally occur on summits with soils that are not well drained. The MAAT > 45 degree F. This ecosite is found in Alleghany plateau in MLRA 126. Steep slopes are dominant, but level to gently rolling plateau remnants exist. The area is dominantly forest, containing large blocks of state forest, game lands, and national forest. Less than one-tenth of the MLRA consists of urban areas

Classification relationships

USDA-NRCS (USDA 2006): Land Resource Region (LRR): N—East and Central Farming and Forest Region Major Land Resource Area (MLRA): 126—Central Allegheny Plateau USDA-FS (Cleland et al. 2007) Province: 221 - Eastern Broadleaf Province Section: 221E - Southern Unglaciaded Allegheny Plateau Subsection: 221Ea - Pittsburgh Low Plateau 221Eb - Teays Plateau 221Ec - Ohio Valley Lowland 221Ed - East Hocking Plateau This site crosswalks to Landfire biophysical setting (BpS) Northeastern Interior Dry-Mesic Oak Forest NatureServe's description (2007) for the equivalent ecological system CES202.592 -- Northeastern Interior Dry-Mesic Oak Forest notes this system is found from central New England west through OH and PA and south to VA. Component Associations Association Unique ID Association Name CEGLO02059 Quercus alba - Quercus rubra - Quercus prinus - Acer saccharum / Lindera benzoin Forest CEGLO02067 Quercus alba - Quercus rubra - Carya (alba, ovata) / Cornus florida Acidic Forest CEGLO02068 Quercus alba - Quercus rubra - Carya ovata Glaciaded Forest CEGLO04793 Quercus muehlenbergii - Quercus (alba, rubra) - Carya cordiformis / Viburnum prunifolium Forest CEGLO06057 Quercus prinus - Quercus rubra / Hamamelis virginiana Forest CEGLO06216 Quercus alba - Carya glabra - Fraxinus americana / Cercis canadensis / Muhlenbergia sobolifera - Elymus hystrix Forest CEGLO06241 Quercus bicolor / Vaccinium corymbosum / Carex stipata Forest CEGLO06293 Pinus strobus - Quercus (rubra, velutina) - Fagus grandifolia Forest CEGLO06301 Quercus rubra - Carya (glabra, ovata) / Ostrya virginiana / Carex lucorum Forest CEGLO06304 Liriodendron tulipifera - Pinus strobus - Tsuga canadensis - Quercus (rubra, alba) / Polystichum acrostichoides Forest CEGLO06336 Quercus (alba, rubra, velutina) - Carya spp. / Viburnum acerifolium Forest CEGLO06454 Quercus (rubra, velutina, alba) - Betula lenta - (Pinus strobus) Forest CEGLO06921 Fagus grandifolia - Betula lenta - Quercus (alba, rubra) / Carpinus caroliniana Forest CEGLO08475 Quercus alba - Quercus rubra - Carya alba / Cornus florida / Vaccinium stamineum / Desmodium nudiflorum Piedmont Forest CEGLO08514 Quercus rubra - Quercus prinus - Carya ovalis / (Cercis canadensis) / Solidago (caesia, curtisii) Forest CEGLO08516 Quercus prinus - Quercus rubra - Carya ovalis / Carex pensylvanica - (Calamagrostis porteri) Forest CEGLO08517 Quercus rubra - Acer saccharum / Ostrya virginiana / Cardamine concatenata Forest CEGLO08518 Quercus rubra - Carya (ovata, ovalis) - Fraxinus americana / Actaea racemosa - Hydrophyllum virginianum Forest CEGLO08528 Tilia americana - Fraxinus americana / Acer pensylvanicum - Ostrya virginiana / Parthenocissus quinquefolia - Impatiens pallida Woodland See also NatureServe's description (2007): Allegheny-Cumberland Dry Oak Forest and Woodland CES202.359 Component Associations Association Unique ID Association Name CEGLO04761 Pinus echinata - Quercus prinus - Quercus (coccinea, velutina) Forest CEGLO04786 Quercus prinus - Carya (alba, glabra, ovata) / Juniperus virginiana var. virginiana Forest CEGLO05023 Quercus prinus - Quercus (alba, coccinea, velutina) / Viburnum acerifolium - (Kalmia latifolia) Forest CEGLO06557 Pinus rigida - Quercus coccinea / Vaccinium angustifolium Woodland CEGLO07119 Pinus virginiana - Pinus (rigida, echinata) - (Quercus prinus) / Vaccinium pallidum Forest CEGLO07231 Quercus alba - Quercus velutina - Carya (ovata, alba, glabra) - Pinus sp. Forest CEGLO07240 Quercus alba - Quercus rubra - Carya ovata / Cercis canadensis - Juniperus virginiana var. virginiana Forest CEGLO07244 Quercus falcata - Quercus alba - Carya alba / Oxydendrum arboreum / Vaccinium stamineum Forest CEGLO07247 Quercus falcata - Quercus (coccinea, stellata) / Vaccinium (pallidum, stamineum) Forest CEGLO07261 Quercus prinus - Carya spp. - Quercus velutina / Vaccinium arboreum / Iris verna var. smalliana Forest CEGLO07269 Quercus prinus - Quercus rubra - Carya (ovata, glabra) - Pinus virginiana Forest CEGLO07517 Pinus strobus - Quercus alba - (Carya alba) / Gaylussacia ursina Forest CEGLO07519 Pinus strobus - Quercus (coccinea, prinus) / (Gaylussacia ursina, Vaccinium stamineum) Forest CEGLO07700 Quercus prinus - Quercus spp. / Vaccinium arboreum - (Kalmia latifolia, Styrax grandifolius) Forest CEGLO07795 Quercus alba - Carya alba - (Quercus velutina) / Desmodium nudiflorum - (Carex picta) Forest CEGLO08406 Quercus stellata - Pinus virginiana / (Schizachyrium scoparium, Piptochaetium avenaceum) Woodland CEGLO08430 Quercus alba - (Quercus prinus) / (Hydrangea quercifolia) - Viburnum acerifolium / Carex picta - Piptochaetium avenaceum Forest CEGLO08431 Quercus prinus - (Quercus coccinea) / Carya pallida / Vaccinium arboreum

- Vaccinium pallidum Forest CEG008443 Quercus alba - Quercus stellata / Ostrya virginiana - Acer barbatum / Chasmanthium sessiliflorum Forest CEG008521 Quercus alba - Quercus (coccinea, velutina, prinus) / Gaylussacia baccata Forest CEG008567 Quercus alba - Quercus falcata / Vaccinium (arboreum, hirsutum, pallidum) Forest

Ecological site concept

From Landfire <http://www.landfire.gov/index.php>: This ecological section comprises part of the Appalachian Plateaus geomorphic province (USDA Forest Service 1995). It is a maturely dissected plateau characterized by high hills, sharp ridges, and narrow valleys. Bedrock is overlain by Quaternary residuum on the ridges and hilltops, colluvium on the slopes, and either or both alluvium and Pleistocene lacustrine materials in the valleys. Udalfs, Udults, and Ochrepts dominate, in combination with mesic soil temperature regime, an udic soil moisture regime, and mixed or illitic mineralogy. Soils formed in parent materials are divided into five groups: residual material, which developed in place by the weathering of underlying bedrock; colluvial material which weathered from bedrock strata transported by water and gravity to the lower slopes; alluvium, lacustrine sediments and outwash deposited by water; and loess deposited by wind. Precipitation averages 35 to 45in (900 to 1,150mm); it occurs mainly during summer, winter, and spring. Rain on snow is common during winter and early spring. Summers are dry with low humidity. Temperature averages 52 degrees F (11 degrees C). The growing season is 120 to 180 days. This ecological section is characterized by a relatively high density of streams, with gradients ranging from high, steep headwaters streams to low gradient rivers that flow into the OH River. NatureServe (2007) provides the following description pertaining to this system type. These oak-dominated forests are one of the matrix forest systems in the northeastern and north-central US. Occurring in dry-mesic settings, they are typically closed-canopy forests, though there may be areas of patchy-canopy woodlands. They cover large expanses at low to mid elevations, where the topography is flat to gently rolling, occasionally steep. Soils are acidic and relatively infertile but not strongly xeric.

Table 1. Dominant plant species

Tree	(1) <i>Quercus Alba</i> (2) <i>Quercus Rubra</i>
Shrub	(1) <i>Cornus Florida</i> (2) <i>Viburnum acerifolium</i>
Herbaceous	(1) <i>Carex</i>

Physiographic features

Residuum of not coarse or skeletal soils.

Table 2. Representative physiographic features

Landforms	(1) Plateau > Hill (2) Plateau > Ridge
Runoff class	Low to very high
Elevation	240 – 460 m
Slope	0 – 30 %
Water table depth	30 – 180 cm
Aspect	Aspect is not a significant factor

Climatic features

The regional climate of the unglaciated Central Allegheny Plateau is predominately a humid continental climate grading at the extreme southwestern corner a to humid temperate climate with hot summers and cool winters (Beck et al., 2018; Bailey, 2014). However, the local climate is highly influenced by the dissected terrain, where climatic variations may be greater at the local scale, e.g., cooler temperatures and shorter growing season at higher elevations and more northerly latitudes. High-intensity, convective thunderstorms are common in summer. Winter precipitation is mostly snow.

Climate change is occurring, and the resiliency of any ecological site will depend upon the direct and indirect effects upon component species and shifting atmospheric and soil conditions.

Table 3 Representative climatic features

Frost-free period (characteristic range)	140-170 days
Freeze-free period (characteristic range)	170-200 days
Precipitation total (characteristic range)	1,020-1,090 mm
Frost-free period (actual range)	130-170 days
Freeze-free period (actual range)	160-200 days
Precipitation total (actual range)	990-1,140 mm
Frost-free period (average)	150 days
Freeze-free period (average)	180 days
Precipitation total (average)	1,070 mm

- (1) WHEELING [USC00469482], Wheeling, WV
- (2) SENECAVILLE LAKE [USC00337559], Senecaville, OH
- (3) CALDWELL 3 SE [USC00331175], Caldwell, OH
- (4) HANNIBAL L&D [USC00333500], New Martinsville, OH
- (5) WAYNESBURG 1 E [USC00369367], Waynesburg, PA
- (6) WEST UNION 2 [USC00469458], West Union, WV
- (7) PARKERSBURG [USW00013867], Parkersburg, WV
- (8) WINFIELD LOCKS [USC00469683], Red House, WV
- (9) PEA RIDGE PSD [USC00466912], Huntington, WV
- (10) HUNTINGTON SEWAGE PLT [USC00464397], Kenova, WV
- (11) LOYALHANNA LAKE [USC00365212], New Alexandria, PA
- (12) PITTSBURGH INTL AP [USW00094823], Coraopolis, PA

Influencing water features

Water features are not typically associated with this ecological site, but can be incidental.

Wetland description

Soil features

Soils are acidic and relatively infertile but not strongly xeric.

Table 4. Representative soil features

Parent material	(1) Residuum
Drainage class	Somewhat poorly drained to moderately well drained
Permeability class	Very slow to moderately slow
Depth to restrictive layer	60 – 180 cm
Surface fragment cover ≤3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (Depth not specified)	5.08 – 20.32 cm
Soil reaction (1:1 water) (Depth not specified)	3.8 – 7.8
Subsurface fragment volume ≤3" (Depth not specified)	0 – 40 %
Subsurface fragment volume >3" (Depth not specified)	0 – 50 %

Ecological dynamics

Information contained in this section was adapted from several sources. The information presented is representative of very complex vegetation communities. Key indicator plants, animals and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The reference plant community is not necessarily the management goal. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

From Landfire <http://www.landfire.gov/index.php>:

This ecological section comprises part of the Appalachian Plateaus geomorphic province. It is a maturely dissected plateau characterized by high hills, sharp ridges, and narrow valleys. Bedrock is overlain by Quaternary residuum on the ridges and hilltops, colluvium on the slopes, and either or both alluvium and Pleistocene lacustrine materials in the valleys. Udalfs, Udufts, and Ochrepts dominate, in

combination with mesic soil temperature regime, an udic soil moisture regime, and mixed or illitic mineralogy. Soils formed in parent materials are divided into five groups:residual material, which developed in place by the weathering of underlying bedrock; colluvial material which weathered from bedrock strata transported by water and gravity to the lower slopes; alluvium, lacustrine sediments and outwash deposited by water; and loess deposited by wind. Precipitation averages 35 to 45in (900 to 1,150mm0150mm); it occurs mainly during summer, winter, and spring. Rain on snow is common during winter and early spring. Summers are dry with low humidity. Temperature averages 52 degrees F (11 degrees C). The growing season is 120 to 180 days. This ecological section is characterized by a relatively high density of streams, with gradients ranging from high, steep headwaters streams to low gradient rivers that flow into the OH River.

NatureServe (2007) provides the following description pertaining to this system type. These oak-dominated forests are one of the matrix forest systems in the northeastern and north-central US Occurring in dry-mesic settings, they are typically closed-canopy forests, though there may be areas of patchy-canopy woodlands. They cover large expanses at low to mid elevations, where the topography is flat to gently rolling, occasionally steep. Soils are acidic and relatively infertile but not strongly xeric.

Vegetation Description

The vegetation consisted of forests dominated by oaks of dry-mesic conditions, especially white oak (*Quercus alba*) and red oak (*Quercus rubra*), and, on drier sites, chestnut oak (*Quercus prinus*), black oak (*Quercus velutina*), and scarlet oak (*Quercus coccinea*). Scarlet oak is absent at the northern edge of the system range. Along with oaks are varying amounts of hickory (*Carya* spp.), red maple (*Acer rubrum*), and other species such as white pine (*Pinus strobus*) and white ash (*Fraxinus americana*). American chestnut (*Castanea dentata*) was a prominent tree in these forests before chestnut blight eradicated it as a canopy constituent (NatureServe 2007). Common shrubs include mountain laurel (*Kalmia* spp.), greenbriar (*Smilax* spp.), blueberries (*Vaccinium* spp.), and huckleberries (*Gaylussacia* spp.). In the Ridge and Valley region, bear oak is an important shrub component. Herbs, forbs, and ferns are usually sparse to moderate in density. Areas experiencing frequent fire had a greater abundance of grasses and sedges.

State and transition model

Figure 7. State and Transition

Figure 8. Legend

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Site Development and Testing Plan Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

Other references

Anderson, D. M. 1982. Plant Communities of Ohio: A Preliminary Classification. Division of Natural Areas and Preserves, Ohio Department of Natural Resources, Columbus, OH. (https://www.lm.doe.gov/cercla/documents/fernald_docs/cat/112509.pdf).

Apsley, D., and B.C. McCarthy. 2004. White-tailed deer herbivory on forest regeneration following fire and thinning treatments in southern Ohio mixed oak forests. P. 461–471. In: Yaussy, D.A., D.M. Hix, R.P. Long, and P.C. Goebel (eds.) Proceedings, 14th Central Hardwood Forest Conference, Wooster, OH. 16-19 March 2004. Gen. Tech. Rep. NE-316. USDA Forest Service, Northeastern Research Station, Newtown Square, PA.

Bailey, R. 2014. Ecoregions: the ecosystem geography of the oceans and continents. 2nd ed. New York, NY: Springer-Verlag.

Beck, H.E., N.E. Zimmermann, T.R. McVicar, N. Vergopolan, A. Berg, E.F. Wood. 2018. Present and future Köppen-Geiger climate classification maps at 1-km resolution. Scientific Data 5(1):1-12.

Butler, P.R., L. Iverson, F.R. Thompson, L. Brandt, S. Handler, M. Janowiak, P.D. Shannon, C. Swanston, K. Karriker, J. Bartig, and S. Connolly. 2015. Central Appalachians Forest Ecosystem Vulnerability Assessment and Synthesis: a Report From The Central Appalachians Climate Change Response Framework Project. Gen. Tech. Rep. NRS-146, US Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C.A. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. [Map. presentation scale 1:3,500,000, colored; A.M. Sloan, cartographer] Gen. Tech. Report WO-76D. U.S. Department of Agriculture, Forest Service, Washington, DC. (<https://www.fs.fed.us/research/publications/misc/73326-wo-gtr-76d-cleland2007.pdf>)

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, and K. Snow. 2003. Ecological Systems of the United States: A Working Classification of US Terrestrial Systems. NatureServe, Arlington, VA. (https://www.natureserve.org/sites/default/files/pcom_2003_ecol_systems_us.pdf).

FGDC (Federal Geographic Data Committee). 2008. National Vegetation Classification Standard, Version 2. VGDC?STD?005?2008 (Version 2). FGDC Vegetation Subcommittee, Reston, Virginia. (https://www.fgdc.gov/standards/projects/vegetation/NVCS_V2_FINAL_2008-02.pdf).

Lafon, C.W., A.T. Naito, H.D. Grissino-Mayer, S.P. Horn, and T.A. Waldrop. 2017. Fire History of the Appalachian Region: a Review and Synthesis. Gen. Tech. Rep. SRS-219., U.S. Department of Agriculture, Forest Service, Southern Research Station, Asheville, NC.

NatureServe 2007. NatureServe Explorer: An Online Encyclopedia of Life [web application]. NatureServe, Arlington, VA. Available: <http://explorer.natureserve.org> (Accessed: April 2020).

Nowacki, G.J. and M.D. Abrams. 2008. The demise of fire and “mesophication” of forests in the eastern United States. Bioscience 58(2):123–138.

Ohio Division of Wildlife. 2015. Ohio's State Wildlife Action Plan. Columbus, Ohio, USA. (https://ohiodnr.gov/static/documents/wildlife/wildlife-management/OH_SWAP_2015.pdf).

Soil Survey Staff-USDA-NRCS [United States Department of Agriculture, Natural Resources Conservation Service] 2016. National Soils Information Service (NASIS Data Model Version 7.3.4) Lincoln, NE. Available description: https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/soils/survey/tools/?cid=nrcs142p2_053552 (Accessed January 2020).

USDA-NRCS [United States Department of Agriculture, Natural Resources Conservation Service]. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. (https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051845.pdf).

USNVC [United States National Vegetation Classification]. 2019. United States National Vegetation Classification Database, V2.03. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. <http://usnvc.org> (accessed April 2020).

Zimmerman, E., T. Davis, G. Podniesinski, M. Furedi, J. McPherson, S. Seymour, B. Eichelberger, N. Dewar, J. Wagner, and J. Fike (editors). 2012. Terrestrial and Palustrine Plant Communities of Pennsylvania, 2nd Edition. Pennsylvania Natural Heritage Program, Pennsylvania Department of Conservation and Natural Resources, Harrisburg, PA.

Contributors

Jason Teets

Approval

Greg Schmidt, 9/27/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
--------------------------	--

Contact for lead author	
Date	10/05/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
