

Ecological site F134XY306LA

West Central Well Drained Loamy Ridge

Last updated: 3/24/2025

Accessed: 09/19/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 134X–Southern Mississippi Valley Loess

MLRA 134, Southern Mississippi Valley Loess, is in Mississippi (39 percent), Tennessee (23 percent), Louisiana (15 percent), Arkansas (11 percent), Kentucky (9 percent), Missouri (2 percent), and Illinois (1 percent). It makes up about 26,520 square miles (68,715 square kilometers). The northern part of the area includes Paducah and Murray, Kentucky; Paragould, Jonesboro, and Forrest City, Arkansas; and Memphis, Dyersburg, Bartlett, and Germantown, Tennessee. The southern part includes Yazoo City, Clinton, and Jackson, Mississippi, and Baton Rouge, Opelousas, Lafayette, and New Iberia, Louisiana. This portion is the central western part of the MLRA in Louisiana and Arkansas. It is in the Macon Ridge Section of the EPA Ecoregions in sub-section 73j. The dissected plains in this MLRA have a loess mantle that is thick at the valley wall and thins rapidly as distance from the valley wall increases. This portion of the MLRA is distinct from other portions of the MLRA because of the influences of the Mississippi River and its series of entrenchments and adjacent old channels of the Arkansas River, such channels as Bayou Bartholomew, Bayou Bonne Idee, Boeuf River, and segments of the Ouachita River. The Macon Ridge has been inhabited prior to European Settlement, Poverty Point is located on the east central portion of the Macon Ridge and has earthworks dating back to 1700-1100 BC.

Classification relationships

Major Land Resource Area (MLRA) and Land Resource Unit (LRU) (USDA-Natural Resources Conservation Service, 2006) The Natural Communities of Louisiana - (Louisiana Natural Heritage Program - Louisiana Department of Wildlife and Fisheries) EPA Level IV Ecoregion

Ecological site concept

Very deep, well drained, moderately permeable soils that formed in loamy sediments on stream braid bars within the Macon Ridge Loess with little to no loess influence. These sites are found on long narrow slight rises on terraces where thin loess deposits are mixed throughout with loamy underlying braided stream material. Slopes are 5% or less. Soils limitations could include high exchangeable Al (Aluminum). This soil chemistry limitation is significant because, Aluminum (Al) toxicity limits production on strongly acidic soils, pH values at or below 5, toxic forms of Al solubilized into the soil solution, inhibiting root growth and function (Kochian, 2005). Common species on this site are Cherrybark oak, Loblolly pine, Pecan, Eastern Cottonwood and Sweetgum.

Associated sites

F134XY304LA	West Central Somewhat Poorly Drained Flats The Somewhat Poorly Drained Flats sites will be at a lower level on the landscape.
F134XY305LA	West Central Somewhat Poorly Drained Loess Terrace The Somewhat Poorly Drained Loess Terrace sites will be at a lower level on the landscape.

F134XY307LA	West Central Loess Ridge The Loess Ridge site will be found at a higher landscape position.
--------------------	---

Similar sites

F134XY307LA	West Central Loess Ridge The West Central Loess Ridge is found at higher landscape position.
--------------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

Macon Ridge in extreme northeastern Louisiana and southeastern Arkansas is a 135-mile long prominent ridge that lies between the Boeuf and Tensas Basins (Saucier, 1994). Macon Ridge is a terrace that features level to gently undulating topography with steep scarps of uplands, floodplains, depressions, and drainageways. The entire area is located within the Mississippi Alluvial Valley section of the Coastal Plain Province of the Atlantic Plain. It reaches a maximum width of about 25 miles in northeastern Louisiana about 30 miles north of Sicily Island (Saucier, 1994). Elevation ranges from 50 to 150 feet (15 to 46 m). The ridge is consistently higher on its eastern side where elevations are 20 to 30 feet higher than in the adjacent Tensas Basin (floodplain). It is bounded on the eastern edge by Bayou Macon. On the western side, bounded by the Boeuf River, elevations of the ridge are approximately the same as those in the Boeuf Basin, and it is sometimes difficult to distinguish the two at the surface (Saucier, 1994). Both Bayou Macon and Boeuf River are underfit streams occupying ancient Arkansas River meanders. The entire Macon Ridge is underlain by Pleistocene-aged loamy and clayey braided stream alluvium from the "old" Arkansas River. Macon Ridge consists almost entirely of Early Wisconsin age glacial outwash and is a continuation of the valley train in the Western Lowlands (Saucier, 1994). The area mantled by loess on the eastern edge of the terrace rises 10 to 30 feet above the floodplains. The loess thins toward the west, and elevation decreases. The loess in the western part of Macon Ridge contains small mixtures of the older underlying braided-stream terrace alluvium, and in even lower elevations, the loess contains mixtures of recent clayey alluvium or is buried completely beneath recent alluvium (Allen, 1993).

This PES occurs on silty and loamy elongated, narrow convex ridges on nearly level to gently sloping stream terraces on the loess-mantled Macon Ridge in Louisiana. Slopes are nearly level to gently sloping (0 to 5 percent). These sites include ridges of stream terraces of the Boeuf River and other former channels and distributaries of the Arkansas River such as: Big and Little Colewa Creeks, Big Creek, Bee Bayou, Turkey Creek, Muddy Bayou, and Pine Bayou.

Table 2. Representative physiographic features

Landforms	(1) Ridge (2) Terrace
Flooding frequency	None
Ponding frequency	None
Elevation	20 – 30 m

Slope	0 – 10 %
Ponding depth	0 cm
Water table depth	180 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate of North East Louisiana and South East Arkansas is warm and humid with a monthly precipitation that is well distributed throughout the year. The monthly precipitation mean is between 2.9 and 5.3 inches, with the lowest rainfall occurring from June through November. The following climatic data are averages from the three weather stations listed below. Temperature and precipitation may vary considerably from that listed for each month. Site specific weather data should be used for land management decisions. For site specific weather conditions, obtain data from a weather station close to the site.

Table 3 Representative climatic features

Frost-free period (average)	230 days
Freeze-free period (average)	260 days
Precipitation total (average)	1,500 mm

- (1) WINNSBORO 5 SSE [USC00169806], Winnsboro, LA
- (2) EUDORA [USC00032355], Eudora, AR
- (3) RAYVILLE [USC00167691], Rayville, LA

Influencing water features

This site is well drained, sheds runoff and the soils are moderately permeable. There is limited amounts of water that influence the site other than the rainfall that is intercepted and allowed to percolate into the soil profile.

Soil features

Soils are well drained, Ultic Hapludalfs (Dexter and Liddieville). These soils formed from mixed loess and loamy and sandy sediments of late Pleistocene Age. Slopes range from 0 to 5 percent. These very deep, moderately permeable soils are found on elongated, narrow convex ridges on nearly level to gently sloping uplands and stream terraces. These soils are not considered hydric and have a water table deeper than 6 feet. These soils are not subject to flooding.

Table 4. Representative soil features

Surface texture	(1) Silt loam
Family particle size	(1) Loamy
Drainage class	Well drained

Permeability class	Moderate
Soil depth	150 – 200 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	0.2 – 0.61 cm
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	4.5 – 7.3
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The pre settlement plant community of this site would have been dominated by hardwood species. Some trees can be uprooted by climatic events, such as strong winds or floods. With these events, openings in the canopy can occur which will set back succession and allow herbaceous and woody shrub species to colonize, these low stature communities are generally short lived and the upper canopy will close as tall growing trees mature. There is generally an age gradient within a forest stand from the herbaceous openings to mature hardwoods.

This site has been altered by human activity and is utilized for multiple production systems such Cropland, Pasture and Tree Farms. Within the alternative uses of the site the transitions will be very similar and require the input of resources such as installation of infrastructure needs and establishment of the desired species.

This PES occurs on Well drained ridges in braided stream terraces of the Loess Plain. Located on the Macon Ridge in Louisiana and Arkansas.

State and transition model

Figure 5. 134XY306 West Central Well Drained Loamy Ridge PES

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
Grass/Grasslike					
1	Warm Season Perennials			2914-14571	
	Bermudagrass	CYDA	<i>Cynodon dactylon</i>	2914-14571	–
	bahiagrass	PANO2	<i>Paspalum notatum</i>	3699-7286	–

Table 8. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Animal community

.

Hydrological functions

.

Recreational uses

.

Wood products

.

Other products

.

Other information

.

Other references

Allen, T. E. (1993), USDA-NRCS Richland Parish Soil Survey Report

Autin, W. J., Burns, S. F., Miller, B. J., Saucier, R. T., & Snead, J. I. (1991). Quaternary geology of the lower Mississippi Valley. The Geology of North America, 2, 547-582.

Cowardin, L. M., Carter, V., Golet, F. C., & LaRoe, E. T. (1979). Classification of wetlands and deepwater habitats of the United States. US Fish and Wildlife Service FWS/OBS, 79(31), 131.

Daigle, J.J., Griffith, G.E., Omernik, J.M., Faulkner, P.L., McCulloh, R.P., Handley, L.R., Smith, L.M., and Chapman, S.S. (2006), Ecoregions of Louisiana (color poster with map, descriptive text, summary tables, and photographs): Reston, Virginia, U.S. Geological Survey (map scale 1:1,000,000).

Emerson, F. V. (1918). Loess-depositing winds in Louisiana. *The Journal of Geology*, 26(6), 532-541.

Ezell, A. W., & Hodges, J. D. (1995). *Bottomland hardwood management: Species Site Relationships*. MSU Extension Service Publication 2004.

Frost, C. C. Presettlement Fire frequency regimes of the United States: A First approximation. *Fire in ecosystem management: Shifting the paradigm from suppression to prescription*, ed. TL Pruden and LA Brennan, 70-81.

Guyette, R. P., Stambaugh, M. C., Dey, D. C., & Muzika, R. M. (2012). Predicting fire frequency with chemistry and climate. *Ecosystems*, 15 (2), 322-335.

Heinrich, P. V., (2008), *Loess Map of LA*, Louisiana Geological Survey

Kochian, L. V., Pineros, M. A., & Hoekenga, O. A. (2005). The physiology, genetics and molecular biology of plant aluminum resistance and toxicity. In *Root Physiology: From Gene to Function* (pp. 175-195). Springer Netherlands.

Latimore S. (1996). *THE RARE AND SENSITIVE NATURAL WETLAND PLANT COMMUNITIES OF INTERIOR LOUISIANA*. Louisiana Natural Heritage Program, Louisiana Department of Wildlife & Fisheries, Baton Rouge, Louisiana.

Louisiana Natural Heritage Program, Louisiana Department of Wildlife & Fisheries, (2009) *The Natural Communities of Louisiana* McCraw, David J., and Whitney J. Autin. *Lower Mississippi Valley, Loess: A Field Guide*. Inqua Commission on Loess, 1989.

Miller, B. J., Day, W. J., & Schumacher, B. A. (1986). *Loesses and loess-derived soils in the Lower Mississippi Valley*. Guidebook for soils- geomorphology tour.

Miller, B. J., Lewis, G. C., Alford, J. J. & Day, W. J. (1984) *Loesses in Louisiana and at Vicksburg, Mississippi*. Guidebook for Friends of the Pleistocene Field Trip.

Muery, E. (1998), *ANALYSIS OF PRESETTLEMENT NATURAL PLANT COMMUNITY TYPES OF THE MACON RIDGE OF LOUISIANA*.

Pettry, D. E., & Switzer, R. E. (1998). *Sodium soils in Mississippi*.

Saucier, R. T. (1994). *Geomorphology and Quarternary Geologic History of the Lower Mississippi Valley*. Volumes 1 and 2. ARMY ENGINEER WATERWAYS EXPERIMENT STATION VICKSBURG MS GEOTECHNICAL LAB.

Schumacher, B. A., B. J. Miller, and W. J. Day. "A chronotoposequence of soils developed in loess in central Louisiana." *Soil Science Society of America Journal* 51.4 (1987): 1005-1010.

Theriot, R. F. (1992). *Flood tolerance of plant species in bottomland forests of the southeastern United States*.

United States Salinity Laboratory Staff, USA, USDA (1954), *Diagnosis and improvement of saline and alkali soils*, USDA Agriculture Handbook 60, 1954, 160 pp.

USDA Agriculture Handbook 296. (2006). <http://soils.usda.gov/MLRAExplorer>. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

USDA Natural Resources Conservation Service. *Published Soil Surveys from Catahoula, Franklin, Richland and West Carroll Parishes*. Various publication dates.

USDA Natural Resources Conservation Service. *Web Soil Survey*, <http://websoilsurvey.nrcs.usda.gov/app>. USDA NRCS Soil Survey Division. Washington, DC. 2008.

Contributors

D Charles Stemmans II
Wayne Roberts
Dwayne Rice
Rachel Stout Evans

Approval

Matthew Duvall, 3/24/2025

Acknowledgments

The Macon Ridge Technical Team did an outstanding job of utilizing existing data and knowledge to develop site concepts. They also provided the needed sections of the descriptions, review and comments of them.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/19/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
