

Ecological site F140XY002NY

Frigid Steep Shallow Till Uplands

Last updated: 10/01/2024
Accessed: 09/25/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 140X–Glaciated Allegheny Plateau and Catskill Mountains

This area is primarily in the Southern New York Section of the Appalachian Plateaus Province of the Appalachian Highlands. The top of the dissected plateau in this MLRA is broad and is nearly level to moderately sloping. The narrow valleys have steep walls and smooth floors. The Catskills in the east have steep slopes. Elevation is typically 650 to 1,000 feet on valley floors; 1,650 to 2,000 feet on the plateau surface; and 3,600 feet or more in parts of the Catskills. The average annual precipitation in most of this area is 30 to 45 inches. Rainfall occurs as high-intensity, convective thunderstorms during the summer, but most of the precipitation in this area occurs as snow. The average annual temperature is 40 to 50 degrees F. The dominant soil order in this MLRA is Inceptisols. The soils in the area dominantly have a mesic soil temperature regime, an aquic or udic soil moisture regime, and mixed mineralogy. Frigid soils are found within the higher elevations. This area supports forest vegetation, particularly hardwood species. Beech-birch-maple and elm-ash-red maple are the potential forest types. The extent of oak species increases from east to west, particularly in areas of shallow and dry soils. In some areas conifers, such as white pine, are important. Aspen, hemlock, northern white-cedar, and black ash grow on the wetter soils. In some parts of the area, sugar maple has potential economic significance. Some of the major wildlife species in this area are white-tailed deer, cottontail, turkey, pheasant, and grouse.

Classification relationships

USDA NRCS: LRR: R - Northeastern Forage and Forest Region MLRA 140 - Glaciated Allegheny Plateau and Catskills Mountains EPA Ecoregions: Level III: 60 - Northern Allegheny Plateau and 58 - Northeastern Highlands Level IV: 60a - Glaciated Low Allegheny Plateau, 60b - Delaware-Neversink Highlands, 60c - Catskills Transition, and 58y - Catskill High Peaks USDA USFS: 200 Humid Temperate Domain 212 Laurentian Mixed Forest Province M212 Adirondack - New England Mixed Forest - Coniferous Forest - Alpine Meadow Province NY Natural Heritage Program Plant Community Classifications: Beech-Maple Mesic Forest Appalachian Oak-Hickory Forest International Vegetation Classification Associations: Sugar Maple - Yellow Birch - American Beech / Hobblebush Forest (CEGL006631) Sugar Maple - American Beech - White Ash / Jack-in-the-Pulpit Forest _CEGL006632) American Beech - Sugar Maple Glaciated Midwest Forest (CEGL005013) Northern Red Oak - Sugar Maple - American Beech / Mapleleaf Viburnum Forest (CEGL006633) Northern Red Oak - Sugar Maple / Mapleleaf Viburnum - Northern Spicebush Forest (CEGL006635) NatureServe Ecological Systems: Laurentian-Acadian Northern Hardwood Forest (CES201.564) North-Central Interior Beech-Maple Forest (CES202.693)

Ecological site concept

Landform/Landscape Position: The site occurs on moderately steep to very steep mountain slopes and ridges. Slopes are typically greater than 30% but range from 16-70%. Soils: The soils consists of shallow (to bedrock), somewhat excessively drained and well drained soils formed in loamy till derived from sandstone, siltstone, and shale. Soil temperature regime is frigid. Reaction is very strongly acid or strongly acid throughout the mineral soil. Soil components Halcott, Hawksnest, and Wasnot with slopes of greater than 15% are correlated to this site. Vegetation: Reference community is a Northern Hardwood Forest. Characteristic vegetation includes: Trees: American beech (*Fagus grandifolia*), sugar maple (*Acer saccharum*), red oak (*Quercus rubra*), yellow birch (*Betula alleghaniensis*), red maple (*Acer rubrum*). Conifers such as eastern white pine (*Pinus strobus*) and hemlock (*Tsuga canadensis*) are minor components. Shrubs: witch-hazel (*Hamamelis virginiana*), hobblebush (*Viburnum lantanoides*), maple-leaved viburnum (*Viburnum acerifolium*) Herbaceous: hay-scented fern (*Dennstaedtia punctilobula*), painted trillium (*Trillium undulatum*), and sessile-leaved bellwort (*Uvularia sessilifolia*)

Associated sites

F140XY003NY	Cold Shallow Till Uplands
-------------	---------------------------

Similar sites

F140XY030NY	Well Drained Dense Till
-------------	-------------------------

Table 1. Dominant plant species

Tree	(1) <i>Acer saccharum</i> (2) <i>Fagus grandifolia</i>
Shrub	(1) <i>Hamamelis virginiana</i> (2) <i>Viburnum lantanoides</i>
Herbaceous	(1) <i>Dennstaedtia punctilobula</i> (2) <i>Trillium undulatum</i>

Physiographic features

The site occurs on moderately steep to very steep mountain slopes and ridges. Slopes are typically greater than 30% but range from 16-70%.

Figure 1. The landscape pattern of upland soils developed in coarse loamy glacial till and ground water flow patterns (arrow shows general flow direction) (Soil Survey of Delaware County, New York; 2006). Halcott soils are correlated to the site.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Upland > Ridge
Runoff class	Low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	430 – 1,100 m
Slope	20 – 60 %
Water table depth	180 cm

Aspect	Aspect is not a significant factor
--------	------------------------------------

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	Not specified
Slope	20 – 70 %
Water table depth	Not specified

Climatic features

Mean annual precipitation is 46 inches and evenly distributed throughout the year. Most of the rainfall occurs as high intensity, convective thunderstorms during the summer. Snowfall is heavy from late in autumn to early spring. Average frost-free and freeze-free days are 98 and 132, respectively.

Table 4 Representative climatic features

Frost-free period (characteristic range)	90-110 days
Freeze-free period (characteristic range)	120-140 days
Precipitation total (characteristic range)	970-1,300 mm
Frost-free period (actual range)	90-110 days
Freeze-free period (actual range)	120-140 days
Precipitation total (actual range)	970-1,520 mm
Frost-free period (average)	100 days
Freeze-free period (average)	130 days
Precipitation total (average)	1,170 mm

- (1) DELHI 2 SE [USC00302036], Delancey, NY
- (2) LANSING MANOR [USC00304575], Gilboa, NY
- (3) SLIDE MTN [USC00307799], Big Indian, NY
- (4) EAST JEWETT [USC00302366], Tannersville, NY
- (5) WELLSVILLE [USC00309072], Wellsville, NY
- (6) ALFRED [USC00300085], Alfred Station, NY

Influencing water features

none

Wetland description

none

Soil features

The soils consists of shallow (to bedrock), somewhat excessively drained and well drained soils formed in loamy till derived from sandstone, siltstone, and shale. Soil temperature regime is frigid. Reaction is very strongly acid or strongly acid throughout the mineral soil. Soil components Halcott, Hawksnest, and Wasnot with slopes of greater than 15% are correlated to this site.

Table 5. Representative soil features

Parent material	(1) Till – sandstone and shale (2) Till – siltstone (3) Till – conglomerate (4) Till – quartzite
Surface texture	(1) Channery silt loam (2) Loam (3) Flaggy sandy loam
Family particle size	(1) Loamy (2) Loamy-skeletal
Drainage class	Somewhat excessively drained
Permeability class	Moderate to slow
Depth to restrictive layer	40 – 50 cm
Soil depth	40 – 50 cm
Surface fragment cover <=3"	20 – 40 %

Surface fragment cover >3"	20 – 30 %
Available water capacity (Depth not specified)	5.08 – 7.62 cm
Soil reaction (1:1 water) (Depth not specified)	3.5 – 5.5
Subsurface fragment volume <=3" (Depth not specified)	10 – 40 %
Subsurface fragment volume >3" (Depth not specified)	0 – 30 %

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	10 – 40 %
Surface fragment cover >3"	10 – 30 %
Available water capacity (Depth not specified)	Not specified
Soil reaction (1:1 water) (Depth not specified)	Not specified
Subsurface fragment volume <=3" (Depth not specified)	Not specified

Subsurface fragment volume >3" (Depth not specified)	Not specified
---	---------------

Ecological dynamics

Reference community is a hardwood forest. Characteristic trees include American beech (*Fagus grandifolia*), sugar maple (*Acer saccharum*), red oak (*Quercus rubra*), yellow birch (*Betula alleghaniensis*), red maple (*Acer rubrum*). Conifers such as eastern white pine (*Pinus strobus*) and hemlock (*Tsuga canadensis*) are minor components.

Shrubs and herbaceous plants include witch-hazel (*Hamamelis virginiana*), hobblebush (*Viburnum lantanoides*), maple-leaved viburnum (*Viburnum acerifolium*), hay-scented fern (*Dennstaedtia punctilobula*), painted trillium (*Trillium undulatum*), and sessile-leaved bellwort (*Uvularia sessilifolia*)

Reference community most likely coincides with NY Natural Heritage Program's Beech-Maple mesic forest and Appalachian oak-hickory forests depending on local site conditions.

Threats to forests in general include changes in land use (e.g., clearing for development), forest fragmentation (e.g., roads), and invasive species (e.g., insects, diseases, and plants). Other threats may include over-browsing by deer, fire suppression, and air pollution (e.g., ozone and acidic deposition) (NYNHP 2019).

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

Other references

Edinger, G. J., D. J. Evans, S. Gebauer, T. G. Howard, D. M. Hunt, and A. M. Olivero (editors). 2014. Ecological Communities of New York State. Second Edition. A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY.

New York Natural Heritage Program. 2019. Online Conservation Guide for Beech-maple mesic forest. Available from: <https://guides.nynhp.org/beech-maple-mesic-forest/>. Accessed September 11, 2019.

Contributors

Joshua Hibit

Approval

Greg Schmidt, 10/01/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/20/2020
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
