

Ecological site F144AY010NH

Sandy High Floodplain

Last updated: 5/20/2025
Accessed: 09/08/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 144A—New England and Eastern New York Upland, Southern Part

MLRA 144A: New England and Eastern New York Upland, Southern Part The eastern half of the eastern part of this MLRA is in the Seaboard Lowland Section of the New England Province of the Appalachian Highlands. The western half of the eastern part and the southeastern half of the western part are in the New England Upland Section of the same province and division. The northwestern half of the western part is in the Hudson Valley Section of the Valley and Ridge Province of the Appalachian Highlands. This MLRA is a very scenic area of rolling to hilly uplands that are broken by many gently sloping to level valleys that terminate in coastal lowlands. Elevation ranges from sea level to 1,000 feet in much of the area, but it is 2,000 feet on some hills. Relief is mostly about 6 to 65 feet in the valleys and about 80 to 330 feet in the uplands. This area has been glaciated and consists almost entirely of till hills, drumlins, and bedrock-controlled uplands with a mantle of till. It is dissected by narrow glacio-fluvial valleys. The southernmost boundary of the area marks the farthest southward extent of Wisconsinian glaciation on the eastern seaboard. The river valleys and coastal plains are filled with glacial lake sediments, marine sediments, and glacial outwash. The bedrock in the eastern half of the area consists primarily of igneous and metamorphic rocks of early Paleozoic age. Granite is the most common igneous rock, and gneiss, schist, and slate are the most common metamorphic rocks. In the parts of the MLRA in eastern and southeastern New York, Devonian- to Pennsylvanian-age sandstone, shale, and limestone are dominant. Carbonate rocks, primarily dolomite and limestone, are the dominant kinds of bedrock in the part of this MLRA in northwestern Connecticut.

Classification relationships

USDA-NRCS (USDA 2006): Land Resource Region (LRR): N—East and Central Farming and Forest Region Major Land Resource Area (MLRA): 144A— New England and Eastern New York Upland, Southern Part. USDA-FS (Cleland et al. 2007) Province: 221 - Eastern Broadleaf Province Section: 221A - Lower New England Subsection: 221Aa – Boston Basin 221Ac – Narragansett-Bristol Lowland and Islands 221Ad – Southern New England Coastal Lowland 221Ae – Hudson Highlands 221Ag - Southeast New England Coastal Hills and Plains 221Ah - Worcester-Monadnock Plateau 221Ai – Gulf of Maine Coastal Plain 221Ak - Gulf of Maine Coastal Lowland Section: 221B – Hudson Valley Subsection: 221Ba – Hudson Limestone Valley 221Bb - Miami – Taconic Foothills 221Bc – Hudson Glacial Lake Plains

Ecological site concept

The site consists of deep, coarse-loamy, well drained, alluvial soils on high floodplains of mostly large and medium sized river valleys. The site is flooded less frequently or for a shorter duration than low floodplains. Representative soils are Occum and Wappinger. The reference community is a sugar maple - white ash (*Acer saccharum* - *Fraxinus americana*) forest with associated plants including American elm (*Ulmus americana*), bitternut hickory (*Carys cordiformis*), American bladdernut (*Staphylea trifolia*), ostrich fern (*Matteuccia struthiopteris*), and sedges such as Sprengel's sedge (*Carex sprengelii*). Unlike low floodplain forests, silver maple is absent from this community. Limited examples of this forest type exist since they have mostly been converted to agricultural use. River types such as large, low gradient and small-medium low and high gradient rivers differ in hydrologic regime and fluvial geomorphology and consequently have different community composition (Marks et al. 2011).

Associated sites

F144AY006CT	High Floodplain Levee
-------------	-----------------------

F144AY015NY	Wet Silty Low Floodplain
-------------	--------------------------

Similar sites

F144AY027MA	Moist Sandy Outwash
-------------	---------------------

Table 1. Dominant plant species

Tree	(1) <i>Acer saccharum</i> (2) <i>Fraxinus americana</i>
Shrub	Not specified
Herbaceous	(1) <i>Carex sprengeii</i>

Physiographic features

The site occurs on high floodplains of mostly large to medium sized river valleys. The site is flooded less frequently or for a shorter duration than low floodplains.

Table 2. Representative physiographic features

Landforms	(1) Hills > Drumlin (2) Till plain > Ground moraine (3) Upland > Till plain (4) Hill (5) Drumlinoid ridge (6) Ridge (7) Knoll (8) Mountain (9) Hillslope
Runoff class	Negligible to very high
Flooding frequency	None to very rare
Ponding frequency	None
Elevation	0 – 620 m

Slope	0 – 60 %
Water table depth	30 – 180 cm
Aspect	Aspect is not a significant factor

Climatic features

The Koppen-Geiger climate classification of the area in which this MLRA occurs varies between Dfb (Warm-summer humid continental) in the North, and Dfa (Hot-summer humid continental) to the southern portion of the MLRA. Precipitation is usually uniformly distributed throughout the year. Near the coast, however, it is slightly lower in summer. It is slightly higher in spring and fall in inland areas. Rainfall occurs as high-intensity, convective thunderstorms during the summer. During the winter, most of the precipitation occurs as moderate-intensity storms (northeasters) that produce large amounts of rain or snow. The freeze-free period increases in length to the south.

Table 3 Representative climatic features

Frost-free period (characteristic range)	120-150 days
Freeze-free period (characteristic range)	170-180 days
Precipitation total (characteristic range)	1,140-1,320 mm
Frost-free period (actual range)	110-170 days
Freeze-free period (actual range)	140-200 days
Precipitation total (actual range)	1,020-1,350 mm
Frost-free period (average)	140 days
Freeze-free period (average)	170 days
Precipitation total (average)	1,220 mm

- (1) DANBURY [USC00061762], Bethel, CT
- (2) MASSABESIC LAKE [USC00275211], Manchester, NH
- (3) WOONSOCKET [USC00379423], Manville, RI
- (4) WORCESTER RGNL AP [USW00094746], Leicester, MA
- (5) WEST POINT [USC00309292], Cold Spring, NY
- (6) STORRS [USC00068138], Storrs Mansfield, CT
- (7) CHARLOTTEBURG RESERVE [USC00281582], Newfoundland, NJ
- (8) ALBANY AP [USW00014735], Latham, NY

Influencing water features

The site is flooded very rarely or for a shorter duration than low floodplains.

Wetland description

NONE

Soil features

The site consists of very deep, well drained coarse loamy soils formed in alluvial sediments. They are nearly level soils on flood plains, subject to common flooding. Slope ranges from 0 to 3 percent. Saturated hydraulic conductivity is moderately high or high in the loamy layers and high or very high in the sandy substratum. Soil pH ranges from very strongly acid to neutral.

Representative soils are Occum, and Wappinger.

Figure 7. Sandy High Floodplain (Occum soils)

Table 4. Representative soil features

Parent material	(1) Alluvium – granite and gneiss (2) Schist (3) Shale (4) Slate (5) Sandstone and siltstone
Surface texture	(1) Very fine sandy loam (2) Fine sandy loam (3) Loam (4) Silt loam
Family particle size	(1) Coarse-loamy over sandy or sandy-skeletal (2) Coarse-loamy
Drainage class	Well drained
Permeability class	Very slow to moderate
Depth to restrictive layer	Not specified
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified

Available water capacity (0-101.6cm)	10.16 – 15.24 cm
Soil reaction (1:1 water) (0-101.6cm)	3.6 – 8.4
Subsurface fragment volume <=3" (Depth not specified)	0 – 40 %
Subsurface fragment volume >3" (Depth not specified)	0 – 10 %

Ecological dynamics

[Caveat: The vegetation information contained in this section and is only provisional, based on concepts, not yet validated with field work.*]

The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer 2003). Terrestrial ecological SYSTEMS are specifically defined as a group of plant community-types called ASSOCIATIONS that tend to [co-]occur within landscapes with similar ecological processes, substrates, and/or environmental gradients. They are intended to provide a classification unit that is readily mappable, often from terrain and remote imagery, and readily identifiable by conservation and resource managers in the field. A given system will typically manifest itself in a landscape at intermediate geographic scales of tens-to-thousands of hectares and will persist for 50 or more years. A vegetation association is a plant community that is much more specific to a given soil, geology, landform, climate, hydrology, and disturbance history. It is the basic unit for vegetation classification and recognized by the US National Vegetation Classification (US FDGC 2008). Each association will be named by the diagnostic and often dominant species that occupy the different height strata (tree, sapling, shrub, and herb). Within the NatureServe Explorer database, ecological systems are numbered by a Community Ecological System Code (CES) and individual vegetation associations are assigned an identification number called a Community Element Global Code (CEGL).

Additional and more localized vegetation information is provided by the State Natural Heritage Programs of Connecticut (Metzler and Barrett 2001) and Massachusetts (Swain and Kearsley 2001), New Hampshire (Sperduto and Nichols, 2011), and New York (Edinger et al., 2014).

The Laurentian-Acadian Floodplain Forest system (CES201.587) is characteristic of this ecological site (NatureServe 2015). This floodplain forest develops along medium to large river systems with a medium to low gradient. Disturbances are related to flood magnitude, frequency, and seasonal timing. At higher elevations in the floodplains and floodplain terraces, much of this ecological site has been converted to agriculture. The vegetation is often a mosaic of forest, woodland, shrub land, and herbaceous communities. However, with less the understory is more developed. The characteristic plant community is a Terrace Hardwood Floodplain Forest.

[*Caveat] The information presented is representative of very complex vegetation communities. Key indicator plants and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and geography. The reference plant community is not necessarily the management goal. The drafts of species lists are merely representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 3.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 3.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

Other references

REFERENCES

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C.A. Carpenter, and W.H.McNab. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. [Map. presentation scale 1:3,500,000, colored; A.M. Sloan, cartographer] Gen. Tech. Report WO-76D. U.S. Department of Agriculture, Forest Service, Washington, DC. (<https://www.fs.fed.us/research/publications/misc/73326-wo-gtr-76d-cleland2007.pdf>)

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K., Snow, and J.Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia.

Cowardin, L.M. et. al. 1979. Classification of Wetlands and Deepwater habitats of the United States. FWS/OBS-79/31, U.S. Department of the Interior, Fish and Wildlife Service, Washington, DC.

Edinger, G.J., Evans, D.J., Gebauer, S., Howard, T.G., Hunt, D.M., and A.M. Olivero, A.M. (eds.). 2014. Ecological Communities of New York State, Second Edition: A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY.

FGDC [Federal Geographic Data Committee]. 2008. National Vegetation Classification Standard, Version 2. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC..

Marks, C.O., K.A. Lutz, A.P. Olivero-Sheldon. 2011. Ecologically important floodplain forests in the Connecticut River watershed. The Nature Conservancy, Connecticut River Program. 44pp.

Metzler, K.J. and Barrett, J.P., 2006. The Vegetation of Connecticut, a Preliminary Classification. Department of Environmental Protection, State Geological and Natural History Survey of Connecticut. Rpt of Investigations No. 12.

NatureServe. 2009. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central

Databases. Arlington, VA, U.S.A. Data current as of 06 February 2009.

NatureServe 2015. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available <http://explorer.natureserve.org>. (Accessed: December 2015).

PRISM Climate Group, Oregon State University. Available <http://prism.oregonstate.edu>, (created February 26, 2013).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. Agricultural Handbook 296

Soil Survey Staff-USDA-NRCS [United States Department of Agriculture, Natural Resources Conservation Service] 2016. National Soils Information Service (NASIS Data Model Version 7.3.4) Lincoln, NE. Available description: https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/soils/survey/tools/?cid=nrcs142p2_053552 (Accessed January 2015).

Sperduto, D.D., & Nichols, W.F. 2011. Natural Communities of New Hampshire, Second Ed. NH Natural Heritage Bureau, Concord, NH. Publ. UNH Cooperative Extension.

Swain, P.C. and Kearsley, J.B., 2001. Classification of the natural communities of Massachusetts. Natural Heritage & Endangered Species Program, Massachusetts Division of Fisheries and Wildlife.

United States Environmental Protection Agency, 2013. Level III ecoregions of the continental United States. Corvallis, Oregon, U.S. EPA-National health and Environmental Effects Research Laboratory, map scale 1:7,500,000, http://www.epa.gov/wed/pages/ecoregions/level_iii_iv.htm.

Woods, A.J., J.O. Omernik, D.D. Brown, C.W. Kiilsgaard. 1996. Level IV Ecoregions of EPA Region 3. US Environmental Protection Agency National Health and Environmental Effects Research Laboratory, Corvallis, Oregon. Map scale 1:250,000.

Contributors

Nels Barrett, Ph.D. (vegetation)

Approval

Greg Schmidt, 5/20/2025

Acknowledgments

Michael Margo and tech team provided earlier drafts. Josh Hibit made compliance updates w/ 2021 Checklist V.2

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/15/2025
Approved by	
Approval date	

Composition (Indicators 10 and 12) based on	Annual Production
---	-------------------

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

- 14. Average percent litter cover (%) and depth (in):**

- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

- 17. Perennial plant reproductive capability:**
