

Ecological site F144BY009ME

Sandy Beach (reserved)

Last updated: 9/27/2024
Accessed: 08/20/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 144B—New England and Eastern New York Upland, Northern Part

This major land resource area (MLRA) is in Maine (56 percent), New Hampshire (22 percent), Vermont (14 percent), Massachusetts (6 percent), Connecticut (1 percent), and New York (1 percent). It makes up about 22,728 square miles (58,864 square kilometers). The MLRA consists of a relatively young landscape shaped by the Laurentide Ice Sheet, which covered the region from 35,000 to 10,000 years ago. Rolling hills of dense basal till converge on ridges of shallow bedrock that were scoured by glacial ice. River valleys that were flooded by melting glacial water or seawater house large expanses of glacial outwash and stratified drift in inland areas and, to a lesser extent, glaciomarine and glaciolacustrine sediment deposits in coastal areas. Organic bogs, ablation till, and alluvial flood plains make up the remaining portions of the MLRA. The soils in this region are dominantly Entisols, Spodosols, and Inceptisols. They commonly have a fragipan. The dominant suborders are Ochrepts, Orthods, Aquepts, Fluvents, and Saprist. The soils in the region dominantly have a frigid soil temperature regime with some cryic areas at higher elevation, a udic soil moisture regime, and mixed mineralogy. Most of the land is forested, and 98 percent is privately owned. Significant amounts of forest products are produced including lumber, pulpwood, Christmas trees, and maple syrup. Principal agricultural crops include forage and grains for dairy cattle, potatoes, apples, and blueberries. Wildlife habitat and recreation are important land uses. Stoniness, steep slopes, and poor drainage limit the use of many of the soils. Maritime Sandy Beaches are restricted to the Maine coast.

Classification relationships

USDA-NRCS (USDA 2006) Land Resource Region (LRR): R—Northeastern Forage and Forest Region Major Land Resource Area (MLRA): 144B—New England and Eastern New York Upland, Northern Part USDA-FS (Cleland et al. 2007) Province: 211 - Northeastern Mixed Forest Province (in part – eastern) Section: 211D - Central Maine Coastal and Embayment Subsection: 221Da - Central Maine Embayment Subsection: 211Db - Casco Bay Coast Subsection: 221Dc - Penobscot Coast

Ecological site concept

Additional research and on-site data collection is needed to further develop interpretations for this ecological site. Maritime Sandy Beach ecological sites are beach strands occurring just at and above the limits high tide line on sandy beaches. Beaches are flooded daily by the tides and during storms. With accelerated sea-level rise, beaches are expected to migrate landward. Beach strand vegetation is typically sparsely vegetated and more often dominated by forbs than grasses.

Associated sites

F144BY010ME	Coastal Dune (reserved) Coastal Dunes often are developed just landward of the Sandy Beaches.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
------	---------------

Shrub	Not specified
Herbaceous	(1) <i>Suaeda maritima</i> (2) <i>Lathyrus japonicus</i> var. <i>maritimus</i>

Physiographic features

Maritime Sandy Beaches are depositional features of the coast where the land meets the sea.

Table 2. Representative physiographic features

Landforms	(1) Beach
Runoff class	High
Flooding duration	Extremely brief (0.1 to 4 hours)
Flooding frequency	Very frequent
Elevation	0 m
Slope	0 %
Aspect	Aspect is not a significant factor

Climatic features

The climate is humid and temperate and is characterized by warm summers and cold winters. Precipitation generally is evenly distributed throughout the year. Near the coast, it is slightly lower in summer. In inland areas, it is slightly higher in spring and fall. Rainfall occurs during high-intensity, convective thunderstorms in summer. In winter, most of the precipitation occurs as moderate-intensity storms (northeasters) that produce large amounts of rain or snow. Heavy snowfalls commonly occur late in winter. Temperatures and the length of the freeze-free period increase from north to south and closer to the coast.

Table 3 Representative climatic features

Frost-free period (characteristic range)	130-150 days
Freeze-free period (characteristic range)	170-180 days
Precipitation total (characteristic range)	1,170-1,320 mm
Frost-free period (actual range)	130-160 days
Freeze-free period (actual range)	150-190 days

Precipitation total (actual range)	1,140-1,400 mm
Frost-free period (average)	140 days
Freeze-free period (average)	170 days
Precipitation total (average)	1,250 mm

- (1) BELFAST [USC00170480], Belfast, ME
- (2) LINCOLN SAN DIST WTP [USC00174683], Blue Hill, ME
- (3) EASTPORT [USC00172426], Eastport, ME
- (4) ROBBINSTON [USC00177238], Robbinston, ME
- (5) PORT CLYDE [USC00176881], Tenants Harbor, ME
- (6) ELLSWORTH 3SSW [USC00172623], Surry, ME
- (7) ACADIA NP [USC00170100], Bar Harbor, ME
- (8) WEST ROCKPORT 1 NNW [USC00179593], Rockport, ME
- (9) WESTPORT ISLAND [USC00179547], Wiscasset, ME
- (10) WISCASSET AP [USW00094623], Wiscasset, ME

Influencing water features

Beaches are subject to the ebb and flow of oceanic tides.

Wetland description

According to the National Wetlands Inventory, Sandy Beaches are classified as: Marine or Estuarine (system) , intertidal (subsystem), unconsolidated shore (class), and sand (subclass).

Soil features

Typical soil mapunits are beaches and coastalbeaches.

Table 4. Representative soil features

Parent material	(1) Beach sand
Surface texture	(1) Sand

Ecological dynamics

Caveat: The vegetation information contained in this section and is only provisional, based on concepts, and future projects support validation through field work. *) The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer et al., 2003) and localized associations provided by the New York Natural Heritage Program (Edinger et al., 2014), Maine Natural Areas Program (Gawler and Cutko, 2010), New Hampshire Natural Heritage Program (Sperduto and Nichols, 2011), and Massachusetts Division of Fisheries and Wildlife (Swain, 2020).

The maritime sandy beach ecological site is subject overwash during high tides and storms as well as salt spray, desiccating winds, and sun exposure.

Additional research and on-site data collection is needed to further develop interpretations for this ecological site.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Future work is needed, as described in a future project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

Other references

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C.A. Carpenter, and W.H.McNab. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. [Map. presentation scale 1:3,500,000, colored; A.M. Sloan, cartographer] Gen. Tech. Report WO-76D. U.S. Department of Agriculture, Forest Service, Washington, DC. (<https://www.fs.fed.us/research/publications/misc/73326-wo-gtr-76d-cleland2007.pdf>)

Comer, P., D. Faber-Langendoen, R. Evans, S. Grawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schultz, K. Snow, and J. Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia

Edinger, G. J., D. J. Evans, S. Gebauer, T. G. Howard, D. M. Hunt, and A. M. Olivero (editors). 2014. Ecological Communities of New York State. Second Edition. A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY.

Gawler, S. and A. Cutko. 2010. Natural Landscapes of Maine: A Guide to Natural Communities and Ecosystems. Maine Natural Areas Program, Maine Department of Conservation, Augusta, Maine.

NatureServe. 2021. NatureServe Explorer: An online encyclopedia of life [web application]. NatureServe, Arlington, Virginia. <https://explorer.natureserve.org/>. (accessed 2022).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. Agricultural Handbook 296.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions. Available online. (accessed 11 Aug. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Climate Research Station Data. Available online. (accessed 23 June. 2021).

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) Database for [MLRA 141, Maine]. Available online. (accessed 14 Oct. 2021).

Sperduto, D.D. and William F. Nichols. 2011. Natural Communities of New Hampshire. 2nd Ed. NH Natural Heritage Bureau, Concord, NH. Pub. UNH Cooperative Extension, Durham, NH.

Swain, P. C. 2020. Classification of the Natural Communities of Massachusetts. Massachusetts Division of Fisheries and Wildlife, Westborough, MA

USNVC [United States National Vegetation Classification]. 2017. United States National Vegetation Classification Database V2.01. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. Available The U.S. National Vegetation Classification (usnvc.org) (accessed 2022).

Contributors

Approval

Nels Barrett, 9/27/2024

Acknowledgments

Nels Barrett and Nick Butler provided considerable review of this ecological site concept.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	08/20/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future

establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
