

Ecological site F152BY001TX

Depressional

Last updated: 9/22/2023
Accessed: 08/31/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 152B–Western Gulf Coast Flatwoods

Major Land Resource Area (MLRA) 152B, Western Gulf Coast Flatwoods, is in eastern Texas and western Louisiana. Locally termed the Flatwoods, the area is dominated by coniferous forest covering 5,681 square miles (14,714 square kilometers). The region is a hugely diverse transition zone between the northern and eastern mixed forests and southern and western coastal prairies and grasslands.

Classification relationships

Major Land Resource Area (MLRA) (USDA-Natural Resources Conservation Service, 2006)

Ecological site concept

The Depressional ecological site has very deep, very poorly or poorly drained soils that are ponded to 0.5 feet from November to May. The depressions occur as isolated system on uplands and terraces. Their isolation and inundation of water form their hydrophytic plant communities.

Associated sites

F152BY010TX	Terrace Soils have better drainage and on terrace position.
F152BY005TX	Seasonally Wet Loamy Upland Soils are on a slightly higher landform.
F152BY006TX	Well Drained Loamy Upland Soils are well drained and on higher landform.
F152BY007TX	Poorly Drained Loamy Upland Soils have better drainage and on slightly higher landform.

Similar sites

F152BY011TX	Swamp Soils are ponded almost year-round and on a lower landform.
F152BY007TX	Poorly Drained Loamy Upland Soils have better drainage and on slightly higher landform.

Table 1. Dominant plant species

Tree	(1) <i>Nyssa sylvatica</i> (2) <i>Quercus nigra</i>
Shrub	(1) <i>Smilax laurifolia</i>
Herbaceous	(1) <i>Carex cherokeensis</i>

Physiographic features

The ecological site includes areas on flat to nearly level depressions on uplands and terraces. Slope is 0 to 1 percent, but mainly 0.5 percent. Elevation ranges from 10 to 150 feet. Water table depth fluctuates throughout the year. From November to May, the depth to the top of water table will be 0 to 63 inches. The water table will deepen during the warmer months of the year. The sites are typically ponded from November to May and have aquic conditions during that time.

Table 2. Representative physiographic features

Landforms	(1) Coastal plain > Depression
Runoff class	Negligible
Flooding duration	Brief (2 to 7 days)
Flooding frequency	None to frequent
Ponding duration	Very long (more than 30 days)
Ponding frequency	None to frequent
Elevation	0 – 50 m
Slope	0 %
Ponding depth	0 – 30 cm

Water table depth	0 – 160 cm
Aspect	Aspect is not a significant factor

Climatic features

The Western Gulf Coast Flatwoods (MLRA 152B) is within the humid subtropical climate zone. The region boasts one of the highest rainfall averages in the southern United States, over 60 inches (152 centimeters) annually. This is due to the gulf currents that carry humid air to the region, where it condenses and precipitates. Rainfall averages are fairly consistent month by month, ranging from the lowest of 3.5 inches (8.9 centimeters) in March and the highest of 5.6 inches (14.3 centimeters) in June.

The area is prone to severe thunderstorms and tornadoes when the proper conditions exist, generally in the springtime. Sometimes excessive rainfall occurs, leading to flooding. Hurricanes also strike the region, generally in late summer or early fall. These extreme weather events can be quite destructive, toppling trees, and serves to naturally reset the vegetation to primary succession. The higher humidity of the region amplifies the feeling of heat during the summer. Prolonged droughts and snowfall events are rare.

Table 3 Representative climatic features

Frost-free period (average)	250 days
Freeze-free period (average)	290 days
Precipitation total (average)	1,600 mm

- (1) DE RIDDER [USC00162367], Deridder, LA
- (2) OBERLIN FIRE TWR [USC00166938], Oberlin, LA
- (3) ORANGE 9 N [USC00416680], Orange, TX
- (4) LIBERTY [USC00415196], Liberty, TX
- (5) LUMBERTON [USC00415435], Silsbee, TX
- (6) TOWN BLUFF DAM [USC00419101], Jasper, TX
- (7) ELIZABETH [USC00162800], Oakdale, LA
- (8) CLEVELAND [USC00411810], Cleveland, TX
- (9) WILDWOOD [USC00419754], Kountze, TX
- (10) DE QUINCY [USC00162361], Dequincy, LA

Influencing water features

Water collects in the upland depressions by rainfall and drainage instead of overflow flooding like bottomland systems. The Depressionals typically pond water from November to May.

Wetland description

This site has hydric soils. Onsite investigation is necessary to determine exact local conditions.

Soil features

The soils of this site are characterized by very deep, very poorly or poorly drained soils formed from loamy alluvium. The soils typically have loam or silty loam surface textures and clay loam or silty clay loam subsurface textures. Water saturation is to be expected during parts of the year and the profile will have gleyed coloring starting as high as the A horizon. Soils within this ecological site include Caneyhead, Jayhawker, and Lelavale. The series fall within the Typic Glossaqualf taxonomic class, with the exception of Jayhawker, which is a Typic Paleaquult.

Table 4. Representative soil features

Parent material	(1) Fluviomarine deposits – igneous, metamorphic and sedimentary rock
Surface texture	(1) Silt loam (2) Loam
Family particle size	(1) Fine-silty (2) Coarse-silty
Drainage class	Poorly drained to very poorly drained
Permeability class	Slow to very slow
Soil depth	200 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-152.4cm)	25.4 – 30.48 cm
Calcium carbonate equivalent (0-152.4cm)	Not specified
Electrical conductivity (0-152.4cm)	Not specified
Sodium adsorption ratio (0-152.4cm)	0 – 10
Soil reaction (1:1 water) (0-152.4cm)	3.5 – 10

Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The information in this ecological site description (ESD), including the state-and-transition model (STM), was developed using archeological and historical data, professional experience, and scientific studies. The information is representative of a complex set of plant communities. Not all scenarios or plants are included. Key indicator plants, animals, and ecological processes are described to inform land management decisions.

Introduction – In southeastern Texas and southwestern Louisiana the transition from coastal grasslands to the large expanse of coniferous forest has been deemed the “Flatwoods”. As the name suggests, the region is relatively flat and, with many transitional areas, highly diverse in flora and fauna. Historically, the area was covered by pines with mixed hardwoods, sparse shrubs, and a diverse understory of grasses and forbs. Fire and drainage patterns play a significant role in shaping the plant communities and their development. Fire suppression, drainage alterations, and land conversion have reduced the amount of historical communities in existence today.

Background – Prior to settlement by the Europeans, the reference state for the Depressional ecological site was Blackgum/Water Oak Forests. Remnants of this presumed historic plant community still exist where natural conditions are intact. Evidence of the reference state is found in accounts of early historic explorers to the area, historic forest and biological survey teams, as well as recent ecological studies in the last 30 years. The age of this community varies, and has a diverse flora.

Settlement Management – As human settlement increased throughout the area, so did the increase in logging and grazing by domestic livestock. The logging became so extensive that by the 1930's most of the region had been cut-over. Replanting trees to historic communities was not common and early foresters began planting loblolly pine (*Pinus taeda*) for its quick growth. As more people colonized they began suppressing fire, which allowed dense thickets of shrubs to replace the herbaceous understory.

Current Management and State – Today much of the historic forest is gone, replaced by pine plantations, crops, and pastures. The areas that were not converted have been fire-suppressed so long that loblolly pine and fire intolerant hardwoods populate the overstory structure. Currently, federally-managed properties are the best place to view the remnant sites (National Park Service, U.S. Fish and Wildlife Service, etc.). Some private individuals have begun restoring communities through selective tree planting and retention of communities that remain. Other restoration efforts include mimicking natural-disturbance regimes through gap-phase regeneration on plantation sites.

Fire Regimes – Fire was a natural and important disturbance throughout the region. Fire occurred naturally from lightning strikes, by Native Americans for game movement, and eventually early European settlers. Fires throughout the Flatwoods occurred at two different times. Early in the year, they would occur during late winter and early spring, removing senescent vegetation, recycling nutrients and minerals, and spurring new plant growth. Late summer and early fall fires occurred as well, but with a different community effect. Summer fires burned hotter and with more intensity, greatly suppressing the shrub canopy layer. The summer fires also shifted the ecological site transitional state by decreasing grass densities and increasing forb densities. The topography, fuel loads, and other conditions caused patchy burns throughout the region resulting in mosaic patterns of plant communities and a heterogeneous landscape.

Disturbance Regimes – Extreme weather events occur occasionally throughout the region. Tornados uproot trees and open canopies in the spring months. In the late summer and early fall, hurricanes or tropical depressions can make landfall, dumping excessive amounts of rain and toppling trees with high winds. Another cause of large canopy openings is the effects of the southern pine beetle (*Dendroctonus frontalis*). Starting in the late 1950's, beetle outbreaks have occurred every 6 to 9 years (although a major attack has not occurred in some time); usually when the trees are stressed due to multiple environmental factors.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height M	Canopy Cover (%)	Diameter Cm	Basal Area (square M/hectare)
Tree							
willow oak	QUPH	<i>Quercus phellos</i>	Native	–	–	–	–
red maple	ACRU	<i>Acer rubrum</i>	Native	–	–	–	–
American holly	ILOP	<i>Ilex opaca</i>	Native	–	–	–	–
loblolly pine	PITA	<i>Pinus taeda</i>	Native	–	–	–	–
sweetgum	LIST2	<i>Liquidambar styraciflua</i>	Native	–	–	–	–
water oak	QUNI	<i>Quercus nigra</i>	Native	–	–	–	–
blackgum	NYSY	<i>Nyssa sylvatica</i>	Native	–	–	–	–

Table 7. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (m)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
Frank's sedge	CAFR3	<i>Carex frankii</i>	Native	–	–
bushy bluestem	ANGL2	<i>Andropogon glomeratus</i>	Native	–	–
sedge	CAREX	<i>Carex</i>	Native	–	–
grassleaf rush	JUMA4	<i>Juncus marginatus</i>	Native	–	–
Cherokee sedge	CACH3	<i>Carex cherokeensis</i>	Native	–	–
Forb/Herb					
knotweed	POLYG4	<i>Polygonum</i>	Native	–	–
Fern/fern ally					
western brackenfern	PTAQ	<i>Pteridium aquilinum</i>	Native	–	–
Shrub/Subshrub					
swamp titi	CYRA	<i>Cyrilla racemiflora</i>	Native	–	–
wax myrtle	MOCE2	<i>Morella cerifera</i>	Native	–	–
dwarf palmetto	SAMI8	<i>Sabal minor</i>	Native	–	–
Vine/Liana					
laurel greenbrier	SMLA	<i>Smilax laurifolia</i>	Native	–	–
lanceleaf greenbrier	SMSM	<i>Smilax smallii</i>	Native	–	–

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Wood products

These soils belong to the 3w9 woodland suitability group. Ponding of water is common during wet months. They have a moderate potential for woodland management, both pine and hardwood. The 50-year site index for loblolly pine averages between 80 and 85 feet (approximately 55 to 57 feet on a 25-year curve), but can range from 75 to 90 feet depending on drainage. The 50-year site index for bottomland oaks can range from 70 to 80 feet. The yield from an unmanaged, natural stand of loblolly pine, over a 50-year period, is approximately 250 board feet (Doyle Rule), 2 tons, or 80 cubic feet per acre per year. Management can substantially increase this yield. Access and equipment operability on these soils is poor during wet periods, when rutting will occur due to saturation of the soil. Harvesting and other operations may need to be suspended during such periods. Wetness also makes these soils poorly suited for log landings and roads. Low strength makes them moderately suited for road construction. Raising and crowning the road bed will be necessary and care must be taken to avoid interrupting the natural drainage. Site preparation operations should be limited to the dry

months and planting should be planned for the drier part of the planting season. Use of herbicides for site preparation must also take into consideration the poor drainage and high seasonal water tables on these soils. Wetness may cause a moderate loss in seedling survival. Bedding may be needed.

Inventory data references

This site description was developed as part of the provisional ecological site initiative using historic soil survey manuscripts, available range site descriptions, and low intensity field sampling.

Other references

Ajilvsgi, G. 2003. Wildflowers of Texas. Revised edition. Shearer Publishing, Fredericksburg, TX.

Ajilvsgi, G. 1979. Wildflowers of the Big Thicket. Texas A&M University Press, College Station, TX.

Allen, J. A., B. D. Keeland, J. A. Stanturf, and A. F. Kennedy Jr. 2001. A guide to bottomland hardwood restoration. Technical report, USGS/BRD/ITR-2000-0011.

Bray, W. L. 1904. Forest resources of Texas. Bureau of Forestry Bulletin 47, Government Printing Office, Washington D.C.

Diggs, G. M., B. L. Lipscomb, M. D. Reed, and R. J. O'Kennon. 2006. Illustrated flora of East Texas. Second edition. Botanical Research Institute of Texas & Austin College, Fort Worth, TX.

Jones, S. D., J. K. Wipff, and P. M. Montgomery. 1997. Vascular plants of Texas: a comprehensive checklist including synonymy, bibliography, and index. University of Texas Press, Austin.

Liu, C., P. A. Harcombe, and I. S. Elsie. 1990. Fire study report, including Roy E. Larsen Preserve species list. Summer 1990. Department of Ecology and Evolutionary Biology, Rice University, Houston, TX.

Marks, P. L., and P. A. Harcombe. 1981. Forest Vegetation of the Big Thicket, southeast Texas. Ecological Monographs 51:287-305.

Matos, J. A. 1985. Roy E. Larsen Sandylands Sanctuary vascular plant species list. Master thesis, Stephen F. Austin University, Nacogdoches, TX.

NatureServe. 2002. International classification of ecological communities: Terrestrial vegetation of the United States. National forests in Texas final report. NatureServe, Arlington, VA.

Nixon, E. S. 2000. Trees, shrubs & woody vines of East Texas. Second edition. Bruce Lyndon Cunningham Productions, Nacogdoches, TX.

Randall, J. M., and J. Marinelli. 1996. Invasive plants: weeds of the global garden. Volume 149. Brooklyn Botanic Garden, Brooklyn, NY.

Stanturf, J. A., S. H. Schoenholtz, C. J. Schweitzer, and J. P. Shepard. 2001. Achieving restoration success: Myths in bottomland hardwood forests. Restoration Ecology, 9:189-200.

Stringham, T. K., W. C. Krueger, and P. L. Shaver. 2003. State and transition modeling: An ecological process approach. Journal of Range Management 56:106-113.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) Database.

Truett, J. C. 1984. Land of bears and honey: A natural history of East Texas. The University of Texas Press, Austin, TX.

Van Kley, J. E., R. L. Turner, L. S. Smith, and R. E. Evans. 2007. Ecological classification system for the national forests and adjacent areas of the West Gulf Coastal Plain. Second approximation. Stephen F. Austin University and The Nature Conservancy, Nacogdoches, TX.

USDA-NRCS Ag Handbook 296 (2006).

Vines, R. A. 1960. Trees, shrubs, and woody vines of the Southwest. University of Texas Press, Austin, TX.

Watson, G. E. 2006. Big Thicket Plant Ecology. Third Edition. University of North Texas Press, Denton, TX.

Contributors

Tyson Hart

Approval

Bryan Christensen, 9/22/2023

Acknowledgments

Thanks to all involved during the preparation, sampling, and reviewing of the Flatwoods project. Thanks to Josh Berry, Dennis Brezina, Kenny Hall, Jason Hohlt, Stacey Kloesel, Ricky Lambert, Cody Langston, Mark Moseley, Ramiro Molina, Mike Oliver, Alan Peer, Sara Russell, Don Sabo, Mary Webb-Marek, and Jon Wiedenfeld for all their help.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be

verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/20/2021
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
