

Ecological site F152BY002TX

Sodic Flats

Last updated: 9/22/2023
Accessed: 08/28/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 152B–Western Gulf Coast Flatwoods

Major Land Resource Area (MLRA) 152B, Western Gulf Coast Flatwoods, is in eastern Texas and western Louisiana. Locally termed the Flatwoods, the area is dominated by coniferous forest covering 5,681 square miles (14,714 square kilometers). The region is a hugely diverse transition zone between the northern and eastern mixed forests and southern and western coastal prairies and grasslands.

Classification relationships

Major Land Resource Area (MLRA) (USDA-Natural Resources Conservation Service, 2006)

Ecological site concept

The Sodic Flat ecological site has very deep, poorly drained soils influenced by salt and bioturbation. The churning of the soil is typically caused by crawfish; coupled with the presence of salts, only plants adapted to these unique conditions survive and grow.

Associated sites

F152BY007TX	Poorly Drained Loamy Upland Soils are loamy and have no salt concentrations or bioturbation.
F152BY005TX	Seasonally Wet Loamy Upland Soils are loamy and have no salt concentrations or bioturbation.
F152BY006TX	Well Drained Loamy Upland Soils are loamy, well drained, and have no salt concentrations or bioturbation.

Similar sites

F152BY007TX	Poorly Drained Loamy Upland Soils are loamy and have no salt concentrations or bioturbation.
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>Pinus palustris</i> (2) <i>Pinus taeda</i>
Shrub	Not specified
Herbaceous	(1) <i>Drosera brevifolia</i> (2) <i>Xyris</i>

Physiographic features

The ecological site includes areas on flats on uplands. Slope is 0 to 1 percent. Elevation ranges from 10 to 151 feet. From November to May, the depth to the top of water table will be 0 to 9 inches. The water table will deepen during the warmer months of the year.

Table 2. Representative physiographic features

Landforms	(1) Coastal plain > Flat
Runoff class	High
Flooding frequency	None
Ponding frequency	None
Elevation	0 – 50 m
Slope	0 %
Water table depth	0 – 20 cm
Aspect	Aspect is not a significant factor

Climatic features

The Western Gulf Coast Flatwoods (MLRA 152B) is within the humid subtropical climate zone. The region boasts one of the highest rainfall averages in the southern United States, over 60 inches (152 centimeters) annually. This is due to the gulf currents that carry humid air to the region, where it condenses and precipitates. Rainfall averages are fairly consistent month by month, ranging from the lowest of 3.5 inches (8.9 centimeters) in March and the highest of 5.6 inches (14.3 centimeters) in June.

The area is prone to severe thunderstorms and tornadoes when the proper conditions exist, generally in the springtime. Sometimes excessive rainfall occurs, leading to flooding. Hurricanes also strike the region, generally in late summer or early fall. These extreme weather events can be quite destructive, toppling trees, and serves to naturally reset the vegetation to primary succession. The higher humidity of the region amplifies the feeling of heat during the summer. Prolonged droughts and snowfall events are rare.

Table 3 Representative climatic features

Frost-free period (average)	250 days
-----------------------------	----------

Freeze-free period (average)	290 days
Precipitation total (average)	1,600 mm

- (1) WILDWOOD [USC00419754], Kountze, TX
- (2) LIBERTY [USC00415196], Liberty, TX
- (3) LUMBERTON [USC00415435], Silsbee, TX
- (4) TOWN BLUFF DAM [USC00419101], Jasper, TX
- (5) DE QUINCY [USC00162361], Dequincy, LA
- (6) DE RIDDER [USC00162367], Deridder, LA
- (7) ELIZABETH [USC00162800], Oakdale, LA
- (8) OBERLIN FIRE TWR [USC00166938], Oberlin, LA
- (9) CLEVELAND [USC00411810], Cleveland, TX
- (10) ORANGE 9 N [USC00416680], Orange, TX

Influencing water features

While no ponding or flooding occurs, the soils within the ecological site have a seasonally high water table from November to May and all are hydric. The sites have potential for being wetlands, but specific determinations will need to take place in the field by observing water, plants, and soils.

Wetland description

This site has hydric soils. Onsite investigation is necessary to determine exact local conditions.

Soil features

The soils of this site consist of very deep, poorly drained soils formed from fluviomarine deposits. The soils typically have loam or silty loam surface textures and clay loam or silty clay loam subsurface textures. Soils within this ecological site include Nona, Plank, and Vidor. Nona and Plank within the taxonomic class of Natric Vermaqualf and Vidor a Vermic Natraqualf. The term natric refers to salt concentrations within the soil and vermic refers to bioturbation caused by animals, typically crawfish.

Table 4. Representative soil features

Parent material	(1) Fluviomarine deposits – igneous and metamorphic rock
Surface texture	(1) Loam (2) Silt loam
Family particle size	(1) Fine-silty (2) Coarse-silty
Drainage class	Poorly drained
Permeability class	Very slow
Soil depth	200 cm

Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-152.4cm)	20.32 – 27.94 cm
Calcium carbonate equivalent (0-152.4cm)	Not specified
Electrical conductivity (0-152.4cm)	0 – 10 mmhos/cm
Sodium adsorption ratio (0-152.4cm)	0 – 20
Soil reaction (1:1 water) (0-152.4cm)	3.5 – 10
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Ecological dynamics

The information in this ecological site description (ESD), including the state-and-transition model (STM), was developed using archeological and historical data, professional experience, and scientific studies. The information is representative of a complex set of plant communities. Not all scenarios or plants are included. Key indicator plants, animals, and ecological processes are described to inform land management decisions.

Introduction – In southeastern Texas and southwestern Louisiana the transition from coastal grasslands to the large expanse of coniferous forest has been deemed the “Flatwoods”. As the name suggests, the region is relatively flat and, with many transitional areas, highly diverse in flora and fauna. Historically, the area was covered by pines with mixed hardwoods, sparse shrubs, and a diverse understory of grasses and forbs. Fire and drainage patterns play a significant role in shaping the plant communities and their development. Fire suppression, drainage alterations, and land conversion have reduced the amount of historical communities in existence today.

Background – Prior to settlement by the Europeans, the reference state for the Sodic Flats were Longleaf Pine/Loblolly Pine Woodlands. Remnants of this presumed historic plant community still exist where natural conditions are intact. Evidence of the reference state is found in accounts of early historic explorers to the area, historic forest and biological survey teams, as well as recent ecological studies in the last 30 years. The age of this woodland community varies, and has a diverse flora.

Settlement Management – As human settlement increased throughout the area, so did the increase in logging and grazing by domestic livestock. The logging became so extensive that by the 1930's most of the region had been cut-over. Replanting trees to historic communities was not common and early foresters began planting loblolly pine (*Pinus taeda*) for its quick growth. As more people colonized they began suppressing fire, which allowed dense thickets of shrubs to replace the herbaceous understory.

Current Management and State – Today much of the historic forest is gone, replaced by pine plantations, crops, and pastures. The areas that were not converted have been fire-suppressed so long that loblolly pine and fire intolerant hardwoods populate the overstory structure. Currently, federally-managed properties are the best place to view the remnant sites (National Park Service, U.S. Fish and Wildlife Service, etc.). Some private individuals have begun restoring communities through selective tree planting and retention of communities that remain. Other restoration efforts include mimicking natural-disturbance regimes through gap-phase regeneration on plantation sites.

Fire Regimes – Fire was a natural and important disturbance throughout the region. Fire occurred naturally from lightning strikes, by Native Americans for game movement, and eventually early European settlers. Fires throughout the Flatwoods occurred at two different times. Early in the year, they would occur during late winter and early spring, removing senescent vegetation, recycling nutrients and minerals, and spurring new plant growth. Late summer and early fall fires occurred as well, but with a different community effect. Summer fires burned hotter and with more intensity, greatly suppressing the shrub canopy layer. The summer fires also shifted the ecological site transitional state by decreasing grass densities and increasing forb densities. The topography, fuel loads, and other conditions caused patchy burns throughout the region resulting in mosaic patterns of plant communities and a heterogeneous landscape.

Disturbance Regimes – Extreme weather events occur occasionally throughout the region. Tornados uproot trees and open canopies in the spring months. In the late summer and early fall, hurricanes or tropical depressions can make landfall, dumping excessive amounts of rain and toppling trees with high winds. Another cause of large canopy openings is the effects of the southern pine beetle (*Dendroctonus frontalis*). Starting in the late 1950's, beetle outbreaks have occurred every 6 to 9 years (although a major attack has not occurred in some time); usually when the trees are stressed due to multiple environmental factors.

State and transition model

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height M	Canopy Cover (%)	Diameter Cm	Basal Area (square M/hectare)
Tree							
longleaf pine	PIPA2	<i>Pinus palustris</i>	Native	—	—	—	—
loblolly pine	PITA	<i>Pinus taeda</i>	Native	—	—	—	—

Table 7. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (m)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
grassleaf rush	JUMA4	<i>Juncus marginatus</i>	Native	—	—
crabgrass	DIGIT2	<i>Digitaria</i>	Native	—	—
needleleaf rosette grass	DIAC	<i>Dichanthelium aciculare</i>	Native	—	—
sedge	CAREX	<i>Carex</i>	Native	—	—
little bluestem	SCSC	<i>Schizachyrium scoparium</i>	Native	—	—
yelloweyed grass	XYRIS	<i>Xyris</i>	Native	—	—
crowngrass	PASPA2	<i>Paspalum</i>	Native	—	—
panicgrass	PANIC	<i>Panicum</i>	Native	—	—
Forb/Herb					
blazing star	LIATR	<i>Liatris</i>	Native	—	—
pipewort	ERIOC	<i>Eriocaulon</i>	Native	—	—
eryngo	ERYNG	<i>Eryngium</i>	Native	—	—
anisescented goldenrod	SOOD	<i>Solidago odora</i>	Native	—	—
Shrub/Subshrub					

yaupon	ILVO	<i>Ilex vomitoria</i>	Native	–	–
wax myrtle	MOCE2	<i>Morella cerifera</i>	Native	–	–
Tree					
longleaf pine	PIPA2	<i>Pinus palustris</i>	Native	–	–
sweetbay	MAVI2	<i>Magnolia virginiana</i>	Native	–	–
loblolly pine	PITA	<i>Pinus taeda</i>	Native	–	–
Vine/Liana					
greenbrier	SMILA2	<i>Smilax</i>	Native	–	–

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Wood products

These soils occur in the Woodland Suitability Group 3w9. They have moderate potential for woodland management, both pine and hardwood. The 50-year site index for loblolly pine averages 80 feet (approximately 55 feet on a 25-year curve). The site index for bottomland oaks ranges from 65 to 75 feet. The yield from an unmanaged, natural stand of loblolly pine, over a 50-year period, is approximately 230 board feet (Doyle Rule), 1.84 tons, or 75 cubic feet per acre per year. Management can substantially increase this yield. Access and equipment operability on these soils is poor during wet periods due to saturation of the soil. Harvesting and other operations may need to be suspended during such periods, when rutting will occur. Wetness also makes these soils poorly suited for log landings and roads. Low strength makes them moderately suited for road construction material. Road layout should avoid these soils whenever possible. Site preparation operations should be limited to the dry months and planting should be planned for the drier part of the planting season. Use of herbicides for site preparation must also take into consideration the poor drainage on these soils. Applications should not be made during wet periods. Wetness may cause a moderate loss in pine seedling survival. Slash pine may be suited to these soils.

Type locality

Location 1: Hardin County, TX	
UTM zone	N
UTM northing	30.5458611
UTM easting	-94.411444
General legal description	Sundew Trail – Big Thicket National Forest

Other references

- Ajilvsgi, G. 2003. Wildflowers of Texas. Revised edition. Shearer Publishing, Fredericksburg, TX.
- Ajilvsgi, G. 1979. Wildflowers of the Big Thicket. Texas A&M University Press, College Station, TX.
- Allen, J. A., B. D. Keeland, J. A. Stanturf, and A. F. Kennedy Jr. 2001. A guide to bottomland hardwood restoration. Technical report, USGS/BRD/ITR-2000-0011.
- Bray, W. L. 1904. Forest resources of Texas. Bureau of Forestry Bulletin 47, Government Printing Office, Washington D.C.

Diggs, G. M., B. L. Lipscomb, M. D. Reed, and R. J. O'Kennon. 2006. Illustrated flora of East Texas. Second edition. Botanical Research Institute of Texas & Austin College, Fort Worth, TX.

Jones, S. D., J. K. Wipff, and P. M. Montgomery. 1997. Vascular plants of Texas: a comprehensive checklist including synonymy, bibliography, and index. University of Texas Press, Austin.

Liu, C., P. A. Harcombe, and I. S. Elisk. 1990. Fire study report, including Roy E. Larsen Preserve species list. Summer 1990. Department of Ecology and Evolutionary Biology, Rice University, Houston, TX.

Marks, P. L., and P. A. Harcombe. 1981. Forest Vegetation of the Big Thicket, southeast Texas. Ecological Monographs 51:287-305.

Matos, J. A. 1985. Roy E. Larsen Sandylands Sanctuary vascular plant species list. Master thesis, Stephen F. Austin University, Nacogdoches, TX.

NatureServe. 2002. International classification of ecological communities: Terrestrial vegetation of the United States. National forests in Texas final report. NatureServe, Arlington, VA.

Nixon, E. S. 2000. Trees, shrubs & woody vines of East Texas. Second edition. Bruce Lyndon Cunningham Productions, Nacogdoches, TX.

Randall, J. M., and J. Marinelli. 1996. Invasive plants: weeds of the global garden. Volume 149. Brooklyn Botanic Garden, Brooklyn, NY.

Stanturf, J. A., S. H. Schoenholtz, C. J. Schweitzer, and J. P. Shepard. 2001. Achieving restoration success: Myths in bottomland hardwood forests. Restoration Ecology, 9:189-200.

Stringham, T. K., W. C. Krueger, and P. L. Shaver. 2003. State and transition modeling: An ecological process approach. Journal of Range Management 56:106-113.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) Database.

Truett, J. C. 1984. Land of bears and honey: A natural history of East Texas. The University of Texas Press, Austin, TX.

Van Kley, J. E., R. L. Turner, L. S. Smith, and R. E. Evans. 2007. Ecological classification system for the national forests and adjacent areas of the West Gulf Coastal Plain. Second approximation. Stephen F. Austin University and The Nature Conservancy, Nacogdoches, TX.

USDA-NRCS Ag Handbook 296 (2006).

Vines, R. A. 1960. Trees, shrubs, and woody vines of the Southwest. University of Texas Press, Austin, TX.

Watson, G. E. 2006. Big Thicket Plant Ecology. Third Edition. University of North Texas Press, Denton, TX.

Contributors

Tyson Hart

Approval

Bryan Christensen, 9/22/2023

Acknowledgments

Thanks to all involved during the preparation, sampling, and reviewing of the Flatwoods project. Thanks to Josh Berry, Dennis Brezina, Kenny Hall, Jason Hohlt, Stacey Kloesel, Ricky Lambert, Cody Langston, Mark Moseley, Ramiro Molina, Mike Oliver, Alan Peer, Sara Russell, Don Sabo, Mary Webb-Marek, and Jon Wiedenfeld for all their help.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/20/2021
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
