

# Ecological site F153BY045NC

## Moist Clay Rises and Flats

Last updated: 4/02/2025  
Accessed: 08/20/2026

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 153B–Tidewater Area

The MLRA notes section provides a brief description of the entire MLRA. This brief description of the entire MLRA is intended to provide some context about the MLRA that this ecological site is within. A more complete description of the MLRA can be found in Ag Handbook 296 (USDA-NRCS, 2022). This MLRA stretches along the Atlantic coastline from northern Florida to southern Virginia. It features young marine terrace flats, broad flood plains and deltas, tidal marshes and estuaries, barrier sea islands, and a beach ridge system that spans the length of the MLRA. Its broad, shallow valleys with large rivers, tidal marshes, swamps, estuaries, drowned valleys, sea islands, and beaches, are all features of the Late Quaternary (USDA-NRCS, 2017). The Suffolk Scarp is the upper (western) limit of this MLRA and marks the extent of the ocean shoreline before it retreated during the Wisconsin period of glaciation. Fluctuating ocean levels, along with wave and wind activity, continue to rework sand deposits that comprise the ever-changing barrier sea islands and coastline in this MLRA. The marine terraces are younger to the east and are progressively older and higher inland to the west. The youngest marine terraces adjacent to the coast are very low lying and at high risk of inundation by extreme high tides, wind tides, storm surge, and extreme precipitation events. In addition to the risks of inundation, these low-lying terrestrial and freshwater systems are at high risk of salt water intrusion. The MLRA is characterized by a persistent high water table. The hydraulic gradient across this MLRA is very low. Overall, elevation ranges from sea level to less than 25 feet (0 to 8 meters). Local relief is mainly about 3 feet (1 meter) or less. Most of the surface water in this MLRA is either coming into the MLRA from the piedmont and upper coastal plain, is managed by ditching, or is ponded on the surface. Surface flow channels originating within the MLRA are extremely subtle, typically blackwater, and flow generally channelizes mostly near the shoreline where tidal processes also impact flooding processes. The dominant soil orders in this MLRA are Alfisols and Entisols. Ultisols and Histosols are important but are of lesser extent. The soils in the area are characterized by restricted drainage, a thermic temperature regime, and an aquic moisture regime. The study of subaqueous soils is of increasing importance along nearshore coastal waters. The major soil suborders of the MLRA include: 1) Endoaqualfs, which are very deep and loamy to clayey, 2) Endoaqualts, which are very deep and loamy to clayey, 3) Haplosaprists, which are extensive in North Carolina and Virginia, in the Great Dismal Swamp, and in broad upland wetlands known as pocosins, 4) Hapludults, which are in the higher areas of somewhat better drainage, 5) Psammments, 6) Sulfaquents, which are extensive throughout the brackish tidal marshes protected by the barrier and sea islands, 7) Sulfiwassents (subaqueous soils), which formed in low- to moderate-energy estuarine deposits, and 8) Umbraqualts, which are very deep and loamy to clayey. MLRA 153B has a lengthy north-south extent, and it runs parallel to the Atlantic coast. The MLRA extends from the northeastern corner of Florida to southern Virginia. Five states are intersected by the MLRA, including North Carolina (42 percent), Virginia (21 percent), South Carolina (20 percent), Georgia (14 percent), and Florida (3 percent). The MLRA extent makes up about 11,152 square miles (28,884 square kilometers). Because of climatic differences between the northern and southern reaches of the MLRA, vegetative communities vary with latitude. Loblolly pine, red oak, and white oak are dominant in the uplands, and blackgum, sweetgum, pond pine, laurel oak, water tupelo, and bald cypress are dominant on the bottomland. Longleaf pine and slash pine were dominant historically in the southern part of the area. Understory species common to the MLRA include switchcane, inkberry, large gallberry, greenbrier, wax myrtle, and cabbage palm. Herbaceous understory species include little bluestem, and various panicgrasses. Major wildlife species include alligator, black bear, white-tailed deer, fox, raccoon, opossum, otter, muskrat, rabbit, mink, squirrel, quail, and mourning dove. The red wolf, an endangered species, is being reintroduced in several parts of the MLRA. The nearshore estuaries of the Chesapeake Bay, the Albemarle-Pamlico estuary systems, and Atlantic Ocean provide habitat for diverse populations of terrestrial and aquatic animal species. The subaquatic vegetation in these coastal lagoon areas provides critical habitat and cover for many shellfish and juvenile finfish. The estuaries host numerous migratory waterfowl and wading birds throughout the year and are an integral part of the Atlantic Flyway. (USDA-NRCS, 2022)

### LRU notes

Currently, Ecological Site Descriptions (ESDs) for MLRA 153B cover the full north-south range of the MLRA. However, climate variation across the north-south extent warrants the development of Land Resource Unit (LRU) classifications to support more precise Ecological Site Descriptions.

Classification relationships

MLRA 153B has overlap with two level III EPA ecoregion concepts: 63) the Middle Atlantic Coastal Plain and 75) the Southern Coastal Plain. Under ecoregions 63 and 75 are a number of lower level (IV) concepts, of which several apply to MLRA 153B. These include: 63b) Chesapeake-Pamlico Lowlands and Tidal Marshes, 63c) Swamps and Peatlands, 63d) Virginia Barrier Islands and Coastal Islands, 63f) Delmarva Uplands), 63g) Carolinian Barrier Islands and Coastal Marshes, and 75j) Sea Islands/Coastal Marsh. (U.S. EPA, 2013) MLRA 153B overlaps a portion of the US Forest Service Outer Coastal Plain Mixed Forest province (232). The MLRA roughly corresponds to the easternmost portions of the Atlantic Coastal Flatwoods (232C) and the southeastern portion of the Northern Atlantic Coastal Flatwoods (232I) sections. In combination with MLRA 153A, these two MLRAs correspond very closely to the full extent of Sections 232C and 232I. (Cleland et al., 2007) Based on the USGS physiographic classification system, most of MLRA 153B is in the Sea Island section of the Coastal Plain province, in the Atlantic Plain division. The northern quarter is in the Embayed section of the same province and division. The embayed barrier islands extend from the eastern shore of the Chesapeake Bay in Virginia to north of Charleston, South Carolina (Fenneman et al., 1946). The portion in North Carolina is referred to as the Outer Banks. Large bodies of brackish water, such as Pamlico and Albemarle Sounds, are on the inland side of the barrier islands. The sea islands extend from north of Charleston, South Carolina, to Jacksonville, Florida. The reference community for this particular site is approximately aligned with Mesic Pine Flatwoods (Schafale and Weakely, 1990) and Mesic Flatwoods (FNAI, 2010).

Ecological site concept

This site is characterized by clayey and fine silty soils (dominantly Ultisols) on coastal plain rises and flats of somewhat poorly drained and moderately well drained soils. This site represents those locations where soils do not meet hydric criteria, but this site is often associated with wet sites that do meet hydric criteria, and, in this broad flat MLRA, the transition can be exceptionally subtle. This site has the potential to support a variety of vegetation communities including flatwoods, and mixed hardwood forests, but most of this site has been converted to alternative states. Historically, the vegetation communities on this site have been maintained by frequent low-intensity surface fires. Although this site is on moderately high ground within the MLRA, this site is still at risk of inundation associated with extreme tides, storm surge, and extreme precipitation.

Associated sites

|             |                                                                                                                                                                                                                                                                                                                                                                                 |
|-------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| F153BY060NC | <p><b>Wet Loamy Flats and Depressions</b></p> <p>A moist clayey flat is often associated with and difficult to distinguish from a wet loamy flat. Much of the difference between moist and wet sites is driven by variations in subsurface drainage patterns, which are difficult to see at the surface, but these sites can be distinguished by soil moisture and texture.</p> |
| F153BY065NC | <p><b>Wet Clay Flats and Depressions</b></p> <p>A moist clayey flat is often associated with and difficult to distinguish from a wet clayey flat. Much of the difference between moist and wet sites is driven by variations in subsurface drainage patterns, which are difficult to see at the surface, but these sites can be distinguished by soil moisture.</p>             |

Similar sites

|             |                                                                                                                                                                                                             |
|-------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| F153BY040NC | <p><b>Moist Loamy Rises and Flats</b></p> <p>This site is on very similar landforms but is comprised of loamy soils. The loamy soils are likely more productive than the clay soils.</p>                    |
| F153AY045NC | <p><b>Moist Clay Rises and Flats</b></p> <p>This site is on very similar landforms but in an adjacent MLRA where the marine terrace surfaces are older, more dissected, and removed from tidal impacts.</p> |

Table 1. Dominant plant species

|      |                                                                       |
|------|-----------------------------------------------------------------------|
| Tree | <p>(1) <i>Pinus palustris</i></p> <p>(2) <i>Fagus grandifolia</i></p> |
|------|-----------------------------------------------------------------------|

|            |                                                                |
|------------|----------------------------------------------------------------|
| Shrub      | (1) <i>Kalmia latifolia</i><br>(2) <i>Vaccinium arboreum</i>   |
| Herbaceous | (1) <i>Aristida stricta</i><br>(2) <i>Aristida beyrichiana</i> |

### Physiographic features

This ecological site represents rises and flats of clayey and fine-silty mineral soils (mostly Ultisols) that are currently isolated from most flood plain processes. In general, these landforms developed by marine deposition, or ancient fluvial reworking and redeposition. In comparison to it's surroundings, topographic relief is relatively high on this site. Most slopes are less than 2 percent with slopes ranging up to 6 percent.

**Table 2. Representative physiographic features**

|                    |                                                                                               |
|--------------------|-----------------------------------------------------------------------------------------------|
| Hillslope profile  | (1) Summit<br>(2) Shoulder                                                                    |
| Landforms          | (1) Coastal plain > Marine terrace<br>(2) Marine terrace > Flat<br>(3) Marine terrace > Ridge |
| Runoff class       | Very low to low                                                                               |
| Flooding frequency | None                                                                                          |
| Ponding frequency  | None                                                                                          |
| Elevation          | 0 – 10 m                                                                                      |
| Slope              | 0 %                                                                                           |
| Water table depth  | 30 – 90 cm                                                                                    |
| Aspect             | Aspect is not a significant factor                                                            |

**Table 3. Representative physiographic features (actual ranges)**

|                    |                   |
|--------------------|-------------------|
| Runoff class       | Very low to low   |
| Flooding frequency | None to very rare |

|                   |             |
|-------------------|-------------|
| Ponding frequency | None        |
| Elevation         | 0 – 10 m    |
| Slope             | 0 – 10 %    |
| Water table depth | 20 – 110 cm |

### Climatic features

The climate of MLRA 153B is generally warm, temperate, and humid with maritime influences along the coast. The maximum precipitation occurs in summer, and the minimum occurs in autumn. Rainfall is usually of moderate intensity. Occasionally, extreme weather events (e.g., northeasters, tropical storms, and hurricanes) produce large amounts of precipitation and destructive winds. Snowfall may occur in the northern end of the area. The average annual temperature is 57 to 70 degrees F (14 to 21 degrees C), increasing to the south. (USDA-NRCS, 2022)

**Table 4 Representative climatic features**

|                                            |                |
|--------------------------------------------|----------------|
| Frost-free period (characteristic range)   | 210-260 days   |
| Freeze-free period (characteristic range)  | 260-340 days   |
| Precipitation total (characteristic range) | 1,140-1,220 mm |
| Frost-free period (actual range)           | 200-270 days   |
| Freeze-free period (actual range)          | 240-360 days   |
| Precipitation total (actual range)         | 1,140-1,250 mm |
| Frost-free period (average)                | 240 days       |
| Freeze-free period (average)               | 300 days       |
| Precipitation total (average)              | 1,190 mm       |

- (1) ELIZABETH CITY CGAS [USW00013786], Elizabeth City, NC
- (2) CHARLESTON CITY [USW00013782], Charleston, SC
- (3) BRUNSWICK 23 S [USW00063856], Saint Marys, GA

### Influencing water features

This MLRA is dominated by a persistent high water table, but this site is characterized by better internal drainage. Fluctuations in water table depth are mostly driven by variability in subsurface drainage patterns, which can be difficult to identify on the surface, so this site

can be difficult to distinguish from wet flats and depressions. This site is at risk of exposure to storm surge only under the most extreme conditions.

## Wetland description

This site is not a wetland.

## Soil features

The soils of this site are all primarily clayey or fine-silty in texture, and most are Ultisols formed in deep marine and fluviomarine mineral soil deposits. The soils are acidic and very deep. Soils on this site do not meet hydric criteria. This site commonly abuts ecological sites with hydric soils and the transition between these sites can be exceptionally subtle. The soils on this site are primarily somewhat poorly drained and moderately well drained. In this MLRA, these soils are at risk of inundation during extreme tidal, precipitation, or storm surge events.

Soil series on this site include: Barclay, Craven, Dogue, Dunbar, Eulonia, Fairhope, Lenoir, Nemours, Nevarc, Newflat, Okeetee, Peawick, and Wahee.

Eulonia and Wahee are modal.

**Table 5. Representative soil features**

|                                         |                                                    |
|-----------------------------------------|----------------------------------------------------|
| Parent material                         | (1) Marine deposits<br>(2) Fluviomarine deposits   |
| Surface texture                         | (1) Fine sandy loam<br>(2) Silt                    |
| Drainage class                          | Somewhat poorly drained to moderately well drained |
| Permeability class                      | Moderately slow to moderately rapid                |
| Soil depth                              | 200 cm                                             |
| Surface fragment cover <=3"             | Not specified                                      |
| Surface fragment cover >3"              | Not specified                                      |
| Available water capacity<br>(0-101.6cm) | 10.67 – 15.49 cm                                   |
| Soil reaction (1:1 water)<br>(0-25.4cm) | 0 – 6.5                                            |

|                                                |               |
|------------------------------------------------|---------------|
| Subsurface fragment volume <=3"<br>(0-101.6cm) | 0 – 10 %      |
| Subsurface fragment volume >3"<br>(0-101.6cm)  | Not specified |

**Table 6. Representative soil features (actual values)**

|                                                |                                                    |
|------------------------------------------------|----------------------------------------------------|
| Drainage class                                 | Somewhat poorly drained to moderately well drained |
| Permeability class                             | Moderately slow to moderately rapid                |
| Soil depth                                     | 200 cm                                             |
| Surface fragment cover <=3"                    | 0 %                                                |
| Surface fragment cover >3"                     | 0 %                                                |
| Available water capacity<br>(0-101.6cm)        | 6.1 – 19.3 cm                                      |
| Soil reaction (1:1 water)<br>(0-25.4cm)        | 3.5 – 6.5                                          |
| Subsurface fragment volume <=3"<br>(0-101.6cm) | 0 – 10 %                                           |
| Subsurface fragment volume >3"<br>(0-101.6cm)  | 0 %                                                |

## Ecological dynamics

The most dominant ecological drivers on this site are fire, soil moisture dynamics, tides, and landuse conversion. This site is relatively fertile and nearly all of these locations have been converted to agriculture and other development.

This site tends to occur in locations that are somewhat sheltered from fire, but frequent low intensity fire on a return interval of roughly 2 to 10 years is thought to be necessary to maintain the reference community. This site likely experiences seasonal saturation and seasonal dryness. On this ecological site, longleaf pine uplands is the reference community, because it represents the dominant precolonial forest community. It is probable that longleaf pine uplands were a cultural state maintained by indigenous civilizations but, in most locations today, they no longer dominate the landscape. Historically, the use of fire by indigenous civilizations may have been extensive. Some limited wildfire and prescribed fire occur today, but fire suppression has been the norm since the 20th century.

Across this MLRA, tidal influences and salinization also exert a primary influence on vegetation communities. This site is on relatively high ground in the context of this MLRA, but the entire MLRA is at risk of inundation associated with the most extreme tidal, precipitation, or storm surge events.

(Peat and Allard, 1993; Schafale and Weakley, 1990)

State and transition model

Additional community tables

Table 7. Community 3.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 8. Community 3.2 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 9. Community 3.3 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 10. Community 3.4 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Inventory data references

Data collection and analysis of field data will be performed during the Verification Stage of ESD development.

Other references

Ash, A., E. McDonald, E. Kane, and C. Pories. 1983. Natural and modified pocosins: Literature synthesis and management options. U.S. Fish and Wildlife Services, Dept. Biol., Tech. Rep. FWS/OBS-83/04. U.S. Fish and Wildlife Services, Washington, D.C.

Caldwell, P., M. Vepraskas, J.D. Gregory, R.W. Skaggs, and R.L. Huffman. 2011. Linking Plant Ecology and Long-Term Hydrology to Improve Wetland Restoration Success. Transactions of the ASABE. 54: 2129-2137. DOI: 10.13031/2013.40662

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C.A. Carpenter, W.H. McNab. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. General Technical Report WO-76D. U.S. Department of Agriculture, Forest Service. Washington, D.C.

Dimick, B. P., J. Stucky, W. Wall, M. Vepraskas, T. Wentworth, and C. Arellana. 2010. Plant?soil? hydrology relationships in three Carolina bays in Bladen County, North Carolina, USA. Castanea 75(4): 407?420

Fenneman, N.M., and D.W. Johnson. 1946. Physical divisions of the United States. U.S. Geological Survey, Physiographic Committee. Scale 1:700,000.

Florida Chapter, Soil and Water Conservation Society. 1989. 26 Ecological Communities of Florida. 147 pp.

Florida Natural Areas Inventory (FNAI). 2010. Guide to the natural communities of Florida: 2010 edition. Florida Natural Areas Inventory, Tallahassee, FL.

McNab, W.H.; D.T. Cleland, J.A Freeouf, J.E. Keys Jr., G.J. Nowacki, C.A. Carpenter, comps. 2007. Description of ecological subregions: sections of the conterminous United States [CD-ROM]. Gen. Tech. Report WO-76B. Washington, DC: U.S. Department of Agriculture, Forest Service. 80 pp.

Moritz, C. 2021. Evaluating mitigation sites in Carolina bay wetlands that were previously converted to agriculture. Institutional Repository at North Carolina State University, North Carolina State University, Raleigh, NC, 1–323.

Moritz C., M. Vepraskas, and M. Ricker. 2022. Hydrology and Vegetation Relationships in a Carolina Bay Wetland 15 Years after Restoration. Wetlands. 42. DOI: 10.1007/s13157-022-01530-0.

Nelson, J.B. 1986. The Natural Communities of South Carolina Initial Classification and Description, South Carolina Wildlife and Marine Resources Department, Division of Wildlife and Freshwater Fisheries.

Peet, R.K., and D.J. Allard. 1993. Longleaf Pine Vegetation of the Southern Atlantic and Eastern Gulf Coast Regions: A Preliminary Classification. In Proceedings of the Tall Timbers Fire Ecology Conference, No. 18, The Longleaf Pine Ecosystem: ecology, restoration and management, edited by Sharon M. Hermann, Tall Timbers Research Station, Tallahassee, FL, 1993.

Ross, T.E. 2003. Pocosins and Carolina Bays Compared, The North Carolina Geographer, Volume 11: 22-32

Schafale, M.P., and A.S. Weakley. 1990. Classification of the Natural Communities of North Carolina Third Approximation. North Carolina Natural Heritage Program. 321 pp.

Sharitz R.R., and J.W. Gibbons. 1982. The ecology of southeastern shrub bogs (pocosins) and Carolina bays: a community profile. U.S. Fish and Wildlife Service, Division of Biological Services, Washington, D.C. FWS/OBS-82/04. 93 pp.

Soil Survey Staff. 2023. Web Soil Survey. USDA Natural Resources Conservation Service. <http://websoilsurvey.sc.egov.usda.gov/> (accessed 16 February 2023).

U.S. Department of Agriculture, Natural Resources Conservation Service. 2017. Geomorphic Description System, Version 5.0. Schoeneberger, P.J., and D.A. (eds). USDA-NRCS, National Soil Survey Center, Lincoln, NE.

U.S. Department of Agriculture, Natural Resources Conservation Service. 2022. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. Agriculture Handbook 296.

U.S. Environmental Protection Agency. 2013. Level III and IV ecoregions of the continental United States: Corvallis, Oregon, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000, <https://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>.

## Contributors

Matthew D. Duvall

## Approval

Charles Stemmans, 4/02/2025

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|                                             |                   |
|---------------------------------------------|-------------------|
| Author(s)/participant(s)                    |                   |
| Contact for lead author                     |                   |
| Date                                        | 08/20/2026        |
| Approved by                                 |                   |
| Approval date                               |                   |
| Composition (Indicators 10 and 12) based on | Annual Production |

## Indicators

1. Number and extent of rills:

---

2. Presence of water flow patterns:

---

3. Number and height of erosional pedestals or terracettes:

---

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

5. Number of gullies and erosion associated with gullies:

---

6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---