

Ecological site QX191X01X503

Very Shallow to Moderately Deep Soils on Limestone Plateaus

Last updated: 8/19/2025
Accessed: 10/05/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 191X–High Limestone Plateaus of the Mariana Islands

This MLRA consists of the northern half of Guam and the islands of Saipan, Tinian, Rota, and Aguijan. Topography consists mostly of limestone plateaus. The highest elevations are 1,700 feet (520 meters) on Rota, 1,485 feet (450 meters) on Saipan, 655 feet (200 meters) on Tinian, and 860 feet (260 meters) in the northern part of Guam. The limestone plateaus were uplifted by tectonic activity. Deeply weathered volcanic rock is exposed on some ridges and slopes. Average annual rainfall is 100 inches (2,540 millimeters) on northern Guam and 80 inches (2,030 millimeters) on Saipan, Tinian, Aguijan, and Rota. The rainy season occurs from July through November; the dry season occurs from December through June. Trade winds blow from the northeast. Average annual temperature is 79 degrees F (26 degrees C). Typhoons are frequent. Soils are Alfisols, Andisols, Entisols, Mollisols, or Oxisols. The dominant soil moisture regime is ustic. The soil temperature regime is isohyperthermic. Native vegetation consists of mixed tropical hardwoods; introduced white leadtree or tangantangan (*Leucaena leucocephala*) are abundant. Introduced deer, pigs, goats, and water buffalo are common (USDA-NRCS, 2006). The main human disturbance is clearing land for cultivation. Feral pigs, feral cattle, brown tree snakes, stray dogs (Boonie dogs), and introduced deer are destructive to the native forest (Amand, 2000; CNMI SWARS Council, 2010; Donnegan et al., 2011; Liske-Clark, 2015; Willsey et al., 2019).

Classification relationships

This ecological site occurs within Major Land Resource Area (MLRA) 191 – High Limestone Plateaus of the Mariana Islands.

Ecological site concept

This ecological site occurs on the islands of Guam, Saipan, Tinian, Rota, and Aguijan in the Mariana Islands. It occurs on nearly level to steep (0 to 30 percent slopes) limestone plateaus at elevations ranging from 10 to 1,476 feet (3 to 450 meters) (USDA-SCS, 1988; USDA-SCS, 1989). Soils range from very shallow to the shallower range of the moderately deep class. They are well drained Mollisols (Argiustolls and Haplustolls), Alfisols, and Entisols that formed in sediment or residuum over coralline limestone or argillaceous limestone. Some phases consist of fill deposited over the natural soil surface. Soil temperature regimes are isohyperthermic; soil moisture regimes are ustic. Average annual precipitation is 88 inches (2,240 millimeters) and ranges from 78 to 95 inches (1,980 to 2,410 millimeters) (Prism, 2006). Water runoff is predominately low; permeability is slow to moderately rapid. Effective rooting depth ranges from 2 to 20 inches (5 to 50 centimeters); some roots occur in cracks in the underlying limestone. Available water-holding capacity is 2 to 3 inches (3 to 10 centimeters) in the upper 40 inches (100 centimeters) of soil (USDA-SCS, 1988; USDA-SCS, 1989). Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native forest. Tangantangan (*Leucaena leucocephala*) is abundant in places (Amidon et al., 2017; Falanruw et al., 1989; Fosberg, 1960; Liu and Fischer, 2008; Stone, 1970; Wagner and Grether, 1948; Willsey et al., 2019).

Associated sites

QX191X01X506	Somewhat Poorly and Poorly Drained Valley Bottoms and Coastal Plains Soils of QX191X01X506 (The Somewhat Poorly and Poorly Drained Valleys and Alluvial Coastal Plains Ecological Site) are in broad river valley bottoms and alluvial coastal plains. They are poorly to somewhat poorly drained. They formed in alluvium or beach deposits, have aquic moisture regimes, shallow water tables, areas of slightly brackish groundwater, and are occasionally flooded. They have moderately high (4 inches) or high (5 inches) water holding capacity. Most of the area is vegetated by grasses, forbs, and wetland forest. Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) occur on limestone plateaus. They are well drained, very shallow to moderately deep (8 to 26 inches) clays, clay loams and very gravelly sandy loams. Available water-holding capacity is very low (2 to 3 inches). Sub surface soils contain 0 to 13 percent rock fragments greater than 3 inches. Much of the area is farmed. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest.
QX191X01X004	Somewhat Poorly and Poorly Drained Limestone Basins Soils of QX191X01X004 (The Somewhat Poorly and Poorly Drained Limestone Basins Ecological Site) occur on basin floors; they rarely flood or pond. They are moderately deep and deep, somewhat poorly and poorly drained, have low to moderate (5 inches) available water holding capacity, and somewhat neutral to slightly basic soils (pH ranges from 5.8 to 7.0). Most of the area is vegetated by water- tolerant grasses and forest. Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) occur on plateaus. They are well drained, very shallow to moderately deep (8 to 26 inches) clays, clay loams and very gravelly sandy loams. Available water-holding capacity is very low (2 to 3 inches). Sub surface soils contain 0 to 13 percent rock fragments greater than 3 inches. Much of the area is farmed. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest.
QX192X01X501	Somewhat Poorly Drained Alluvium Soils of QX192X01X501 (The Somewhat Poorly Drained Alluvium Ecological Site) occur in stream channels, depressions, and seeps on volcanic uplands. The soils, which formed in alluvium from volcanic tuff and tuff breccia, are somewhat poorly drained, have effective rooting depths greater than 60 inches, high available water holding capacity, shallow water tables during the rainy season, and are occasionally flooded. Most of the area is vegetated by wetland forest, grasses, and sedges. Soils of Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) formed in fine mineral materials over porous coralline or argillaceous limestone. Soils have ustic moisture regimes, deep water tables, only fresh groundwater, and no flooding or ponding. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest.
QX191X01X005	Very Poorly Drained Muck Soils of QX191X01X005 (The Very Poorly Drained Muck Ecological Site) occurs in depressions on nearly level (0 to 2 percent) slopes at elevations ranging from 0 to 20 feet. The surface soil is 8 inches of muck over mucky clay loam. The soil moisture regimes are aquic; the soils are wet at the surface (0 inches). Groundwater varies from fresh to saline (EC 8 to 16 dS/m) depending on location. Flooding is frequent. Most of the area is vegetated by marsh grasses and, in a few locations, small stands of hibiscus (<i>Hibiscus tiliaceus</i>) or mangrove forest. Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) formed in fine mineral materials over porous coralline or argillaceous limestone. Soils have ustic moisture regimes, deep water tables, only fresh groundwater, and no flooding or ponding. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest.

QX192X01X001	Moderately Deep Oxidic Soils on Volcanic Uplands Soils of QX192X01X001 (The Very Deep Oxidic Soils on Volcanic Uplands Ecological Site) formed in volcanic materials and have effective rooting depths greater than 60 inches, moderately high available water holding capacity, low base saturation, low pH, and aluminum toxicity. Most of the area is vegetated by Pacific silver grass or swordgrass (<i>Miscanthus floridulus</i>). Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) formed in fine materials over porous coralline or argillaceous limestone. Effective rooting depth is 2.5 to 26 inches, available water holding capacity is very low to low, base saturation is high, pH is high, and there is no aluminum toxicity. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest. Tangantangan is abundant in places.
QX192X01X002	Deep Alfic Soils on Volcanic Uplands Soils of QX192X01X002 (The Deep Alfic Soils on Volcanic Uplands Ecological Site) formed in volcanic materials and have effective rooting depths greater of 40 to 60 inches, low available water holding capacity, fairly high base saturation, pH above 5.0, and no aluminum toxicity. Most of the area is vegetated by forest, grasses, and forbs. Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) formed in fine materials over porous coralline or argillaceous limestone. Effective rooting depth is 2 to 20 inches, available water holding capacity is very low to low, base saturation is high, pH is high, and there is no aluminum toxicity. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest. Tangantangan is abundant in places.
QX192X01X003	Shallow Soils on Volcanic Uplands Soils of QX192X01X003 (The Shallow and Shallow Soils on Volcanic Uplands Ecological Site) formed in volcanic materials, have effective rooting depths from 4 to 15 inches, and have very low available water holding capacity. Most of the area is vegetated by grasses, forbs, and forest. Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) formed in fine materials over porous coralline or argillaceous limestone. Effective rooting depth is 2 to 20 inches and available water holding capacity is very low to low. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest. Tangantangan is abundant in places.
QX191X01X505	Very Low Available Water Capacity Soils on Limestone Plateaus and Escarpments Soils of QX191X01X505 (The Very Low Available Water Capacity Soils on Limestone Plateaus and Escarpments Ecological Site) occur on plateaus. They are well drained, shallow, very-cobbly clays and cobbly clay loams with very low (1 inch) available water holding capacity. Sub surface soils contain 13 to 40 percent rock fragments greater than 3 inches. Most of the area is vegetated by native forest. Soils of QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) also occur on limestone plateaus. They are well drained, very shallow to moderately deep (8 to 26 inches) clays, clay loams, and very gravelly sandy loams. Available water-holding capacity is very low (2 to 3 inches). Sub surface soils contain 0 to 13 percent rock fragments greater than 3 inches. Much of the area is farmed. Areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest.

Similar sites

QX191X01X502	Deep and Very Deep Soils on Limestone Plateaus QX191X01X502 (Deep and Very Deep Soils on Limestone Plateaus Ecological Site) formed in fine materials over porous coralline or argillaceous limestone. Effective rooting depth is 40 to 60 inches or deeper and available water holding capacity is moderate. Most of this ecological site is used for farmland; areas not farmed are vegetated by mostly introduced grass, forb, and tree species. Tangantangan (<i>Leucaena leucocephala</i>) is abundant in places. Soils of QX191X01X503 (Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site) formed in fine materials over porous coralline or argillaceous limestone. Effective rooting depth is 2 to 20 inches and available water holding capacity is very low to low. For both ecological sites, areas not farmed are vegetated by mostly introduced grass, forb, and tree species, with some areas of native upland forest. Tangantangan is abundant in places.
---------------------	---

Table 1. Dominant plant species

Tree	(1) <i>Intsia bijuga</i> (2) <i>Guamia mariannae</i>
Shrub	(1) <i>Cycas circinalis</i>
Herbaceous	Not specified

Legacy ID

F191XY503MP

Physiographic features

This ecological site primarily occurs on limestone plateaus. A few soil map unit components occur on basin floor (USDA-SCS, 1988; USDA-SCS, 1989). The depth to water is greater than 72 inches.

Table 2. Representative physiographic features

Landforms	(1) Plateau
Runoff class	Low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	0 – 450 m
Slope	0 – 30 %
Water table depth	180 cm
Aspect	Aspect is not a significant factor

Table 3. Representative physiographic features (actual ranges)

Runoff class	Very low to very high
Flooding frequency	Not specified
Ponding frequency	Not specified

Elevation	Not specified
Slope	0 – 60 %
Water table depth	Not specified

Climatic features

Summary for this Ecological Site

Rainfall statistics were determined from PRISM rainfall raster data (PRISM, 2006). Representative (20th and 80th percentiles) values for mean annual precipitation range from 78 to 95 inches (1,980 to 2,410 millimeters) while actual (10th and 90th percentiles) values for mean annual precipitation range from 75 to 96 inches (1,910 to 2,440 millimeters). Extreme values range from 69 to 104 inches (1,750 to 2,640 millimeters). The average annual precipitation is 88 inches (2,240 millimeters) and the median annual average precipitation is 93 inches (2,360 millimeters).

Temperature statistics were determined from PRISM temperature raster data (PRISM, 2006). Representative (20th and 80th percentiles) values for mean annual temperatures range from 80 to 81 degrees F (26 to 27 degrees C) while actual (10th and 90th percentiles) values for mean annual temperatures also range from 80 to 81 degrees F (26 to 27 degrees C). Extreme values range from 75 to 82 degrees F (24 to 28 degrees C). The average annual temperature is 81 degrees F (27 degrees C), and the median annual temperature is also 81 degrees F (27 degrees C).

The data presented in the climate normals tables below are from select climate stations in Guam and Saipan. I used the data from these stations because they are representative of the spatial distribution of this ecological site and because they provide a reasonable approximation of the PRISM data presented above, e.g. the average annual precipitation is 91 inches (2,310 millimeters), and the average annual temperature is 80 degrees F (26 degrees C).

General Principles

The climate is uniformly warm and humid throughout the year. Afternoon temperatures typically are about 86 degrees F (30 degrees C); nighttime temperatures are about 68 F (20 C). Relative humidity is 65 to 75 percent in the afternoon to 85 to 100 percent at night. Though temperature and humidity vary only slightly throughout the year, rainfall and wind conditions vary markedly. There are two main seasons, the dry season from December through June and the rainy season from mid-July to mid-November. Moisture deficit occurs between January and June. Mean annual rainfall ranges from about 98 inches (2,490 millimeters) on the windward (eastern) side of the higher mountains to about 79 inches (2,005 millimeters) along the coast of the western side of the islands. On average, about 15 percent of the annual rainfall occurs during the dry season and 55 percent during the rainy season (Fosberg, 1960; USDA-SCS, 1988; USDA-SCS, 1989).

Throughout the year, the dominant winds are the trade winds that blow from the east or northeast. The trade winds are strongest and most constant during the dry season, when windspeeds of 15 to 25 mph (25 to 40 kph) are common. Coastal areas that have east, and northeast exposures are subject to salt spray and buffeting winds (Fosberg, 1960; USDA-SCS, 1988; USDA-SCS, 1989).

During the rainy season the trade winds may break down and be replaced by a weak, westerly monsoon influence that brings heavy showers or steady and sometimes torrential rains. The islands lie in the path of typhoons from the southeast and east. They bring heavy rains and violent winds that may result in a surge of water onto low-lying coastal areas. They occur most frequently during the latter half of the year. The chance of having one or more typhoons pass close to the islands in any particular year is about once in three years. The chance of having a typhoon move directly across an island is about once in eight years (Fosberg, 1960; USDA-SCS, 1988; USDA-SCS, 1989).

Table 4 Representative climatic features

Frost-free period (characteristic range)	370 days
Freeze-free period (characteristic range)	370 days
Precipitation total (characteristic range)	1,980-2,410 mm

Frost-free period (actual range)	370 days
Freeze-free period (actual range)	370 days
Precipitation total (actual range)	1,910-2,440 mm
Frost-free period (average)	370 days
Freeze-free period (average)	370 days
Precipitation total (average)	2,240 mm

- (1) CAPITOL HILL 1 [CQC00914080], MP
- (2) ANDERSEN AFB GUAM [GQC00914025], GU
- (3) GUAM INTL AP [GQW00041415], GU

Influencing water features

Number of National Wetland Inventory (NWI) features overlapping ecological site: Riverine (156), freshwater emergent wetlands (123), freshwater forested/shrub wetlands (78), estuarine and marine wetland (48), freshwater pond (21), and estuarine and marine deepwater (20) (USFWS, 2023).

Soil features

The soil components associated with this ecological site are Banaderu (Mollisols), Chinen (Mollisols), Dandan (Inceptisols), Pulantat (Mollisols), Guam (Entisols), and Chacha (Alfisols) (USDA-SCS, 1988; USDA-SCS, 1989). Soil temperature regimes are isohyperthermic; soil moisture regimes are ustic. They are very shallow to the shallower end of moderately deep, and well drained. The underlying limestone is porous. Two phases of Chinen series consist of very gravelly sandy loam fill deposited over the natural soil surface.

Table 5. Representative soil features

Parent material	(1) Residuum – coral limestone (2) Residuum – argillaceous limestone
Surface texture	(1) Clay (2) Clay loam (3) Cobbly clay loam
Family particle size	(1) Fine (2) Clayey
Drainage class	Well drained

Permeability class	Slow to moderate
Depth to restrictive layer	10 – 50 cm
Soil depth	10 – 50 cm
Surface fragment cover <=3"	0 – 10 %
Surface fragment cover >3"	0 – 10 %
Available water capacity (0-101.6cm)	3.56 – 9.91 cm
Calcium carbonate equivalent (0-101.6cm)	0 – 100 %
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-25.4cm)	6.5 – 7.9
Subsurface fragment volume <=3" (0-101.6cm)	0 – 20 %
Subsurface fragment volume >3" (0-101.6cm)	0 – 10 %

Ecological dynamics

The main human disturbance is clearing land for cultivation. Much of the area is currently used for farmland. Feral pigs, feral cattle, and introduced deer are destructive to the forest (CNMI SWARS Council, 2010; Donnegan et al., 2011; Liske-Clark, 2015; Willsey et al., 2019).

The main natural disturbance is strong storms that can damage or kill vegetation by high wind speeds. Natural fire is rare in forested areas due to the low amounts of fine fuels on the ground (CNMI SWARS Council, 2010; Donnegan et al., 2011; Liske-Clark, 2015; Willsey et al., 2019).

State and transition model

Figure 7. State-and-Transition Model for F191XY503MP (Very Shallow to Moderately Deep Soils on Limestone Plateaus)

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 7. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 11. Community 4.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 12. Community 5.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Other references

References for QX191X01X503 (The Very Shallow to Moderately Deep Soils on Limestone Plateaus Ecological Site)

Amand, A. (2000). *Boiga irregularis* (Brown Tree Snake) on Guam and Its Effect on Fauna. *Restoration and Reclamation Review*, 6 1-6.

Amidon F., Metevier, M., and Miller, S. E. (2017). *Vegetation Mapping of the Mariana Islands: Commonwealth of the Northern Mariana Islands and Territory of Guam*. US Fish and Wildlife Service and Pacific Islands Climate Change Cooperative. Final Report November 2017.

CNMI SWARS Council. (2010). *Commonwealth of the Northern Mariana Islands (CNMI) Statewide Assessment and Resource Strategy 2010-2015+*. CNMI Forestry.

Donnegan, J. A., Butler, S. L., Kuegler, O., and Hiserote, B. A. (2011). *Commonwealth of the Northern Mariana Islands' Forest resources, 2004*. Resource. Bull. PNW-RB-261. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 40 p.

Falanruw, M. C., Cole, T. G., and Ambacher, A. H. (1989). *Vegetation survey of Rota, Tinian, and Saipan, Commonwealth of the Northern Mariana Islands*. Resource. Bull. PSW-27. Berkeley, CA: Pacific Southwest Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture.

Fosberg, F. R. (1960). *The Vegetation of Micronesia. I. General descriptions, the vegetation of the Marianas Islands, and a detailed consideration of the vegetation of Guam*. *Bulletin of the American Museum of Natural History* 199:1.

Liske-Clark, J. (2015). *Wildlife Action Plan for the Commonwealth of the Northern Mariana Islands, 2015-2025*. CNMI DLRN-Division of Fish and Wildlife, Saipan, MP.

Liu, Z. and Fischer, L. (2008). *Commonwealth of the Northern Mariana Islands Vegetation Mapping Using Very High Spatial Resolution Imagery*. USDA Forest Service Pacific Southwest Region, Forest Health Protection, McClellan, CA.

PRISM Climate Group, Oregon State University. (2006). <https://prism.oregonstate.edu>, data created 4 Feb 2014, accessed 09 Oct 2024.

Pacific Islands Project: Average monthly precipitation, minimum and maximum temperature, relative humidity, and mean dew point temperature for the period 1971-2000 (completed 2006). Covers Hawaii, Kosrae, Manua, Pohnpei, Tutuila, Guam, CNMI, and Palau. Project sponsored by USDA Natural Resources Conservation Service.

Stone, B. C. (1970). The Flora of Guam. A Manual for the Identification of the Vascular Plants of the Island. Micronesica – Journal of the University of Guam, vol. 6, July 1970.

USDA-NRCS. (2006). Major Land Resource Regions. USDA Agriculture Handbook 296. <http://soils.usda.gov/MLRAExplorer>

USDA-Soil Conservation Service. (1989). Soil Survey of the Islands of Aguijan, Rota, Saipan, and Tinian, Commonwealth of the Northern Mariana Islands.

USDA-Soil Conservation Service. (1988). Soil Survey of Territory of Guam.

U. S. Department of Interior, Fish & Wildlife Service. (2023). Download seamless wetlands data by state. National Wetlands Inventory website. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C. Accessed April 24, 2024. [<https://www.fws.gov/program/national-wetlands-inventory/download-state-wetlands-data>].

Wagner, W. H. and Grether, D. F. (1948). The Pteridophytes of Guam. Occasional Papers of Bernice P. Bishop Museum, volume XIX, number 2. Honolulu, Hawaii.

Willsey, T., Kwon, J. A., Reeves, M. K., Amidon, F., and Miller, S. E. (2019). Mariana Islands Forest. Encyclopedia of the World's Biomes. <https://doi.org/10.1016/B978-0-12-409548-9.12012-3>

Contributors

David Clausnitzer PhD
John Proctor
Ann Tan
Sarah Quistberg
Kendra Moselely
Brendan Brazee
Amy Koch
Jill Fick-Beaton
Daniel Block
Daniel Bowman

Approval

Curtis Talbot, 8/19/2025

Acknowledgments

Assistance, advice, review, or insights: Michael Constantinides, NRCS-PIA Jennifer Higashino, USFWS and NRCS Michael Kolman, NRCS Bart Lawrence, NRCS Pamela Sablan, NRCS

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	08/19/2025

Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
