

# Ecological site F236XY116AK

## Boreal Forest Loamy Wet Slopes

Last updated: 2/13/2024  
Accessed: 10/03/2026

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 236X–Bristol Bay-Northern Alaska Peninsula Lowlands

The Bristol Bay-Northern Alaska Peninsula Lowland Major Land Resource Area (MLRA 236) is located in Western Alaska. This MLRA covers approximately 19,500 square miles and is defined by an expanse of nearly level to rolling lowlands, uplands and low to moderate hills bordered by long, mountain footslopes. Major rivers include the Egegik, Mulchatna, Naknek, Nushagak, and Wood River. MLRA 236 is in the zone of discontinuous permafrost. It is primarily in areas with finer textured soils on terraces, rolling uplands and footslopes. This MLRA was glaciated during the early to middle Pleistocene. Moraine and glaciofluvial deposits cover around sixty percent of the MLRA. Alluvium and coastal deposits make up a large portion of the remaining area (Kautz et al., 2012; USDA, 2006). Climate patterns across this MLRA shift as one moves away from the coast. A maritime climate is prominent along the coast, while continental weather, commonly associated with Interior Alaska, is more influential inland. Across the MLRA, summers are general short and warm while winters are long and cold. Mean annual precipitation is 13 to 50 inches, with increased precipitation at higher elevations and areas away from the coast. Mean annual temperatures is between 30 and 36 degrees F (USDA, 2006). The Bristol Bay-Northern Alaska Peninsula MLRA is principally undeveloped wilderness. Federally managed land includes parts of the Katmai and Aniakchak National Parks, and the Alaska Peninsula, Becharof, Togiak and Alaska Maritime National Wildlife Refuges. The MLRA is sparsely populated. Principal communities include Dillingham, Naknek, and King Salmon. Commercial fishing in Bristol Bay and the Bering Sea comprises a major part of economic activity in the MLRA. Other land uses include subsistence activities (fishing, hunting, and gathering) and sport hunting and fishing (USDA, 2006).

### Ecological site concept

This boreal ecological site is on linear to convex plains slopes. Site elevation is between 40 and 150 feet above sea level. Slopes are nearly level to gentle (0 – 4 percent). Site hydrology and soil development and fertility shape the vegetation on this landform. Ponding is an occasional and brief event. A shallow (17 – 33 inches) water table is present in June, restricting plant species during the important early growing season. While hydrologically influenced, the understory of this site is less restricted by wet soils than other similar sites, like R236XY117AK. Soil development is weak, characterized by an ochric epipedon, but is likely balanced by the presence of volcanic ash deposits that may increase soil fertility. The reference state supports two communities. The reference plant community is characterized as a mixed open forest (Viereck et al., 1992). It is composed of an open canopy of white spruce and paper birch with an open understory of facultative upland to facultative wetland shrub and herbaceous species. The second community is a closed willow scrubland.

### Associated sites

|             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|-------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| F236XY117AK | <p><b>Boreal Forest Wet Loamy Plain Drainageways</b></p> <p>Both are forested sites on wet plain soils. F236XY117AK is associated with wetter soils that are somewhat poorly drained and support a shallow water table throughout the growing season. Vegetation is restricted to facultative wet to obligate species in the understory. This site is moderately well drained with a water table in June. While hydrologically influenced, the understory is less restricted by wet soils.</p> |
|-------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

|                    |                                                                                                                                                                                                                                                                                                              |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>F236XY115AK</b> | <b>Boreal Forest Loamy Moist Slopes</b><br><br>Both sites are on plains and hills. F236XY115AK describes similar areas that are not adjacent to wetlands. Wetland hydrology influences the site characteristics, disturbances, and supported vegetation of F236XY116AK, differentiating it from F236XY115AK. |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

### Similar sites

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>F236XY117AK</b> | <b>Boreal Forest Wet Loamy Plain Drainageways</b><br><br>Both are forested sites on wet plain soils. F236XY117AK is associated with wetter soils that are somewhat poorly drained and support a shallow water table throughout the growing season. Vegetation is restricted to facultative wet to obligate species in the understory. This site is moderately well drained with a water table in June. While hydrologically influenced, the understory is less restricted by wet soils. |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**Table 1. Dominant plant species**

|            |                                                                      |
|------------|----------------------------------------------------------------------|
| Tree       | (1) <i>Betula papyrifera</i><br>(2) <i>Picea glauca</i>              |
| Shrub      | (1) <i>Spiraea stevenii</i><br>(2) <i>Vaccinium vitis-idaea</i>      |
| Herbaceous | (1) <i>Dryopteris expansa</i><br>(2) <i>Calamagrostis canadensis</i> |

### Physiographic features

This site is on linear to convex slopes of plains. Elevation typically ranges from 40 to 150 feet above sea level. Slopes are nearly level to gentle (0 – 4 percent). This site is found at all aspects. Ponding is an occasional and brief occurrence and a shallow (17 – 33 inches) water table is present in June.

**Table 2. Representative physiographic features**

|                            |                                         |
|----------------------------|-----------------------------------------|
| Geomorphic position, flats | (1) Talf<br>(2) Rise                    |
| Landforms                  | (1) Plains > Plain<br>(2) Plains > Rise |
| Runoff class               | Low to medium                           |
| Flooding frequency         | None                                    |
| Ponding duration           | Brief (2 to 7 days)                     |
| Ponding frequency          | Occasional                              |

|                   |                            |
|-------------------|----------------------------|
| Elevation         | 10 – 50 m                  |
| Slope             | 0 %                        |
| Water table depth | 40 – 80 cm                 |
| Aspect            | W, NW, N, NE, E, SE, S, SW |

**Table 3. Representative physiographic features (actual ranges)**

|                    |                     |
|--------------------|---------------------|
| Runoff class       | Low to high         |
| Flooding frequency | None                |
| Ponding duration   | Brief (2 to 7 days) |
| Ponding frequency  | None to occasional  |
| Elevation          | 0 – 300 m           |
| Slope              | 0 – 20 %            |
| Water table depth  | 40 – 100 cm         |

### Climatic features

The climate of this site reflects that of the MLRA, which is described as maritime polar (EPA, 2013). Temperatures are moderated by the nearby Bristol Bay and northern Pacific bodies of water. Annual precipitation ranges from 21 – 34 inches with approximately 40 percent occurring during the June-September growing season (PRISM, 2018).

**Table 4 Representative climatic features**

|                                            |             |
|--------------------------------------------|-------------|
| Frost-free period (characteristic range)   | 80-100 days |
| Freeze-free period (characteristic range)  | 70-90 days  |
| Precipitation total (characteristic range) | 530-860 mm  |
| Frost-free period (actual range)           | 80-100 days |
| Freeze-free period (actual range)          | 70-90 days  |

|                                    |              |
|------------------------------------|--------------|
| Precipitation total (actual range) | 380-1,040 mm |
| Frost-free period (average)        | 90 days      |
| Freeze-free period (average)       | 80 days      |
| Precipitation total (average)      | 740 mm       |

### Influencing water features

Site hydrology shapes the vegetation found in the forest understory. However, due to its landscape position, this site is not influenced by wetland or riparian water features. Precipitation and throughflow are the main sources of water.

### Soil features

Soils are weakly developed and weakly weather Spodosols (Soil Survey Staff, 2013). Soils are very deep and moderately well drained. They support a cryic temperature regime and an udic moisture regime. Parent material is organic material over volcanic ash over loess.

Soil characteristics affecting vegetation include hydrology, soil development and the presence of volcanic ash. A moderately deep water table is present in June and aquic conditions are present beginning at 33 inches. Wet soils restrict which plant species can grow here during the important early growing season. Soil development is weak, characterized by an ochric epipedon. Weak soil development typically restricts plant recruitment. However, the presences of andic soil properties, the result of volcanic ash deposits, likely increase soil fertility and the resulting vegetation.

Correlated soil components in MLRA 236: Muklung, Oxyaquic Haplocryods

Table 5. Representative soil features

|                             |                               |
|-----------------------------|-------------------------------|
| Parent material             | (1) Loess<br>(2) Volcanic ash |
| Surface texture             | (1) Highly organic silt loam  |
| Drainage class              | Moderately well drained       |
| Permeability class          | Slow                          |
| Soil depth                  | 150 cm                        |
| Surface fragment cover <=3" | Not specified                 |
| Surface fragment cover >3"  | Not specified                 |

|                                                         |                |
|---------------------------------------------------------|----------------|
| Available water capacity<br>(0-25.4cm)                  | 5.33 – 6.35 cm |
| Soil reaction (1:1 water)<br>(0-25.4cm)                 | 4.2 – 5.8      |
| Subsurface fragment volume ≤3"<br>(Depth not specified) | Not specified  |
| Subsurface fragment volume >3"<br>(Depth not specified) | Not specified  |

Table 6. Representative soil features (actual values)

|                                                         |                         |
|---------------------------------------------------------|-------------------------|
| Drainage class                                          | Moderately well drained |
| Permeability class                                      | Slow                    |
| Soil depth                                              | 150 cm                  |
| Surface fragment cover ≤3"                              | 0 %                     |
| Surface fragment cover >3"                              | 0 %                     |
| Available water capacity<br>(0-25.4cm)                  | 5.33 – 7.62 cm          |
| Soil reaction (1:1 water)<br>(0-25.4cm)                 | 4.2 – 5.9               |
| Subsurface fragment volume ≤3"<br>(Depth not specified) | 0 – 20 %                |
| Subsurface fragment volume >3"<br>(Depth not specified) | 0 %                     |

## Ecological dynamics

The boreal upland plains and hills support several ecological sites. The swales and areas near the Nushagak River support unique sites, primarily due to differences in the soils, disturbances, and local climate.

F236XY116AK is in linear to convex areas of upland plains. It generally is at the edge of tree islands and is surrounded by low-lying wetlands. The soils and hydrology in these areas support a unique ecological site. The annual precipitation is 24 to 55 inches, and the annual frost-free period is 85 to 140 days.

Spatial patterns in soil and site hydrology create two communities. The typical soil in this site supports a aquic conditions below 33 inches and a moderately deep water table in June. This soil supports a mixed forest canopy, while the understory is a mix of facultative upland to facultative wet wetland species. Low-lying areas on this site have wetter soils. Wetter soils are typically along the tree island edges, though they can be within the tree island. These wet areas do not support trees and the vegetation is comprised of facultative to obligate wetland species.

The communities on this site are stable. The communities on this site are primarily shaped by soil wetness. Site hydrology appear to be stable over short periods. Site hydrology is typically regulated by the surrounding wetland. Long term changes to wetland hydrology may influence this site, causing a community shift.

Willows are likely browsed by moose throughout the year. This does not appear to affect the ecological processes of the site.

The information in this Ecological Dynamics section, including the state-and-transition model (STM), was developed based on current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Table 8. Community 1.2 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production () | Foliar Cover (%) |
|-------|-------------|--------|-----------------|----------------------|------------------|
|-------|-------------|--------|-----------------|----------------------|------------------|

Inventory data references

Modal points for Community 1.1 07CS11204 07CS12004 07CS17705 07SS08601 07SS11305 07SS12506 08SS15301 Modal points for community 1.2 07SS06209 07CS15204 10SS01804

References

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Other references

Hook, D., and R.M.M. Crawford. 1978. Plant life in anaerobic environments. Ann Arbor Science Publishers, Ann Arbor, MI.

Jackson, M.B., D.D. Davies, and H. Lambers (editors). 1991. Plant life under oxygen deprivation: Ecology, physiology, and biochemistry. SPB Academic Publication, The Hague, Netherlands.

Kautz, D.R., P. Taber, and S. Nield (editors). 2004. Land resource regions and major land resource areas of Alaska. U.S. Department of Agriculture, Natural Resources Conservation Service, Palmer, AK. Revised 2012.

Schoeneberger, P.J., and D.A. Wysocki. 2012. Geomorphic description system. Version 4.2. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. Field book for describing and sampling soils. Version 3.0. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Soil Science Division Staff. 2017. Soil survey manual. Ditzler, C., K. Scheffe, and H.C Monger, editors. U.S. Department of Agriculture Handbook 18. Government Printing Office, Washington, D.C.

U.S. Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

US Environmental Protection Agency (EPA). Level III Ecoregions of the Conterminous United States. UP ESP Office of Research and Development. Corvallis, OR. <http://edg.epa.gov/>. Created 16 Apr 2013. Accessed 20 Mar 2023.

Vartapetian, B.B., and M.B. Jackson. 1996. Plant adaptations to anaerobic stress. *Annals of Botany* 79 (Supplement A): 3-20.

Viereck, L.A., C.T. Dyrness, A.R. Batten, and K.J. Wenzlick. 1992. The Alaska vegetation classification. Gen. Tech. Rep. PNW-GTR-286. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 278 p. vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station General Technical Report PNW-GTR-286. Portland, OR.

## Contributors

Phil Barber  
Stephanie Schmit  
Michael Margo  
Sue Tester  
Jamin Johanson  
Steff Shoemaker

## Approval

Kirt Walstad, 2/13/2024

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|                                             |                   |
|---------------------------------------------|-------------------|
| Author(s)/participant(s)                    |                   |
| Contact for lead author                     |                   |
| Date                                        | 10/03/2026        |
| Approved by                                 |                   |
| Approval date                               |                   |
| Composition (Indicators 10 and 12) based on | Annual Production |

## Indicators

### 1. Number and extent of rills:

---

2. Presence of water flow patterns:

---

3. Number and height of erosional pedestals or terracettes:

---

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

5. Number of gullies and erosion associated with gullies:

---

6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---