

Ecological site F236XY202AK

Boreal Forest Frozen Loamy Slopes

Last updated: 2/13/2024
Accessed: 09/26/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 236X–Bristol Bay-Northern Alaska Peninsula Lowlands

The Bristol Bay-Northern Alaska Peninsula Lowland Major Land Resource Area (MLRA 236) is located in Western Alaska. This MLRA covers approximately 19,500 square miles and is defined by an expanse of nearly level to rolling lowlands, uplands and low to moderate hills bordered by long, mountain footslopes. Major rivers include the Egegik, Mulchatna, Naknek, Nushagak, and Wood River. MLRA 236 is in the zone of discontinuous permafrost. It is primarily in areas with finer textured soils on terraces, rolling uplands and footslopes. This MLRA was glaciated during the early to middle Pleistocene. Moraine and glaciofluvial deposits cover around sixty percent of the MLRA. Alluvium and coastal deposits make up a large portion of the remaining area (Kautz et al., 2012; USDA, 2006). Climate patterns across this MLRA shift as one moves away from the coast. A maritime climate is prominent along the coast, while continental weather, commonly associated with Interior Alaska, is more influential inland. Across the MLRA, summers are general short and warm while winters are long and cold. Mean annual precipitation is 13 to 50 inches, with increased precipitation at higher elevations and areas away from the coast. Mean annual temperatures is between 30 and 36 degrees F (USDA, 2006). The Bristol Bay-Northern Alaska Peninsula MLRA is principally undeveloped wilderness. Federally managed land includes parts of the Katmai and Aniakchak National Parks, and the Alaska Peninsula, Becharof, Togiak and Alaska Maritime National Wildlife Refuges. The MLRA is sparsely populated. Principal communities include Dillingham, Naknek, and King Salmon. Commercial fishing in Bristol Bay and the Bering Sea comprises a major part of economic activity in the MLRA. Other land uses include subsistence activities (fishing, hunting, and gathering) and sport hunting and fishing (USDA, 2006).

Classification relationships

Alaska Vegetation Classification: Needleleaf woodland (I.A.3 - level III) / Black spruce open forest (I.A.3.d - level IV) (Viereck et al., 1992)
Circumboreal Vegetation Map – Alaska-Yukon Region: Southern Alaska Spruce-Birch-Herb Forests (Jorgensen and Meidinger, 2015)
BioPhysical Settings: 7616220 - Western North American Boreal Black Spruce Wet-Mesic Slope Woodland (LANDFIRE biophysical settings, 2009)

Ecological site concept

This boreal ecological site is on mountain backslopes and footslopes. It is typically at elevations between 400 and 1,500 feet. Soils contain permafrost and are poorly drained. Soil hydrology, low soil fertility caused by soil acidity, and a fire regime shape the vegetation on this landform. The reference state supports three communities. The reference plant community is characterized as a black spruce woodland (Viereck et al., 1992). It is composed of a black spruce canopy with an open understory of ericaceous and other woody shrubs and mosses. Post-fire communities are typically absent a forest overstory and are comprised of extant shrubs and colonizing trees.

Associated sites

R236XY107AK	Western Alaska Maritime Scrub Gravelly Drainages R236XY107AK describes scrub communities in drainages and swales. These hydrologically influenced areas are typically too wet to support trees.
-------------	--

F236XY115AK	Boreal Forest Loamy Moist Slopes F236XY115AK describes mixed forests on rounded mountain backslopes. This ecological site is associated with well drained soils, which are more suitable for trees such as white spruce and paper birch. The soil in F236XY202AK is too wet and cold to support those species.
F236XY201AK	Boreal Open Forest Wet Loamy Warm Mountain Slopes F236XY201AK is associated with warmer soils than F236XY202AK. This ecological site is found primarily at warmer aspects and is associated with soils with a large percentage of subsurface rock fragments, which help transport heat into the soil profile. These conditions allow for the growth of white spruce, which is typically restricted from the permafrost soils of F236XY202AK.

Similar sites

F236XY201AK	Boreal Open Forest Wet Loamy Warm Mountain Slopes These ecological sites are both associated with poorly drained soils on mountain backslopes and footslopes. F236XY201AK is restricted to warmer soils, due to southerly aspects and greater than 50% subsurface rock fragments. F236XY201AK may support black spruce in areas, particularly as a result of local conditions and fire disturbance(s). It can be recognized by the absence of permafrost and the presence of other conifer and broadleaf tree species.
--------------------	---

Table 1. Dominant plant species

Tree	(1) <i>Picea mariana</i>
Shrub	(1) <i>Betula nana</i> (2) <i>Empetrum nigrum</i>
Herbaceous	(1) <i>Carex</i>

Physiographic features

This site is on rugged and glaciated mountain backslopes and toeslopes. Elevation typically ranges from 400 to 1,500 feet (2,200 feet maximum) above sea level. Slopes are usually gently to strongly sloped (4 – 14 percent). This site is found at all aspects. A water table is present at the beginning of the growing season between March and June. Ponding and flooding do not occur.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope
Runoff class	Low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	120 – 460 m

Slope	0 – 10 %
Water table depth	20 – 150 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	10 – 670 m
Slope	0 – 30 %
Water table depth	20 – 150 cm

Climatic features

The climate of this site reflects that of the MLRA, which is described as maritime polar (EPA, 2013). Temperatures are moderated by the nearby Bristol Bay and northern Pacific bodies of water. Annual precipitation ranges from 21 – 34 inches with approximately 40 percent occurring during the June-September growing season (PRISM, 2018).

Table 4 Representative climatic features

Frost-free period (characteristic range)	80-100 days
Freeze-free period (characteristic range)	70-90 days
Precipitation total (characteristic range)	530-860 mm
Frost-free period (actual range)	80-100 days
Freeze-free period (actual range)	70-90 days
Precipitation total (actual range)	380-1,040 mm
Frost-free period (average)	90 days

Freeze-free period (average)	80 days
Precipitation total (average)	740 mm

Influencing water features

Due to its landscape position, this site is not influenced by wetland or riparian water features. Precipitation is the main source of water. Snow melt and seasonal ice melt contribute to a water table during the early months of the growing season.

Soil features

Soil is very deep, poorly drained, and supports permafrost. It is moderately deep (22 to 39 inches) to the restrictive permafrost layer. Parent material is comprised of mossy organic material over loamy cryoturbate.

Soil characteristics affecting vegetation include soil hydrology, soil temperature, and a thick organic layer. Soils are wet during the beginning of the growing season, due to increased inputs from snow melt and seasonal frost melt. The poorly drained soil supports a very shallow water table from March through June, which restricts vegetation. Soil temperature, which remains low in the profile throughout the year, further restricts vegetation. Low temperatures decrease decomposition rates, allowing a thick organic layer to build which serves as ideal substrate for moss ground cover. The top ten inches of soil is strongly to extremely acidic, which favors ericaceous shrubs.

Correlated soil components in MLRA 236: E36-Boreal taiga-loamy frozen till slopes

Table 5. Representative soil features

Parent material	(1) Cryoturbate
Surface texture	(1) Peat
Drainage class	Poorly drained
Permeability class	Moderate to rapid
Depth to restrictive layer	60 – 100 cm
Soil depth	150 cm
Surface fragment cover ≤3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-25.4cm)	1.27 – 4.32 cm

Soil reaction (1:1 water) (0-25.4cm)	3.5 – 10
Subsurface fragment volume <=3" (Depth not specified)	Not specified
Subsurface fragment volume >3" (Depth not specified)	Not specified

Table 6. Representative soil features (actual values)

Drainage class	Poorly drained
Permeability class	Moderate to rapid
Depth to restrictive layer	60 – 100 cm
Soil depth	150 cm
Surface fragment cover <=3"	0 %
Surface fragment cover >3"	0 %
Available water capacity (0-25.4cm)	1.27 – 4.32 cm
Soil reaction (1:1 water) (0-25.4cm)	3.5 – 10
Subsurface fragment volume <=3" (Depth not specified)	0 %
Subsurface fragment volume >3" (Depth not specified)	0 %

Ecological dynamics

This boreal site is on mountain backslopes and footslopes. Elevation ranges from 400 to 1,500 feet above sea level. Slope gradients are strong (4 to 14 percent). Soil characteristics, including hydrology and strong acidity, and a fire regime shape the vegetation.

The reference plant community is a black spruce woodland (Viereck et al., 1992). It is composed of a black spruce canopy with an open understory of ericaceous and other woody shrubs and mosses. Vegetation on this site is linked with soil characteristics. Snow and seasonal ice melt combine with a poorly drained soil and a restrictive permafrost layer to support a very shallow water table from March through June. Soil temperature remains low throughout the year, causing low rates of organic decomposition. The resulting thick organic layer is excellent substrate for black spruce and ericaceous shrubs. Additionally, the top ten inches of soil is strongly to extremely acidic, which also favors ericaceous shrubs.

Fire is a major disturbance. Fire dynamics and successional pathways in Alaskan spruce forests are complex (Viereck et al., 1992; LANDFIRE Biophysical Settings, 2009). The fire cycle and post-fire communities are shaped by factors including existing vegetation, soil characteristics, climate, and fire characteristics such as frequency and severity. Fire is responsible for three documented communities in the reference state. Post fire communities are typically comprised of extant shrubs and colonizing trees, along with fast-growing herbaceous species immediately preceding the fire event.

The information in this Ecological Dynamics section, including the state-and-transition model (STM), was developed based on current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

State and transition model

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 9. Community 1.3 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Inventory data references

Vegetative communities and transitions are described using existing models and expert knowledge. There are no vegetation inventory data points in NASIS associated with this ecological site. External data sources: The Alaska Vegetation Classification (Viereck et al., 1992) The Alaska-Yukon Region of the Circumboreal Vegetation Map (CBVM) (Jorgensen and Meidinger, 2015) LANDFIRE Biophysical Settings Models (LANDFIRE Biophysical Settings, 2009)

References

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Other references

Fryer, Janet L. 2014. *Picea mariana*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <https://www.fs.fed.us/database/feis/plants/tree/picmar/all.html>. Accessed Dec 13, 2021.

Jorgensen, T., and D. Meidinger. 2015. The Alaska Yukon Region of the Circumboreal Vegetation map (CBVM). CAFF Strategies Series Report. Conservation of Arctic Flora and Fauna, Akureyri, Iceland. ISBN: 978-9935-431-48-6.

Kautz, D.R., P. Taber, and S. Nield, editors. 2012. Land Resource Regions and Major Land Resource Areas of Alaska. United States Department of Agriculture, Natural Resources Conservation Service (USDA–NRCS).

LANDFIRE Biophysical Settings. 2009. Biophysical Setting 7616220 - Western North American Boreal Black Spruce Wet-Mesic Slope Woodland. In: LANDFIRE Biophysical Setting Model: Map zone 76, [Online]. In: Vegetation Dynamics Models. In: LANDFIRE. Washington, DC: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory; U.S.

USDA. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Contributors

Phil Barber
Steff Shoemaker
Stephanie Schmit
Jamin Johanson

Approval

Kirt Walstad, 2/13/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/26/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

14. Average percent litter cover (%) and depth (in):

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. Perennial plant reproductive capability:
-