Ecological site R236XY121AK
Boreal Tall Scrub Loamy Flood Plains

Last updated: 2/13/2024
Accessed: 11/03/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 236X–Bristol Bay-Northern Alaska Peninsula Lowlands

The Bristol Bay-Northern Alaska Peninsula Lowland Major Land Resource Area (MLRA 236) is located in Western Alaska. This MLRA covers approximately 19,500 square miles and is defined by an expanse of nearly level to rolling lowlands, uplands and low to moderate hills bordered by long, mountain footslopes. Major rivers include the Egegik, Mulchatna, Naknek, Nushagak, and Wood River. MLRA 236 is in the zone of discontinuous permafrost. It is primarily in areas with finer textured soils on terraces, rolling uplands and footslopes. This MLRA was glaciated during the early to middle Pleistocene. Moraine and glaciofluvial deposits cover around sixty percent of the MLRA. Alluvium and coastal deposits make up a large portion of the remaining area (Kautz et al., 2012; USDA, 2006). Climate patterns across this MLRA shift as one moves away from the coast. A maritime climate is prominent along the coast, while continental weather, commonly associated with Interior Alaska, is more influential inland. Across the MLRA, summers are general short and warm while winters are long and cold. Mean annual precipitation is 13 to 50 inches, with increased precipitation at higher elevations and areas away from the coast. Mean annual temperatures is between 30 and 36 degrees F (USDA, 2006). The Bristol Bay-Northern Alaska Peninsula MLRA is principally undeveloped wilderness. Federally managed land includes parts of the Katmai and Aniakchak National Parks, and the Alaska Peninsula, Becharof, Togiak and Alaska Maritime National Wildlife Refuges. The MLRA is sparsely populated. Principal communities include Dillingham, Naknek, and King Salmon. Commercial fishing in Bristol Bay and the Bering Sea comprises a major part of economic activity in the MLRA. Other land uses include subsistence activities (fishing, hunting, and gathering) and sport hunting and fishing (USDA, 2006).

Ecological site concept

This ecological site is on lowland low flood plains. Site elevation is between 20 and 810 feet above sea level. Slopes are nearly level (0 – 3 percent). Soil hydrology and post-flood ponding shape the vegetation on this landform. The reference state supports two communities. The reference plant community is characterized as a closed tall scrubland (Viereck et al., 1992). It is composed of a mix of alder and willow in the canopy with bluejoint and diverse forbs in the understory. The second community is wetter and supports more obligate wetland species.

Associated sites

F236XY111AK	**Boreal Forest Loamy Flood Plains** Both sites can be located on the same flood plain. F236XY111AK is rarely flooded with poorly drained soils. It supports a forest community in the reference state that is not present in this site.
F236XY176AK	**Boreal Woodland Loamy Flood Plains** Both sites are on flood plain talfs. F236XY176AK supports a black spruce forest on this site. Soils are more acidic, with an umbric epipedon, and support aquic conditions not found in this site.

R236XY153AK	**Boreal Willow Silty Low Flood Plains** Both sites are flood plains and can be found in the same riverine system. R236XY153AK describes low flood plains that are frequently flooded. These sites support different communities in the reference state.
R236XY156AK	**Subarctic Ericaceous Scrub Loamy Terraces** R236XY156AK described flood plain terraces. These areas do not flood and site hydrology is different than the site described herein.

Similar sites

R236XY153AK	**Boreal Willow Silty Low Flood Plains** Both sites are flood plains and can be found in the same riverine system. R236XY153AK describes low flood plains that are frequently flooded. These sites have similar reference plant communities but they are associated with different soils and unique disturbance regimes.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) *Alnus incana subsp. tenuifolia* (2) *Salix alaxensis*
Herbaceous	(1) *Calamagrostis canadensis* (2) *Impatiens noli-tangere*

Physiographic features

This site is on linear slopes of lowland flood plains.. Elevation typically ranges from 20 to 810 feet above sea level. Slopes are nearly level (0 – 3 percent). This site floods occasionally for brief periods during the growing season. This is followed by occasional, brief ponding. This site is found at all aspects.

Table 2. Representative physiographic features

Geomorphic position, flats	(1) Talf
Landforms	(1) Valley > Flood plain (2) Alluvial plain > Flood plain
Runoff class	Negligible to low
Flooding duration	Brief (2 to 7 days)
Flooding frequency	Occasional

Ponding duration	Brief (2 to 7 days)
Ponding frequency	Occasional
Elevation	10 – 240 m
Slope	0 %
Water table depth	Not specified
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Negligible to low
Flooding duration	Brief (2 to 7 days)
Flooding frequency	Occasional
Ponding duration	Brief (2 to 7 days)
Ponding frequency	Rare to occasional
Elevation	0 – 560 m
Slope	0 %
Water table depth	80 cm

Climatic features

The climate of this site reflects that of the MLRA, which is described as maritime polar (EPA, 2013). Temperatures are moderated by the nearby Bristol Bay and norther Pacific bodies of water. Annual precipitation ranges from 21 – 34 inches with approximately 40 percent occurring during the June-September growing season (PRISM, 2018).

Table 4 Representative climatic features

Frost-free period (characteristic range)	80-100 days
Freeze-free period (characteristic range)	70-90 days

Precipitation total (characteristic range)	530-860 mm
Frost-free period (actual range)	80-100 days
Freeze-free period (actual range)	70-90 days
Precipitation total (actual range)	380-1,040 mm
Frost-free period (average)	90 days
Freeze-free period (average)	80 days
Precipitation total (average)	740 mm

Influencing water features

This site is on flood plains. Flooding events are occasional and brief. Post-flood ponding is brief.

Soil features

Soils are young and weakly developed Inceptisols (Soil Survey Staff, 2013). Soils are very deep and well drained. They support a cryic temperature regime and an udic moisture regime. Parent material is grassy organic material over coarse-loamy alluvium over sandy and gravelly alluvium.

Soil hydrology influences vegetation on this site. Redox concentrations point towards anaerobic soil conditions during some point of the year. Occasional, brief ponding linked with flooding inundates this ecological site. Wet soils favor hydrophygic vegetation on this site.

Correlated soil components in MLRA 236: D36-Boreal scrub loamy flood plains; Lewispoint

Table 5. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Silt
Drainage class	Well drained
Permeability class	Moderate
Soil depth	150 cm
Surface fragment cover <=3"	Not specified

Surface fragment cover >3"	Not specified
Available water capacity (0-25.4cm)	4.06 – 4.83 cm
Soil reaction (1:1 water) (0-25.4cm)	4.7 – 5.6
Subsurface fragment volume <=3" (Depth not specified)	20 %
Subsurface fragment volume >3" (Depth not specified)	Not specified

Table 6. Representative soil features (actual values)

Drainage class	Moderately well drained to well drained
Permeability class	Moderate
Soil depth	150 cm
Surface fragment cover <=3"	0 %
Surface fragment cover >3"	0 %
Available water capacity (0-25.4cm)	4.06 – 5.84 cm
Soil reaction (1:1 water) (0-25.4cm)	4.6 – 6.1
Subsurface fragment volume <=3" (Depth not specified)	0 – 20 %

Subsurface fragment volume >3" (Depth not specified)	0 %

Ecological dynamics

This site is on lowland mid flood plains. Local site factors including local flooding dynamics and soil characteristics support two, often co-occurring plant communities. The reference plant community is a closed tall scrubland (Viereck et al., 1992). Plant species are primarily facultative upland to facultative wetland species.

Flooding is an occasional and brief disturbance on this flood plain. Flooding intensity and frequency are critical in determining the distribution and abundance of vegetation in Alaskan riverine systems (Wohl, 2007). The reference plant community develops under normal flooding circumstances. Areas that undergo post-flood ponding typically support a less diverse community. The hypoxic or anoxic condition that may result from ponding is a major abiotic stress that can determine the presence or absence of vascular plants (Vartapetian and Jackson, 1996). Wetter areas typically exclude the more hydrologically-sensitive species found in the reference plant community.

Slight to sever browsing of willows by moose occurs in the reference plant community. Browsing does not significantly affect the structure and function of this community or the ecological site.

The information in this Ecological Dynamics section, including the state-and-transition model (STM), was developed based on current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

State and transition model

Figure 7. State-and-transition model.

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)

Table 8. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)

Inventory data references

Modal points for Community 1.1 08MM13207 08AO07701 08AO20108 09AO14401 09AO13002 09AO14305 09SC00501 10SS10906 Modal points for community 1.2 09AO14002 10SS03903 10SS07204

References

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Other references

Kautz, D.R., P. Taber, and S. Nield, editors. 2012. Land Resource Regions and Major Land Resource Areas of Alaska. United States Department of Agriculture, Natural Resources Conservation Service (USDA–NRCS).

PRISM Climate Group. (PRISM) Oregon State University. https://prism.oregonstate.edu. Date created October 2018. Accessed 3 Mar 2023.

Rood, S.B., L.A. Goater, J.M. Mahoney, C.M. Pearce, and D.G. Smith. 2007. Floods, fire, and ice: Disturbance ecology of ripariancottonwoods. Canadian Journal of Botany 85(11): 1,019-1,032.

Scenarios Network for Alaska and Arctic Planning (SNAP).? Historical Monthly Temperature – 1km, 1901-2009.

http://ckan.snap.uaf.edu/dataset/. Accessed 20 Mar 2023.

Scenarios Network for Alaska and Arctic Planning (SNAP).? Historical monthly and derived precipitation products downscaled from CRU TS data via the delta methods – 2km, 1901-2009.? http://ckan.snap.uaf.edu/dataset/. Accessed 20 Mar 2023.

Soil Survey Staff. 2013. Simplified Guide to Soil Taxonomy. USDA-Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

US Environmental Protection Agency (EPA). Level III Ecoregions of the Conterminous United States. UP ESP Office of Research and Development. Corvallis, OR. http://edg.epa.gov/. Created 16 Apr 2013. Accessed 20 Mar 2023.

Vartapetian, B.B., and M.B. Jackson. 1996. Plant adaptations to anaerobic stress. Annals of Botany 79 (Supplement A): 3-20.

Wohl, E.E. 2007. Review of effects of large floods in resistant-boundary channels. In Gravel-Bed Rivers VI: From Process Understanding to River Restoration, Volume 11. H. Habersack, H. Piégay, and M. Rinaldi, editors. Elsevier Science, Amsterdam. Pages 181-211.

Yarie, J., L. Viereck, K. Van Cleve, and P. Adams. 1998. Flooding and ecosystem dynamics along the Tanana River. BioScience 48(9): 690-695.

Contributors

Phil Barber
Michael Margo
Sue Tester
Steph Schmit
Kendra Moseley
Jamin Johanson
Steff Shoemaker

Approval

Kirt Walstad, 2/13/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	11/03/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

   ---

2. **Presence of water flow patterns:**

   ---

3. **Number and height of erosional pedestals or terracettes:**

   ---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

   ---

5. **Number of gullies and erosion associated with gullies:**

   ---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

   ---

7. **Amount of litter movement (describe size and distance expected to travel):**

   ---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

   ---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

   ---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

    ---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

    ---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

    Dominant:

    Sub-dominant:

    Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth (in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**
