

# Ecological site R237XY211AK

## Western Alaska Maritime Scrubland Loamy Flood Plains

Last updated: 7/23/2020  
 Accessed: 04/22/2026

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 237X–Ahklun Mountains

The Ahklun Mountains Major Land Resource Area (MLRA 237) is in western Alaska (fig. 1). This MLRA covers approximately 14,555 square miles, and it includes the mountains, hills, and valleys of the Kilbuck Mountains in the north and the Ahklun Mountains in the south. Except for the Kilbuck Mountains and the highest ridges of the Ahklun Mountains, the MLRA was extensively glaciated during the Pleistocene (Kautz et al., 2004). Today, a few small glaciers persist in mountainous cirques (Gallant et al., 1995). The present-day landscape and landforms reflect this glacial history; glacial moraines and glacial drift cover much of the area (USDA-NRCS, 2006). The landscape of the MLRA is primarily defined by low, steep, rugged mountains cut by narrow-to-broad valleys. Flood plains and terraces of varying sizes are common at the lower elevations in the valley bottoms. Glacially carved valleys host many lakes. Togiak Lake is one of the largest lakes in the region. It is 13 miles long and about 9,500 acres in size. Major rivers include the Goodnews, Togiak, Kanektok, Osviak, Eek, and Arolik Rivers. Where the Goodnews and Togiak Rivers reach the coast, the nearly level to rolling deltas support numerous small lakes. This MLRA has two distinct climatic zones: subarctic continental and maritime continental (fig. 2). The high-elevation areas are in the subarctic continental zone. The mean annual precipitation is more than 75 inches, and the mean annual air temperature is below about 27 degrees F (-3 degrees C) in extreme locations. The warmer, drier areas at the lower elevations are in the maritime continental zone. The mean annual precipitation is 20 to 50 inches, and the mean annual air temperature is about 30 to 32 degrees F (-0.2 to 1.2 degrees C) (PRISM). This climatic zone is influenced by both maritime and continental factors. The temperatures in summer are moderated by the open waters of the Bering Sea, and the temperatures in winter are more continental due to the presence of ice in the sea (Western Regional Climate Center, 2017). The seasonal ice reaches its southernmost extent off the coast of Alaska in Bristol Bay (Alaska Climate Research Center, 2017). The western coast of Alaska is also influenced by high winds from strong storms and airmasses in the Interior Region of Alaska (Hartmann, 2002). The Ahklun Mountains MLRA is principally undeveloped wilderness. Federally managed lands include the Togiak and Alaska Maritime National Wildlife Refuges. The MLRA is sparsely populated, but it has several communities, including Togiak, Manokotak, Twin Hills, and Goodnews Bay. Togiak is the largest village. It has a population of approximately 855, most of which are Yup'ik Alaska Natives (U.S. Census Bureau, 2016). Major land uses include subsistence activities (fishing, hunting, and gathering) and wildlife recreation (USDA-NRCS, 2006; Kautz et al., 2004).

### Ecological site concept

Ecological site R237XY211AK is on mid flood plains. The reference state has one community phase. No alternate states are recorded for this ecological site. The reference plant community is open tall scrubland (Viereck et al., 1992). It consists of willows (*Salix* spp.) and an understory of hydrophilic, shade-tolerant forbs and graminoids. Bluejoint (*Calamagrostis canadensis*) and myriad forbs such as field horsetail (*Equisetum arvense*), common cowparsnip (*Heracleum maximum*), and fireweed (*Chamerion angustifolium*) are in the understory.

### Associated sites

<b>R237XY210AK</b>	<p><b>Western Alaska Maritime Scrubland Gravelly Flood Plains</b></p> <p>Ecological site R237XY211AK is on mid flood plains throughout the Ahklun Mountains area. Other ecological sites on flood plains in this area are distinguished by differences in elevation, disturbance, landform, location, associated soils, and type and amount of plants. These sites include R237XY210AK, R237XY212AK, and F237XY216AK. Site R237XY226AK is in depressions of flood plains. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut.</p>
--------------------	--

<p><b>F237XY216AK</b></p>	<p><b>Boreal Woodland Loamy Flood Plains</b></p> <p>Ecological site R237XY211AK is on mid flood plains throughout the Ahklun Mountains area. Other ecological sites on flood plains in this area are distinguished by differences in elevation, disturbance, landform, location, associated soils, and type and amount of plants. These sites include R237XY210AK, R237XY212AK, and F237XY216AK. Site R237XY226AK is in depressions of flood plains. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut.</p>
<p><b>R237XY212AK</b></p>	<p><b>Western Alaska Maritime Scrubland Silty Flood Plains</b></p> <p>Ecological site R237XY211AK is on mid flood plains throughout the Ahklun Mountains area. Other ecological sites on flood plains in this area are distinguished by differences in elevation, disturbance, landform, location, associated soils, and type and amount of plants. These sites include R237XY210AK, R237XY212AK, and F237XY216AK. Site R237XY226AK is in depressions of flood plains. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut.</p>
<p><b>R237XY226AK</b></p>	<p><b>Western Alaska Maritime Grassland Peat Flood Plains, Depression</b></p> <p>Ecological site R237XY211AK is on mid flood plains throughout the Ahklun Mountains area. Other ecological sites on flood plains in this area are distinguished by differences in elevation, disturbance, landform, location, associated soils, and type and amount of plants. These sites include R237XY210AK, R237XY212AK, and F237XY216AK. Site R237XY226AK is in depressions of flood plains. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut.</p>

**Similar sites**

<p><b>R237XY210AK</b></p>	<p><b>Western Alaska Maritime Scrubland Gravelly Flood Plains</b></p> <p>No other ecological sites in the survey area support a similar feltleaf willow community; however, several other ecological sites are on flood plains. These sites are subject to flooding and support a variety of hydrophilic plants. Differences in soil, landform, disturbance regime, and location result in distinct vegetative communities. Site R237XY210AK is on low flood plains that are subject to a different frequency and duration of flooding and are associated with different soils. Because of these differences, unique ecological sites are needed.</p>
<p><b>R237XY212AK</b></p>	<p><b>Western Alaska Maritime Scrubland Silty Flood Plains</b></p> <p>No other ecological sites in the survey area support a similar feltleaf willow community; however, several other ecological sites are on flood plains. These sites are subject to flooding and support a variety of hydrophilic plants. Differences in soil, landform, disturbance regime, and location result in distinct vegetative communities. Site R237XY212AK is in wetter areas that support more hydrophilic plant species. Because of these differences, unique ecological sites are needed.</p>
<p><b>F237XY216AK</b></p>	<p><b>Boreal Woodland Loamy Flood Plains</b></p> <p>No other ecological sites in the survey area support a similar feltleaf willow community; however, several other ecological sites are on flood plains. These sites are subject to flooding and support a variety of hydrophilic plants. Differences in soil, landform, disturbance regime, and location result in distinct vegetative communities. Site F237XY216AK is forested and is on mid to high flood plains along the Izavieknik River. Because of these differences, unique ecological sites are needed.</p>

**Table 1. Dominant plant species**

<p>Tree</p>	<p>Not specified</p>
-------------	----------------------

Shrub	(1) <i>Salix alaxensis</i>
Herbaceous	(1) <i>Calamagrostis canadensis</i> (2) <i>Equisetum arvense</i>

### Physiographic features

Site characteristics specifically relate to the reference plant community phase. Each ecological site has a specific set of site characteristics and disturbance dynamics that results in a unique plant community composition, structure, and function. Site characteristics (climate, geology, topography, and soil characteristics) are dynamic across a landscape. Subtle changes in site characteristics can result in a different plant community phase or ecological site. Definitions of site characteristics are provided in the United States Department of Agriculture Handbook 296 (USDA-NRCS, 2006), Geomorphic Description System (Schoeneberger and Wysocki, 2012), Field Book for Describing and Sampling Soils (Schoeneberger et al., 2012), and Soil Survey Manual (Soil Science Division Staff, 2017).

Figure 1. The Ahklun Mountains area (MLRA 237) is in western Alaska.

Figure 2. High-elevation and low-elevation map units in the area, which illustrate the primary climatic influence.

Table 2. Representative physiographic features

Slope shape across	(1) Linear
Slope shape up-down	(1) Linear
Geomorphic position, flats	(1) Talf
Landforms	(1) Mountain valleys or canyons > Flood plain (2) Plains > Flood plain
Flooding duration	Brief (2 to 7 days)
Flooding frequency	Occasional
Ponding frequency	None
Elevation	10 – 500 m
Slope	0 – 10 %
Water table depth	100 – 150 cm
Aspect	W, NW, N, NE, E, SE, S, SW

## Climatic features

Climate of land resource region (LLR): Maritime continental (Western Regional Climate Center, 2017); short, warm summers and long, cold winters (USDA-NRCS, 2006)

Climate of major land resource area (MLRA): Maritime continental in the lowlands and subarctic continental at higher elevations. The mean annual precipitation is 20 to 30 inches in the lowlands, and it increases to more than 45 inches at the higher elevations. The mean annual air temperature along the coast is about 34 degrees F (1 degree C) (PRISM, 2014). Strong winds are common throughout the year.

Table 3 Representative climatic features

Frost-free period (characteristic range)	80-140 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	

## Influencing water features

### Soil features

Typic Cryofluvents, occasionally flooded, are correlated to this ecological site. These soils are well drained. The saturated hydraulic conductivity is moderately low or moderately high in the stratified silt and fine sand in the upper part of the soils and high or very high in the sandy and gravelly alluvium in the lower part.

Table 4. Representative soil features

Drainage class	Well drained
----------------	--------------

## Ecological dynamics

Three major ecological sites are on flood plains throughout the Ahklun Mountains area. A relatively minor woodland ecological site is on flood plains in the corridor of the Izavieknik River. The three major ecological sites on flood plains are distinguished by landform and elevation above the associated source of floodwater. These sites are characterized as on low, mid, and high flood plains. Landform, flooding characteristics, and soil features primarily determine the vegetation supported in each ecological site.

Ecological site R237XY211AK is on mid flood plains. This site supports one plant community, which is the reference community. The site is subject to flooding, but it does not appear to shift the vegetation to a post-disturbance community.

### Disturbance Dynamics

No known disturbance regime associated with this ecological site results in an early community phase. Based on data collected, the reference plant community is resilient to flooding. Anthropogenic disturbances that remove vegetation, such as construction of trails, may exacerbate the effects of flooding or promote further disturbances such as erosion. These disturbances can alter the reference plant community and result in a different plant community; however, this was not recorded in situ.

### Flooding

Flooding is the major disturbance regime on mid flood plains. The intensity and frequency of flooding greatly affect the distribution and abundance of vegetation in Alaskan riverine systems (Wohl, 2007). Flooding can create barren, moist areas that are ideal for colonization. Floods also bury organic layers, add nutrients to the soil, deposit seed banks, and decrease competition for light and space (Rood et al., 2007; Yarie et al., 1998).

Slope and proximity to a river make these mid flood plains susceptible to occasional, brief periods of flooding, particularly during snowmelt in spring. No unique post-flood community was recorded. The soils are not subject to surface or subsurface ponding because they are well drained, and the elevation above the source of the floodwater typically prevents scouring of the existing vegetation. The wet, well drained soils are ideal for feltleaf willow and the corresponding understory species (Innes, 2014).

## Fire

Although previous wildfires have been mapped in areas of this ecological site, no evidence of fire was recorded in situ.

## Other Observations

In rare instances, balsam poplar (*Populus balsamifera*) will colonize this ecological site. Balsam poplar commonly helps to stabilize the riverbank (Harris, 1990). Once colonized, poplar likely will shade out felleaf willow, which is shade intolerant (Innes, 2014). Although the mechanism and timing by which balsam poplar colonizes and overtakes felleaf willow in this ecological site has not been recorded, it is hypothesized that flooding transports balsam poplar seeds, branches, and trees (colonization via vegetative reproduction) downstream from existing woodland ecological sites. Balsam poplar in this site was observed adjacent to or downriver from existing balsam poplar ecological sites (F237XY216AK and F237XY239AK).

Browsing by moose and caribou on willow has been documented in this ecological site. Browsing may occur during any season, but it appears to be most common in winter and summer. Browsing does not alter the vegetative community significantly; thus, an early browse sere is not required.

No alternate states are associated with this ecological site.

## State and transition model

Figure 3. State-and-transition model.

## Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

## Other references

- Alaska Climate Research Center. 2017. Climatological data—Bristol Bay. <http://oldclimate.gi.alaska.edu>. Accessed September 19, 2017.
- Gallant, A.I., E.F. Binnian, J.M. Omernik, and M.B. Shasby. 1995. Ecoregions of Alaska. U.S. Geological Survey Professional Paper 1567. Government Printing Office, Washington, D.C.
- Harris, H.T. 1990. *Populus balsamifera* subsp. *balsamifera*. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory. <https://www.fs.fed.us/database/feis/plants/tree/popbalb/all.html>. Accessed August 29, 2017.
- Hartmann, B. 2002. Climate regions of Alaska. The Alaska Climate Research Center. <http://oldclimate.gi.alaska.edu/ClimTrends/30year/regions1.html>. Modified August 28, 2002. Accessed September 19, 2017.
- Innes, R. 2014. *Salix alaxensis*. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory. <https://www.fs.fed.us/database/feis/plants/tree/salala/all.html>. Accessed August 29, 2017.
- Kautz, D.R., P. Taber, and S. Nield (editors). 2004. Land resource regions and major land resource areas of Alaska. U.S. Department of Agriculture, Natural Resources Conservation Service, Palmer, AK. Revised 2012.
- PRISM Climate Group. 2014. PRISM climate data. Oregon State University. <http://prism.oregonstate.edu>. Accessed March 27, 2018.
- Rood, S.B., L.A. Goater, J.M. Mahoney, C.M. Pearce, and D.G. Smith. 2007. Floods, fire, and ice: Disturbance ecology of riparian cottonwoods. *Canadian Journal of Botany* 85(11): 1,019-1,032.
- Schoeneberger, P.J., and D.A. Wysocki. 2012. Geomorphic description system. Version 4.2. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.
- Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. Field book for describing and sampling soils. Version 3.0. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.
- Soil Science Division Staff. 2017. Soil survey manual. Ditzler, C., K. Scheffe, and H.C Monger, editors. U.S. Department of Agriculture Handbook 18. Government Printing Office, Washington, D.C.
- U.S. Census Bureau. 2016. Vintage 2016 population estimates: Population estimates. <https://www.census.gov>. Accessed August 14, 2017.

U.S. Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Viereck, L.A., C.T. Dyrness, A.R. Batten, and K.J. Wezlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station General Technical Report PNW-GTR-286. Portland, OR.

Western Regional Climate Center. 2017. Climate of Alaska. <http://wrcc.dri.edu>. Accessed September 19, 2017.

Wohl, E.E. 2007. Review of effects of large floods in resistant-boundary channels. In Gravel-Bed Rivers VI: From Process Understanding to River Restoration, Volume 11. H. Habersack, H. Piégay, and M. Rinaldi, editors. Elsevier Science, Amsterdam. Pages 181-211.

Yarie, J., L. Viereck, K. Van Cleve, and P. Adams. 1998. Flooding and ecosystem dynamics along the Tanana River. *BioScience* 48(9): 690-695.

## Contributors

Kendra Moseley  
Michael Margo  
Stephanie Schmit  
Sue Tester  
Charlotte Crowder

## Approval

Michael Margo, 7/23/2020

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/22/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

2. Presence of water flow patterns:

---

3. Number and height of erosional pedestals or terracettes:

---

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

5. Number of gullies and erosion associated with gullies:

---

6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---