

# Ecological site R237XY224AK

## Western Alaska Maritime Scrubland Gravelly Plains, Berms

Last updated: 7/23/2020  
 Accessed: 04/22/2026

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 237X–Ahklun Mountains

The Ahklun Mountains Major Land Resource Area (MLRA 237) is in western Alaska (fig. 1). This MLRA covers approximately 14,555 square miles, and it includes the mountains, hills, and valleys of the Kilbuck Mountains in the north and the Ahklun Mountains in the south. Except for the Kilbuck Mountains and the highest ridges of the Ahklun Mountains, the MLRA was extensively glaciated during the Pleistocene (Kautz et al., 2004). Today, a few small glaciers persist in mountainous cirques (Gallant et al., 1995). The present-day landscape and landforms reflect this glacial history; glacial moraines and glacial drift cover much of the area (USDA-NRCS, 2006). The landscape of the MLRA is primarily defined by low, steep, rugged mountains cut by narrow-to-broad valleys. Flood plains and terraces of varying sizes are common at the lower elevations in the valley bottoms. Glacially carved valleys host many lakes. Togiak Lake is one of the largest lakes in the region. It is 13 miles long and about 9,500 acres in size. Major rivers include the Goodnews, Togiak, Kanektok, Osviak, Eek, and Arolik Rivers. Where the Goodnews and Togiak Rivers reach the coast, the nearly level to rolling deltas support numerous small lakes. This MLRA has two distinct climatic zones: subarctic continental and maritime continental (fig. 2). The high-elevation areas are in the subarctic continental zone. The mean annual precipitation is more than 75 inches, and the mean annual air temperature is below about 27 degrees F (-3 degrees C) in extreme locations. The warmer, drier areas at the lower elevations are in the maritime continental zone. The mean annual precipitation is 20 to 50 inches, and the mean annual air temperature is about 30 to 32 degrees F (-0.2 to 1.2 degrees C) (PRISM). This climatic zone is influenced by both maritime and continental factors. The temperatures in summer are moderated by the open waters of the Bering Sea, and the temperatures in winter are more continental due to the presence of ice in the sea (Western Regional Climate Center, 2017). The seasonal ice reaches its southernmost extent off the coast of Alaska in Bristol Bay (Alaska Climate Research Center, 2017). The western coast of Alaska is also influenced by high winds from strong storms and airmasses in the Interior Region of Alaska (Hartmann, 2002). The Ahklun Mountains MLRA is principally undeveloped wilderness. Federally managed lands include the Togiak and Alaska Maritime National Wildlife Refuges. The MLRA is sparsely populated, but it has several communities, including Togiak, Manokotak, Twin Hills, and Goodnews Bay. Togiak is the largest village. It has a population of approximately 855, most of which are Yup'ik Alaska Natives (U.S. Census Bureau, 2016). Major land uses include subsistence activities (fishing, hunting, and gathering) and wildlife recreation (USDA-NRCS, 2006; Kautz et al., 2004).

### Ecological site concept

Ecological site R237XY224AK is on the slightly sloping, leeward side of coastal berms. Landform, location, and soil characteristics combine to create this ecological site. The reference state supports a single reference community. Bulldozing of this site for construction results in an alternate state. The reference plant community is a shrub tundra consisting of low and dwarf shrub species and interspersed graminoids. Common plants include black crowberry (*Empetrum nigrum*), marsh Labrador tea (*Ledum palustre* ssp. *decumbens*), dwarf birch (*Betula nana*), bog blueberry (*Vaccinium uliginosum*), lingonberry (*Vaccinium vitis-idaea*), bluejoint (*Calamagrostis canadensis*), and smallawned sedge (*Carex microchaeta*).

### Associated sites

<b>R237XY223AK</b>	<p><b>Western Alaska Maritime Graminoid Gravelly Plains, Berms</b></p> <p>The most common associated ecological site is R237XY223AK. This site is adjacent to site R237XY224AK on the windward side of beach berms. These sites typically are differentiated by one or more criteria, including landform, landform position, associated soils, associated disturbance regimes, and the type and amount of plants. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut.</p>
--------------------	--

## Similar sites

R237XY215AK	<p><b>Western Alaska Maritime Scrubland Loamy Plains</b></p> <p>The reference plant community of several ecological sites in the Ahklun Mountains area resembles that of site R237XY224AK. Sites R237XY208AK, R237XY215AK, and R237XY218AK support a scrubland consisting of low and dwarf shrub species. Landform, elevation, and disturbance regime differentiate these sites from site R237XY224AK. Site R237XY224AK is on coastal shore complexes, and the other sites are restricted dominantly to higher elevation mountain slopes and glaciated plains.</p>
R237XY218AK	<p><b>Western Alaska Maritime Dwarf Scrubland Gravelly Slopes, Concave</b></p> <p>The reference plant community of several ecological sites in the Ahklun Mountains area resembles that of site R237XY224AK. Sites R237XY208AK, R237XY215AK, and R237XY218AK support a scrubland consisting of low and dwarf shrub species. Landform, elevation, and disturbance regime differentiate these sites from site R237XY224AK. Site R237XY224AK is on coastal shore complexes, and the other sites are restricted dominantly to higher elevation mountain slopes and glaciated plains.</p>
R237XY208AK	<p><b>Western Alaska Maritime Scrubland Peat Depressions</b></p> <p>The reference plant community of several ecological sites in the Ahklun Mountains area resembles that of site R237XY224AK. Sites R237XY208AK, R237XY215AK, and R237XY218AK support a scrubland consisting of low and dwarf shrub species. Landform, elevation, and disturbance regime differentiate these sites from site R237XY224AK. Site R237XY224AK is on coastal shore complexes, and the other sites are restricted dominantly to higher elevation mountain slopes and glaciated plains.</p>

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	<p>(1) <i>Empetrum nigrum</i>  (2) <i>Ledum palustre subsp. decumbens</i></p>
Herbaceous	<p>(1) <i>Calamagrostis canadensis</i>  (2) <i>Carex microchaeta</i></p>

## Physiographic features

Site characteristics specifically relate to the reference plant community phase. Each ecological site has a specific set of site characteristics and disturbance dynamics that results in a unique plant community composition, structure, and function. Site characteristics (climate, geology, topography, and soil characteristics) are dynamic across a landscape. Subtle changes in site characteristics can result in a different plant community phase or ecological site. Definitions of site characteristics are provided in the United States Department of Agriculture Handbook 296 (USDA-NRCS, 2006), Geomorphic Description System (Schoeneberger and Wysocki, 2012), Field Book for Describing and Sampling Soils (Schoeneberger et al., 2012), and Soil Survey Manual (Soil Science Division Staff, 2017).

Figure 1. The Ahklun Mountains area (MLRA 237) is in western Alaska.

Figure 2. High-elevation and low-elevation map units in the area, which illustrate the primary climatic influence.

**Table 2. Representative physiographic features**

Slope shape across	(1) Linear
--------------------	------------

Slope shape up-down	(1) Linear
Landforms	(1) Shore complex > Berm
Flooding frequency	Rare
Ponding frequency	None
Elevation	0 – 10 m
Slope	0 %
Water table depth	10 – 100 cm
Aspect	W, NW, N, NE, E, SE, S, SW

### Climatic features

Climate of land resource region (LLR): Maritime continental (Western Regional Climate Center, 2017); short, warm summers and long, cold winters (USDA-NRCS, 2006)

Climate of major land resource area (MLRA): Maritime continental in the lowlands and subarctic continental at higher elevations. The mean annual precipitation is 20 to 30 inches in the lowlands, and it increases to more than 45 inches at the higher elevations. The mean annual air temperature along the coast is about 34 degrees F (1 degree C) (PRISM, 2014). Strong winds are common throughout the year.

Table 3 Representative climatic features

Frost-free period (characteristic range)	90-140 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	

### Influencing water features

#### Soil features

Oxyaquic Cryorthents are correlated to this ecological site. These soils are moderately well drained and have moderately high to very high saturated hydraulic conductivity. The range in texture and stratification of these soils is a result of differences in the intensity of flooding and the energy of waves. The soils are slightly acid to very strongly acid.

Table 4. Representative soil features

Drainage class	Moderately well drained
----------------	-------------------------

## Ecological dynamics

Shore complexes consist of depressions, talfs, and rises. The depressions and talfs support two ecological sites based on whether the soils are proximal to a tidal channel (R237XY236AK) or distal (R237XY208AK). Rises are differentiated by slope length and gradient. Site R237XY224AK is on gentle slopes of wide leeward berms, and site R237XY223AK is on windward rises that have short vertical slopes and long horizontal slopes.

Site R237XY224AK is on gentle slopes of wide beach berms slightly inland from the beach. The reference state is a dense ericaceous tundra plant community. The soils associated with the site are slightly acid to very strongly acid. Acidic soils commonly are associated with ericaceous shrub communities (Viereck et al., 1992). The low pH and natural drainage class of the soils, dense vegetation, and allelopathic effects associated with the *Empetrum* genus appear to prevent colonization by herbaceous plants and larger shrubs (Bråthen et al., 2010; Swanson, 2015). These factors and lack of a pronounced disturbance regime result in a resilient reference state that does not exhibit an early community phase. Natural variations in plant richness and cover may be evident among areas of this site.

### Disturbance Dynamics

No known disturbance regime in this ecological site results in an early community phase. These wide, gently sloping, commonly leeward beach berms generally are protected from coastal flooding and salt spray by the berms and vegetative communities of site R237XY223AK. Although this site may be subject to strong winds, this disturbance does not result in the site surpassing the resilience threshold required to create an early sere. Anthropogenic disturbances that remove vegetation, such as construction of trails, may promote further disturbances such as erosion that can alter the reference plant community and result in a separate community.

### Flooding

This site is subject to rare (1 to 5 times every 100 years) periods of flooding; however, the flooding does not appear to affect the vegetation. A water table is near the surface for a short time in April as the ground thaws and seasonal frost retreats. This may affect the plants that can grow in this site, but it does not create an early sere.

### Fire

No incidence or evidence of fire was recorded in situ for this site, but previous wildfires have been mapped in areas of the site. Historically, the main causes of wildfires in the Ahklun Mountains area are lightning strikes and human activity (AICC, 2017).

### Other Observations

No grazing or browsing was observed in this ecological site. This likely is due to the proximity of the site to coastal villages rather than a lack of edible plants.

One alternate state is in this ecological site. Villages such as Togiak were built on berms. Bulldozing and construction alter the soil surface and lead to a distinct alternate state plant community.

## State and transition model

Figure 3. State-and-transition model.

## Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

Table 6. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

## Other references

Alaska Climate Research Center. 2017. Climatological data—Bristol Bay. <http://oldclimate.gi.alaska.edu>. Accessed September 19, 2017.

Alaska Interagency Coordination Center (AICC). <https://fire.ak.blm.gov/predsvcs/maps.php>. Accessed August 16, 2017.

Bråthen, K.A., C.H. Fodstad, and C. Gallet. 2010. Ecosystem disturbance reduces the allelopathic effects of *Empetrum hermaphroditum* humus on tundra plants. *Journal of Vegetation Science* 21: 786-795.

Gallant, A.I., E.F. Binnian, J.M. Omernik, and M.B. Shasby. 1995. Ecoregions of Alaska. U.S. Geological Survey Professional Paper 1567. Government Printing Office, Washington, D.C.

Hartmann, B. 2002. Climate regions of Alaska. The Alaska Climate Research Center. <http://oldclimate.gi.alaska.edu/ClimTrends/30year/regions1.html>. Modified August 28, 2002. Accessed September 19, 2017.

Kautz, D.R., P. Taber, and S. Nield (editors). 2004. Land resource regions and major land resource areas of Alaska. U.S. Department of Agriculture, Natural Resources Conservation Service, Palmer, AK. Revised 2012.

PRISM Climate Group. 2014. PRISM climate data. Oregon State University. <http://prism.oregonstate.edu>. Accessed March 27, 2018.

Schoeneberger, P.J., and D.A. Wysocki. 2012. Geomorphic description system. Version 4.2. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. Field book for describing and sampling soils. Version 3.0. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Soil Science Division Staff. 2017. Soil survey manual. Ditzler, C., K. Scheffe, and H.C Monger, editors. U.S. Department of Agriculture Handbook 18. Government Printing Office, Washington, D.C.

Swanson, D.K. 2015. Environmental limits of tall shrubs in Alaska's arctic national parks. Eryuan Liang, editor. PLoS One. 10(9): e0138387.

U.S. Census Bureau. 2016. Vintage 2016 population estimates: Population estimates. <https://www.census.gov>. Accessed August 14, 2017.

U.S. Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Viereck, L.A., C.T. Dyrness, A.R. Batten, and K.J. Wezlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station General Technical Report PNW-GTR-286. Portland, OR.

Western Regional Climate Center. 2017. Climate of Alaska. <http://wrcc.dri.edu>. Accessed September 19, 2017.

## Contributors

Kendra Moseley  
Michael Margo  
Stephanie Schmit  
Sue Tester  
Charlotte Crowder

## Approval

Michael Margo, 7/23/2020

## Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/22/2026
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. Number and extent of rills:

---

2. Presence of water flow patterns:

---

3. Number and height of erosional pedestals or terracettes:

---

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

5. Number of gullies and erosion associated with gullies:

---

6. Extent of wind scoured, blowouts and/or depositional areas:

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

**Dominant:**

**Sub-dominant:**

**Other:**

**Additional:**

---

**13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

**14. Average percent litter cover (%) and depth ( in):**

---

**15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

**16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

**17. Perennial plant reproductive capability:**

---