

# Ecological site R237XY226AK

## Western Alaska Maritime Grassland Peat Flood Plains, Depression

Last updated: 7/23/2020

Accessed: 04/22/2026

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 237X–Ahklun Mountains

The Ahklun Mountains Major Land Resource Area (MLRA 237) is in western Alaska (fig. 3). This MLRA covers approximately 14,555 square miles, and it includes the mountains, hills, and valleys of the Kilbuck Mountains in the north and the Ahklun Mountains in the south. Except for the Kilbuck Mountains and the highest ridges of the Ahklun Mountains, the MLRA was extensively glaciated during the Pleistocene (Kautz et al., 2004). Today, a few small glaciers persist in mountainous cirques (Gallant et al., 1995). The present-day landscape and landforms reflect this glacial history; glacial moraines and glacial drift cover much of the area (USDA-NRCS, 2006). The landscape of the MLRA is primarily defined by low, steep, rugged mountains cut by narrow-to-broad valleys. Flood plains and terraces of varying sizes are common at the lower elevations in the valley bottoms. Glacially carved valleys host many lakes. Togiak Lake is one of the largest lakes in the region. It is 13 miles long and about 9,500 acres in size. Major rivers include the Goodnews, Togiak, Kanektok, Osviak, Eek, and Arolik Rivers. Where the Goodnews and Togiak Rivers reach the coast, the nearly level to rolling deltas support numerous small lakes. This MLRA has two distinct climatic zones: subarctic continental and maritime continental (fig. 4). The high-elevation areas are in the subarctic continental zone. The mean annual precipitation is more than 75 inches, and the mean annual air temperature is below about 27 degrees F (-3 degrees C) in extreme locations. The warmer, drier areas at the lower elevations are in the maritime continental zone. The mean annual precipitation is 20 to 50 inches, and the mean annual air temperature is about 30 to 32 degrees F (-0.2 to 1.2 degrees C) (PRISM). This climatic zone is influenced by both maritime and continental factors. The temperatures in summer are moderated by the open waters of the Bering Sea, and the temperatures in winter are more continental due to the presence of ice in the sea (Western Regional Climate Center, 2017). The seasonal ice reaches its southernmost extent off the coast of Alaska in Bristol Bay (Alaska Climate Research Center, 2017). The western coast of Alaska is also influenced by high winds from strong storms and airmasses in the Interior Region of Alaska (Hartmann, 2002). The Ahklun Mountains MLRA is principally undeveloped wilderness. Federally managed lands include the Togiak and Alaska Maritime National Wildlife Refuges. The MLRA is sparsely populated, but it has several communities, including Togiak, Manokotak, Twin Hills, and Goodnews Bay. Togiak is the largest village. It has a population of approximately 855, most of which are Yup'ik Alaska Natives (U.S. Census Bureau, 2016). Major land uses include subsistence activities (fishing, hunting, and gathering) and wildlife recreation (USDA-NRCS, 2006; Kautz et al., 2004).

### Ecological site concept

Ecological site R237XY226AK is in organic soil depressions of lowland flood plains. The reference state supports a single community, which is the reference plant community. Ponding and flooding help to sustain the reference plant community. An alternate state is a result of beaver activity. The reference plant community is a wet sedge-herb meadow tundra (Viereck et al., 1992). Common obligate wetland graminoids are water sedge (*Carex aquatilis*) and beaked sedge (*C. rostrata*), and common hydrophilic forbs are water horsetail (*Equisetum fluviatile*) and purple marshlocks (*Comarum palustre*). Other common species include arctic sweet coltsfoot (*Petasites frigidus*) and western water hemlock (*Cicuta douglasii*).

### Associated sites

<b>R237XY210AK</b>	<p><b>Western Alaska Maritime Scrubland Gravelly Flood Plains</b></p> <p>Ecological site R237XY226AK is on flood plains. Associated sites on flood plains include R237XY210AK, R237XY211AK, and R237XY212AK. Site R237XY226AK is differentiated from these sites by landform, slope shape, soils, and disturbances. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut. This is especially noticeable in smaller depressions because the edge effects are relatively large or make up the total area and species from surrounding ecological sites may intermix.</p>
<b>R237XY211AK</b>	<p><b>Western Alaska Maritime Scrubland Loamy Flood Plains</b></p> <p>Ecological site R237XY226AK is on flood plains. Associated sites on flood plains include R237XY210AK, R237XY211AK, and R237XY212AK. Site R237XY226AK is differentiated from these sites by landform, slope shape, soils, and disturbances. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut. This is especially noticeable in smaller depressions because the edge effects are relatively large or make up the total area and species from surrounding ecological sites may intermix.</p>
<b>R237XY212AK</b>	<p><b>Western Alaska Maritime Scrubland Silty Flood Plains</b></p> <p>Ecological site R237XY226AK is on flood plains. Associated sites on flood plains include R237XY210AK, R237XY211AK, and R237XY212AK. Site R237XY226AK is differentiated from these sites by landform, slope shape, soils, and disturbances. Ecotonal plant communities that have characteristics from more than one ecological site are in areas where these sites abut. This is especially noticeable in smaller depressions because the edge effects are relatively large or make up the total area and species from surrounding ecological sites may intermix.</p>

**Similar sites**

<b>R237XY205AK</b>	<p><b>Western Alaska Maritime Scrubland Loamy Swales</b></p> <p>Although site R237XY205AK is similar to site R237XY226AK, site R237XY205AK is in swales that are not subject to ponding and are not susceptible to beaver damming. Site R237XY205AK supports willows and bluejoint in the reference plant community, which are not in site R237XY226AK. Differences in soils and disturbance regimes lead to different reference state communities.</p>
<b>R237XY208AK</b>	<p><b>Western Alaska Maritime Scrubland Peat Depressions</b></p> <p>Sites R237XY226AK and R237XY208AK are in depressions, and both support a variety of hydrophilic plant species. Site R237XY226AK is in mineral soil depressions of flood plains, and site R237XY208AK is in organic depressions of low- and high-elevation glaciated plains. Differences in soils and disturbance regimes lead to different reference state communities.</p>

Figure 1. Willows encroach on the reference plant community in some areas.

Figure 2. View of alternate beaver-affected state. Multiple beaver dams and lodges are in the large depressions. Beaver dams are at far left and just out of view on right.

Table 3. Dominant plant species

Tree	Not specified
Shrub	Not specified

Herbaceous	(1) <i>Carex</i> (2) <i>Comarum palustre</i>
------------	---

### Physiographic features

Site characteristics specifically relate to the reference plant community phase. Each ecological site has a specific set of site characteristics and disturbance dynamics that results in a unique plant community composition, structure, and function. Site characteristics (climate, geology, topography, and soil characteristics) are dynamic across a landscape. Subtle changes in site characteristics can result in a different plant community phase or ecological site. Definitions of site characteristics are provided in the United States Department of Agriculture Handbook 296 (USDA-NRCS, 2006), Geomorphic Description System (Schoeneberger and Wysocki, 2012), Field Book for Describing and Sampling Soils (Schoeneberger et al., 2012), and Soil Survey Manual (Soil Science Division Staff, 2017).

Figure 1. The Ahklun Mountains area (MLRA 237) is in western Alaska.

Figure 2. High-elevation and low-elevation map units in the area, which illustrate the primary climatic influence.

Table 4. Representative physiographic features

Slope shape up-down	(1) Concave
Slope shape across	(1) Concave
Landforms	(1) Lowland > Flood plain
Flooding frequency	Rare
Ponding duration	Very long (more than 30 days)
Ponding frequency	Frequent
Elevation	10 – 670 m
Aspect	W, NW, N, NE, E, SE, S, SW

### Climatic features

Climate of land resource region (LLR): Maritime continental (Western Regional Climate Center, 2017); short, warm summers and long, cold winters (USDA-NRCS, 2006)

Climate of major land resource area (MLRA): Maritime continental in the lowlands and subarctic continental at higher elevations. The mean annual precipitation is 20 to 30 inches in the lowlands, and it increases to more than 45 inches at the higher elevations. The mean annual air temperature along the coast is about 34 degrees F (1 degree C) (PRISM, 2014). Strong winds are common throughout the year.

Table 5 Representative climatic features

Frost-free period (characteristic range)	80-130 days
--	-------------

Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	

## Influencing water features

### Soil features

Histosols are correlated to this ecological site. The saturated hydraulic conductivity of the soils is moderately high in the upper 12 inches and very high below that depth. These soils are very poorly drained.

Table 6. Representative soil features

Drainage class	Very poorly drained
----------------	---------------------

### Ecological dynamics

Flood plains are throughout the Ahklun Mountains area. The ecological sites on the flood plains differ in location, landform, elevation above the flood source, soil characteristics, and disturbance regime. Flood plain depressions that experience very long ponding support a unique reference state that is differentiated by soil characteristics, disturbance regime, and landform.

Site R237XY226AK is in organic soil depressions of flood plains. These depressions are wet and have a year-round water table at the soil surface, limiting the plants that can grow. The soil associated with this site is very poorly drained. The reference state supports one stable reference plant community. The site is subject to ponding and flooding. Plants are primarily obligate wetland species that are tolerant of ponding and low-energy flooding. Obligate wetland species thrive in wet soils where the high content of soil moisture excludes less hydrophilic plant species. Willow may encroach on this site (fig. 1), but it is too wet to support a willow scrubland community. More willow could colonize this site if it dried, but this was not recorded in situ.

#### Disturbance Dynamics

No disturbance regime in this ecological site results in an early community phase.

#### Ponding

Depressions of the flood plains are wet. The soils are very poorly drained, and a water table is at the soil surface throughout the year. Frequent, very long periods (more than 30 days) of ponding occur during the growing season. Ponding affects the plant community by inhibiting oxygen to susceptible plants (Hook and Crawford, 1978; Jackson et al., 1991). Hypoxic and anoxic conditions are major stressors and partially determine the presence or absence of vascular plants. The temporal tolerance to oxygen deprivation differs among plant species, ranging from many hours to several weeks (Vartapetian and Jackson, 1996). The reference plant community dominantly supports obligate wetland species, which are resilient to ponding. Ponding does not result in an early community sere.

#### Flooding

This site is subject to periodic flooding. The frequency and duration of flooding typically coincide with the local flooding regime of the surrounding flood plains. Flooding does not cause the site to surpass the resilience threshold required to create an early sere.

#### Fire

No incidence or evidence of fire was recorded in situ for this ecological site; however, previous wildfires have been mapped in areas of this site. Historically, the major causes of wildfires in the Ahklun Mountains area are lightning strikes and human activity (AICC, 2017).

#### Other Observations

No browsing or grazing is evident in the reference state,

One alternate state created by beaver activity is recognized in this ecological site. Beaver activity, particularly dam construction, can restructure the plant community in the depressions. Beavers are an introduced species in the Ahklun Mountains area. The extent of their effect on this site has not been fully researched.

## State and transition model

### Additional community tables

**Table 7. Community 1.1 plant community composition**

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

**Table 8. Community 2.1 plant community composition**

Group	Common Name	Symbol	Scientific Name	Annual Production ()	Foliar Cover (%)
-------	-------------	--------	-----------------	----------------------	------------------

## Other references

- Alaska Climate Research Center. 2017. Climatological data–Bristol Bay. <http://oldclimate.gi.alaska.edu>. Accessed September 19, 2017.
- Alaska Interagency Coordination Center (AICC). <https://fire.ak.blm.gov/predsvcs/maps.php>. Accessed August 16, 2017.
- Gallant, A.I., E.F. Binnian, J.M. Omernik, and M.B. Shasby. 1995. Ecoregions of Alaska. U.S. Geological Survey Professional Paper 1567. Government Printing Office, Washington, D.C.
- Hartmann, B. 2002. Climate regions of Alaska. The Alaska Climate Research Center. <http://oldclimate.gi.alaska.edu/ClimTrends/30year/regions1.html>. Modified August 28, 2002. Accessed September 19, 2017.
- Hook, D., and R.M.M. Crawford. 1978. Plant life in anaerobic environments. Ann Arbor Science Publishers, Ann Arbor, MI.
- Jackson, M.B., D.D. Davies, and H. Lambers (editors). 1991. Plant life under oxygen deprivation: Ecology, physiology, and biochemistry. SPB Academic Publication, The Hague, Netherlands.
- Kautz, D.R., P. Taber, and S. Nield (editors). 2004. Land resource regions and major land resource areas of Alaska. U.S. Department of Agriculture, Natural Resources Conservation Service, Palmer, AK. Revised 2012.
- PRISM Climate Group. 2014. PRISM climate data. Oregon State University. <http://prism.oregonstate.edu>. Accessed March 27, 2018.
- Schoeneberger, P.J., and D.A. Wysocki. 2012. Geomorphic description system. Version 4.2. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.
- Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. Field book for describing and sampling soils. Version 3.0. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.
- Soil Science Division Staff. 2017. Soil survey manual. Ditzler, C., K. Scheffe, and H.C Monger, editors. U.S. Department of Agriculture Handbook 18. Government Printing Office, Washington, D.C.
- U.S. Census Bureau. 2016. Vintage 2016 population estimates: Population estimates. <https://www.census.gov>. Accessed August 14, 2017.
- U.S. Department of Agriculture, Forest Service, Alaska Region. 2013. Forest health conditions in Alaska - 2012. Publication R10-PR-32.
- U.S. Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.
- Vartapetian, B.B., and M.B. Jackson. 1996. Plant adaptations to anaerobic stress. *Annals of Botany* 79 (Supplement A): 3-20.
- Viereck, L.A., C.T. Dyrness, A.R. Batten, and K.J. Wezlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station General Technical Report PNW-GTR-286. Portland, OR.
- Western Regional Climate Center. 2017. Climate of Alaska. <http://wrcc.dri.edu>. Accessed September 19, 2017.
- Wolf, E.C., D.J. Cooper, and N.T. Hobbs. 2007. Hydrological regime and herbivory stabilize an alternative state in Yellowstone National Park. *Ecological Applications* 17(6): 1572-1587.

## Contributors

Kendra Moseley  
Michael Margo  
Stephanie Schmit  
Sue Tester  
Charlotte Crowder

## Approval

Michael Margo, 7/23/2020

### Rangeland health reference sheet

**Interpreting Indicators of Rangeland Health** is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/22/2026
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

**1. Number and extent of rills:**

---

**2. Presence of water flow patterns:**

---

**3. Number and height of erosional pedestals or terracettes:**

---

**4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

**5. Number of gullies and erosion associated with gullies:**

---

**6. Extent of wind scoured, blowouts and/or depositional areas:**

---

7. Amount of litter movement (describe size and distance expected to travel):

---

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

---

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

---

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

---

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

---

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---